

Table 7.117A Asthma in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | ASTHMA IN LIFETIME | | | ASTHMA IN PAST YEAR | | |
|---------------|--------------------|--|--------|---------------------|--|--------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 26,946 | 2,834 | 24,111 | 15,293 | 1,478 | 13,815 |
| 12-17 | 4,088 | 322 | 3,766 | 2,454 | 229 | 2,224 |
| 18-25 | 4,415 | 1,038 | 3,377 | 2,231 | 498 | 1,733 |
| 26-34 | 3,947 | 531 | 3,416 | 2,097 | 275 | 1,822 |
| 35-49 | 6,536 | 595 | 5,942 | 3,521 | 339 | 3,182 |
| 50-64 | 4,515 | 255 | 4,260 | 2,764 | 91 | 2,672 |
| 65 or Older | 3,444 | * | 3,351 | 2,227 | * | 2,182 |
| MALE | 11,196 | 1,599 | 9,597 | 5,660 | 781 | 4,879 |
| 12-17 | 2,188 | 159 | 2,029 | 1,220 | 112 | 1,108 |
| 18-25 | 2,118 | 608 | 1,509 | 968 | 270 | 698 |
| 26-34 | 1,762 | 364 | 1,398 | 779 | 177 | 602 |
| 35-49 | 2,361 | 259 | 2,102 | 1,003 | 103 | 899 |
| 50-64 | 1,650 | 164 | 1,485 | 946 | 73 | 873 |
| 65 or Older | 1,117 | * | 1,073 | 743 | * | 699 |
| FEMALE | 15,750 | 1,235 | 14,515 | 9,633 | 697 | 8,936 |
| 12-17 | 1,900 | 163 | 1,737 | 1,234 | 117 | 1,116 |
| 18-25 | 2,297 | 430 | 1,867 | 1,263 | 228 | 1,036 |
| 26-34 | 2,185 | 167 | 2,018 | 1,318 | 98 | 1,220 |
| 35-49 | 4,176 | 336 | 3,840 | 2,518 | 235 | 2,283 |
| 50-64 | 2,865 | * | 2,774 | 1,817 | * | 1,799 |
| 65 or Older | 2,327 | * | 2,278 | 1,483 | * | 1,483 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.117B Asthma in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | ASTHMA IN LIFETIME | | | ASTHMA IN PAST YEAR | | |
|---------------|--------------------|--|------|---------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 11.2 | 13.0 | 11.1 | 6.4 | 6.8 | 6.3 |
| 12-17 | 16.9 | 16.6 | 16.9 | 10.2 | 11.8 | 10.0 |
| 18-25 | 13.9 | 15.0 | 13.6 | 7.0 | 7.2 | 7.0 |
| 26-34 | 11.5 | 12.6 | 11.4 | 6.1 | 6.5 | 6.1 |
| 35-49 | 10.1 | 10.5 | 10.1 | 5.5 | 6.0 | 5.4 |
| 50-64 | 9.1 | 10.7 | 9.0 | 5.6 | 3.8 | 5.6 |
| 65 or Older | 9.8 | * | 9.7 | 6.4 | * | 6.3 |
| MALE | 9.7 | 11.5 | 9.4 | 4.9 | 5.6 | 4.8 |
| 12-17 | 17.9 | 16.9 | 18.0 | 10.0 | 12.0 | 9.9 |
| 18-25 | 13.3 | 14.3 | 13.0 | 6.1 | 6.3 | 6.0 |
| 26-34 | 10.4 | 12.8 | 9.9 | 4.6 | 6.2 | 4.3 |
| 35-49 | 7.5 | 7.1 | 7.5 | 3.2 | 2.8 | 3.2 |
| 50-64 | 6.9 | 9.6 | 6.6 | 3.9 | 4.3 | 3.9 |
| 65 or Older | 7.5 | * | 7.4 | 5.0 | * | 4.8 |
| FEMALE | 12.7 | 15.6 | 12.5 | 7.8 | 8.8 | 7.7 |
| 12-17 | 15.8 | 16.2 | 15.8 | 10.3 | 11.7 | 10.2 |
| 18-25 | 14.4 | 16.1 | 14.1 | 7.9 | 8.5 | 7.8 |
| 26-34 | 12.6 | 12.2 | 12.6 | 7.6 | 7.2 | 7.6 |
| 35-49 | 12.7 | 16.6 | 12.4 | 7.7 | 11.6 | 7.4 |
| 50-64 | 11.2 | * | 11.1 | 7.1 | * | 7.2 |
| 65 or Older | 11.6 | * | 11.4 | 7.4 | * | 7.5 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.118A Bronchitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | BRONCHITIS IN LIFETIME | | | BRONCHITIS IN PAST YEAR | | |
|---------------|------------------------|--|--------|-------------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 26,577 | 2,487 | 24,090 | 10,468 | 1,020 | 9,448 |
| 12-17 | 2,019 | 159 | 1,860 | 884 | 84 | 800 |
| 18-25 | 3,067 | 815 | 2,252 | 1,116 | 344 | 773 |
| 26-34 | 3,511 | 412 | 3,099 | 1,225 | 131 | 1,094 |
| 35-49 | 7,377 | 559 | 6,818 | 2,579 | 179 | 2,400 |
| 50-64 | 6,878 | 426 | 6,452 | 3,001 | 196 | 2,806 |
| 65 or Older | 3,725 | * | 3,608 | 1,662 | * | 1,576 |
| MALE | 9,719 | 1,363 | 8,356 | 3,690 | 589 | 3,101 |
| 12-17 | 977 | 66 | 910 | 449 | 34 | 414 |
| 18-25 | 1,328 | 431 | 897 | 421 | 164 | 257 |
| 26-34 | 1,433 | 245 | 1,187 | 508 | 80 | 427 |
| 35-49 | 2,486 | 302 | 2,184 | 776 | 96 | 679 |
| 50-64 | 2,383 | 251 | 2,133 | 1,011 | 167 | 844 |
| 65 or Older | 1,112 | * | 1,044 | 527 | * | 479 |
| FEMALE | 16,858 | 1,125 | 15,734 | 6,778 | 431 | 6,347 |
| 12-17 | 1,042 | 92 | 950 | 436 | 50 | 386 |
| 18-25 | 1,740 | 385 | 1,355 | 696 | 180 | 516 |
| 26-34 | 2,078 | 166 | 1,912 | 717 | 51 | 667 |
| 35-49 | 4,891 | 257 | 4,634 | 1,803 | 82 | 1,721 |
| 50-64 | 4,494 | * | 4,320 | 1,991 | * | 1,962 |
| 65 or Older | 2,613 | * | 2,564 | 1,136 | * | 1,097 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.118B Bronchitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | BRONCHITIS IN LIFETIME | | | BRONCHITIS IN PAST YEAR | | |
|---------------|------------------------|--|------|-------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 11.1 | 11.4 | 11.1 | 4.4 | 4.7 | 4.3 |
| 12-17 | 8.3 | 8.2 | 8.4 | 3.7 | 4.3 | 3.6 |
| 18-25 | 9.6 | 11.8 | 9.1 | 3.5 | 5.0 | 3.1 |
| 26-34 | 10.2 | 9.7 | 10.3 | 3.6 | 3.1 | 3.6 |
| 35-49 | 11.4 | 9.8 | 11.6 | 4.0 | 3.2 | 4.1 |
| 50-64 | 13.8 | 17.8 | 13.6 | 6.0 | 8.2 | 5.9 |
| 65 or Older | 10.6 | * | 10.5 | 4.8 | * | 4.6 |
| MALE | 8.4 | 9.8 | 8.2 | 3.2 | 4.2 | 3.0 |
| 12-17 | 8.0 | 7.1 | 8.1 | 3.7 | 3.7 | 3.7 |
| 18-25 | 8.4 | 10.1 | 7.7 | 2.6 | 3.8 | 2.2 |
| 26-34 | 8.5 | 8.6 | 8.4 | 3.0 | 2.8 | 3.0 |
| 35-49 | 7.9 | 8.3 | 7.8 | 2.5 | 2.6 | 2.4 |
| 50-64 | 9.9 | 14.7 | 9.5 | 4.2 | 9.8 | 3.8 |
| 65 or Older | 7.4 | * | 7.2 | 3.5 | * | 3.3 |
| FEMALE | 13.6 | 14.2 | 13.6 | 5.5 | 5.4 | 5.5 |
| 12-17 | 8.7 | 9.2 | 8.6 | 3.6 | 5.0 | 3.5 |
| 18-25 | 10.9 | 14.4 | 10.2 | 4.4 | 6.7 | 3.9 |
| 26-34 | 12.0 | 12.2 | 12.0 | 4.1 | 3.7 | 4.2 |
| 35-49 | 14.9 | 12.7 | 15.0 | 5.5 | 4.1 | 5.6 |
| 50-64 | 17.5 | * | 17.3 | 7.8 | * | 7.9 |
| 65 or Older | 13.0 | * | 12.9 | 5.7 | * | 5.5 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.119A Diabetes in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | DIABETES IN LIFETIME | | | DIABETES IN PAST YEAR | | |
|---------------|----------------------|--|--------|-----------------------|--|--------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 16,697 | 533 | 16,164 | 14,118 | 433 | 13,684 |
| 12-17 | 143 | 11 | 131 | 85 | 9 | 76 |
| 18-25 | 325 | 70 | 256 | 241 | 54 | 188 |
| 26-34 | 805 | 108 | 697 | 604 | 76 | 528 |
| 35-49 | 3,053 | 197 | 2,856 | 2,529 | 185 | 2,343 |
| 50-64 | 6,635 | 115 | 6,521 | 5,694 | 88 | 5,606 |
| 65 or Older | 5,735 | * | 5,703 | 4,965 | * | 4,943 |
| MALE | 7,587 | 326 | 7,261 | 6,475 | 295 | 6,180 |
| 12-17 | 59 | 8 | 51 | 29 | 8 | 21 |
| 18-25 | 142 | 45 | 97 | 103 | 36 | 67 |
| 26-34 | 286 | 44 | 242 | 228 | 27 | 201 |
| 35-49 | 1,531 | 168 | 1,363 | 1,280 | 164 | 1,117 |
| 50-64 | 3,147 | 56 | 3,091 | 2,721 | 56 | 2,665 |
| 65 or Older | 2,422 | * | 2,417 | 2,114 | * | 2,109 |
| FEMALE | 9,110 | 207 | 8,903 | 7,643 | 138 | 7,505 |
| 12-17 | 84 | 3 | 81 | 56 | 1 | 55 |
| 18-25 | 184 | 25 | 159 | 139 | 18 | 121 |
| 26-34 | 519 | 64 | 455 | 376 | 49 | 327 |
| 35-49 | 1,523 | 29 | 1,493 | 1,248 | 21 | 1,227 |
| 50-64 | 3,488 | * | 3,429 | 2,973 | * | 2,941 |
| 65 or Older | 3,312 | * | 3,286 | 2,851 | * | 2,834 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.119B Diabetes in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | DIABETES IN LIFETIME | | | DIABETES IN PAST YEAR | | |
|---------------|----------------------|--|------|-----------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 7.0 | 2.4 | 7.4 | 5.9 | 2.0 | 6.3 |
| 12-17 | 0.6 | 0.6 | 0.6 | 0.4 | 0.5 | 0.3 |
| 18-25 | 1.0 | 1.0 | 1.0 | 0.8 | 0.8 | 0.8 |
| 26-34 | 2.3 | 2.6 | 2.3 | 1.8 | 1.8 | 1.8 |
| 35-49 | 4.7 | 3.5 | 4.9 | 3.9 | 3.3 | 4.0 |
| 50-64 | 13.4 | 4.8 | 13.8 | 11.5 | 3.7 | 11.9 |
| 65 or Older | 16.4 | * | 16.6 | 14.2 | * | 14.4 |
| MALE | 6.6 | 2.3 | 7.1 | 5.6 | 2.1 | 6.1 |
| 12-17 | 0.5 | 0.8 | 0.5 | 0.2 | 0.8 | 0.2 |
| 18-25 | 0.9 | 1.1 | 0.8 | 0.6 | 0.8 | 0.6 |
| 26-34 | 1.7 | 1.5 | 1.7 | 1.3 | 0.9 | 1.4 |
| 35-49 | 4.8 | 4.6 | 4.9 | 4.0 | 4.5 | 4.0 |
| 50-64 | 13.1 | 3.3 | 13.8 | 11.3 | 3.3 | 11.9 |
| 65 or Older | 16.2 | * | 16.7 | 14.2 | * | 14.6 |
| FEMALE | 7.4 | 2.6 | 7.7 | 6.2 | 1.7 | 6.5 |
| 12-17 | 0.7 | 0.3 | 0.7 | 0.5 | 0.1 | 0.5 |
| 18-25 | 1.2 | 0.9 | 1.2 | 0.9 | 0.7 | 0.9 |
| 26-34 | 3.0 | 4.7 | 2.8 | 2.2 | 3.6 | 2.0 |
| 35-49 | 4.6 | 1.5 | 4.8 | 3.8 | 1.1 | 4.0 |
| 50-64 | 13.6 | * | 13.7 | 11.6 | * | 11.8 |
| 65 or Older | 16.5 | * | 16.5 | 14.2 | * | 14.2 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.120A Heart Disease in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | HEART DISEASE IN LIFETIME | | | HEART DISEASE IN PAST YEAR | | |
|---------------|---------------------------|--|--------|----------------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 13,450 | 571 | 12,878 | 9,554 | 328 | 9,226 |
| 12-17 | 97 | 8 | 89 | 43 | 8 | 36 |
| 18-25 | 167 | 36 | 131 | 74 | 13 | 61 |
| 26-34 | 279 | 85 | 194 | 181 | 51 | 130 |
| 35-49 | 1,350 | 130 | 1,220 | 921 | 78 | 843 |
| 50-64 | 4,358 | 265 | 4,093 | 2,943 | 162 | 2,782 |
| 65 or Older | 7,199 | * | 7,151 | 5,392 | * | 5,375 |
| MALE | 7,488 | 392 | 7,096 | 5,120 | 224 | 4,896 |
| 12-17 | 47 | 5 | 43 | 24 | 5 | 20 |
| 18-25 | 60 | 13 | 48 | 20 | 4 | 17 |
| 26-34 | 103 | 17 | 85 | 53 | 5 | 48 |
| 35-49 | 711 | 74 | 637 | 476 | 45 | 431 |
| 50-64 | 2,714 | 236 | 2,478 | 1,727 | 149 | 1,579 |
| 65 or Older | 3,853 | * | 3,805 | 2,819 | * | 2,802 |
| FEMALE | 5,961 | 179 | 5,783 | 4,434 | 104 | 4,330 |
| 12-17 | 49 | 3 | 46 | 19 | 3 | 16 |
| 18-25 | 106 | 24 | 83 | 53 | 9 | 44 |
| 26-34 | 176 | 67 | 109 | 128 | * | 82 |
| 35-49 | 639 | 56 | 583 | 444 | 32 | 412 |
| 50-64 | 1,644 | * | 1,615 | 1,216 | * | 1,203 |
| 65 or Older | 3,346 | * | 3,346 | 2,573 | * | 2,573 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.120B Heart Disease in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | HEART DISEASE IN LIFETIME | | | HEART DISEASE IN PAST YEAR | | |
|---------------|---------------------------|--|------|----------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 5.6 | 2.6 | 5.9 | 4.0 | 1.5 | 4.2 |
| 12-17 | 0.4 | 0.4 | 0.4 | 0.2 | 0.4 | 0.2 |
| 18-25 | 0.5 | 0.5 | 0.5 | 0.2 | 0.2 | 0.2 |
| 26-34 | 0.8 | 2.0 | 0.6 | 0.5 | 1.2 | 0.4 |
| 35-49 | 2.1 | 2.3 | 2.1 | 1.4 | 1.4 | 1.4 |
| 50-64 | 8.8 | 11.1 | 8.7 | 5.9 | 6.8 | 5.9 |
| 65 or Older | 20.6 | * | 20.8 | 15.4 | * | 15.6 |
| MALE | 6.5 | 2.8 | 7.0 | 4.4 | 1.6 | 4.8 |
| 12-17 | 0.4 | 0.5 | 0.4 | 0.2 | 0.5 | 0.2 |
| 18-25 | 0.4 | 0.3 | 0.4 | 0.1 | 0.1 | 0.1 |
| 26-34 | 0.6 | 0.6 | 0.6 | 0.3 | 0.2 | 0.3 |
| 35-49 | 2.2 | 2.0 | 2.3 | 1.5 | 1.2 | 1.5 |
| 50-64 | 11.3 | 13.8 | 11.1 | 7.2 | 8.7 | 7.1 |
| 65 or Older | 25.8 | * | 26.3 | 18.9 | * | 19.4 |
| FEMALE | 4.8 | 2.3 | 5.0 | 3.6 | 1.3 | 3.7 |
| 12-17 | 0.4 | 0.3 | 0.4 | 0.2 | 0.3 | 0.1 |
| 18-25 | 0.7 | 0.9 | 0.6 | 0.3 | 0.3 | 0.3 |
| 26-34 | 1.0 | 4.9 | 0.7 | 0.7 | * | 0.5 |
| 35-49 | 1.9 | 2.7 | 1.9 | 1.4 | 1.6 | 1.3 |
| 50-64 | 6.4 | * | 6.5 | 4.7 | * | 4.8 |
| 65 or Older | 16.7 | * | 16.8 | 12.8 | * | 12.9 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.121A Hepatitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | HEPATITIS IN LIFETIME | | | HEPATITIS IN PAST YEAR | | |
|---------------|-----------------------|--|-------|------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 3,537 | 384 | 3,153 | 1,107 | 187 | 919 |
| 12-17 | 38 | 1 | 36 | 9 | 0 | 9 |
| 18-25 | 134 | 46 | 88 | 53 | 17 | 36 |
| 26-34 | 311 | 40 | 271 | 135 | 25 | 109 |
| 35-49 | 1,193 | 238 | 955 | 405 | 110 | 295 |
| 50-64 | 1,282 | 59 | 1,223 | 391 | 35 | 356 |
| 65 or Older | 579 | * | 579 | 113 | * | 113 |
| MALE | 1,852 | 277 | 1,575 | 574 | 134 | 441 |
| 12-17 | 20 | 1 | 19 | 2 | 0 | 2 |
| 18-25 | 72 | 29 | 44 | 35 | 10 | 25 |
| 26-34 | 143 | 39 | 104 | 44 | 25 | 18 |
| 35-49 | 618 | 156 | 462 | 171 | 63 | 108 |
| 50-64 | 727 | 52 | 676 | 285 | 35 | 250 |
| 65 or Older | 271 | * | 271 | 38 | * | 38 |
| FEMALE | 1,685 | 108 | 1,577 | 532 | 54 | 478 |
| 12-17 | 18 | * | 18 | 7 | * | 7 |
| 18-25 | 62 | 18 | 45 | 18 | 7 | 12 |
| 26-34 | 167 | 1 | 166 | 91 | * | 91 |
| 35-49 | 575 | 82 | 493 | 234 | 47 | 187 |
| 50-64 | 555 | * | 548 | 106 | * | 106 |
| 65 or Older | 308 | * | 308 | 76 | * | 76 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.121B Hepatitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | HEPATITIS IN LIFETIME | | | HEPATITIS IN PAST YEAR | | |
|---------------|-----------------------|--|-----|------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 1.5 | 1.8 | 1.4 | 0.5 | 0.9 | 0.4 |
| 12-17 | 0.2 | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 |
| 18-25 | 0.4 | 0.7 | 0.4 | 0.2 | 0.2 | 0.1 |
| 26-34 | 0.9 | 0.9 | 0.9 | 0.4 | 0.6 | 0.4 |
| 35-49 | 1.8 | 4.2 | 1.6 | 0.6 | 1.9 | 0.5 |
| 50-64 | 2.6 | 2.5 | 2.6 | 0.8 | 1.5 | 0.8 |
| 65 or Older | 1.7 | * | 1.7 | 0.3 | * | 0.3 |
| MALE | 1.6 | 2.0 | 1.5 | 0.5 | 1.0 | 0.4 |
| 12-17 | 0.2 | 0.1 | 0.2 | 0.0 | 0.0 | 0.0 |
| 18-25 | 0.5 | 0.7 | 0.4 | 0.2 | 0.2 | 0.2 |
| 26-34 | 0.8 | 1.4 | 0.7 | 0.3 | 0.9 | 0.1 |
| 35-49 | 2.0 | 4.3 | 1.7 | 0.5 | 1.7 | 0.4 |
| 50-64 | 3.0 | 3.0 | 3.0 | 1.2 | 2.1 | 1.1 |
| 65 or Older | 1.8 | * | 1.9 | 0.3 | * | 0.3 |
| FEMALE | 1.4 | 1.4 | 1.4 | 0.4 | 0.7 | 0.4 |
| 12-17 | 0.1 | * | 0.2 | 0.1 | * | 0.1 |
| 18-25 | 0.4 | 0.7 | 0.3 | 0.1 | 0.2 | 0.1 |
| 26-34 | 1.0 | 0.1 | 1.0 | 0.5 | * | 0.6 |
| 35-49 | 1.7 | 4.0 | 1.6 | 0.7 | 2.3 | 0.6 |
| 50-64 | 2.2 | * | 2.2 | 0.4 | * | 0.4 |
| 65 or Older | 1.5 | * | 1.5 | 0.4 | * | 0.4 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.122A High Blood Pressure in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | HIGH BLOOD PRESSURE IN LIFETIME | | | HIGH BLOOD PRESSURE IN PAST YEAR | | |
|---------------|---------------------------------|--|--------|----------------------------------|--|--------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 46,844 | 2,736 | 44,107 | 37,582 | 2,106 | 35,476 |
| 12-17 | 453 | 40 | 413 | 275 | 19 | 256 |
| 18-25 | 1,181 | 241 | 941 | 795 | 164 | 631 |
| 26-34 | 2,476 | 278 | 2,198 | 1,602 | 161 | 1,441 |
| 35-49 | 9,831 | 1,054 | 8,776 | 7,787 | 775 | 7,011 |
| 50-64 | 16,732 | 867 | 15,865 | 13,887 | 766 | 13,122 |
| 65 or Older | 16,172 | * | 15,915 | 13,236 | * | 13,015 |
| MALE | 21,579 | 1,824 | 19,755 | 16,941 | 1,400 | 15,541 |
| 12-17 | 256 | 18 | 238 | 155 | 8 | 148 |
| 18-25 | 583 | 150 | 432 | 384 | 99 | 285 |
| 26-34 | 1,417 | 203 | 1,214 | 914 | 124 | 789 |
| 35-49 | 5,145 | 679 | 4,466 | 4,007 | 496 | 3,511 |
| 50-64 | 7,753 | 597 | 7,156 | 6,421 | 509 | 5,912 |
| 65 or Older | 6,426 | * | 6,250 | 5,060 | * | 4,897 |
| FEMALE | 25,264 | 912 | 24,352 | 20,641 | 706 | 19,935 |
| 12-17 | 197 | 22 | 175 | 120 | 11 | 109 |
| 18-25 | 599 | 90 | 509 | 411 | 65 | 346 |
| 26-34 | 1,058 | 75 | 984 | 688 | 37 | 652 |
| 35-49 | 4,686 | 375 | 4,311 | 3,780 | 279 | 3,501 |
| 50-64 | 8,979 | * | 8,709 | 7,466 | * | 7,210 |
| 65 or Older | 9,746 | * | 9,665 | 8,176 | * | 8,118 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.122B High Blood Pressure in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | HIGH BLOOD PRESSURE IN LIFETIME | | | HIGH BLOOD PRESSURE IN PAST YEAR | | |
|---------------|---------------------------------|--|------|----------------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 19.6 | 12.6 | 20.3 | 15.7 | 9.7 | 16.3 |
| 12-17 | 1.9 | 2.0 | 1.9 | 1.1 | 1.0 | 1.2 |
| 18-25 | 3.7 | 3.5 | 3.8 | 2.5 | 2.4 | 2.5 |
| 26-34 | 7.2 | 6.6 | 7.3 | 4.7 | 3.8 | 4.8 |
| 35-49 | 15.2 | 18.6 | 14.9 | 12.1 | 13.7 | 11.9 |
| 50-64 | 33.7 | 36.3 | 33.5 | 28.0 | 32.0 | 27.8 |
| 65 or Older | 46.2 | * | 46.3 | 37.8 | * | 37.9 |
| MALE | 18.7 | 13.2 | 19.4 | 14.7 | 10.1 | 15.3 |
| 12-17 | 2.1 | 1.9 | 2.1 | 1.3 | 0.8 | 1.3 |
| 18-25 | 3.7 | 3.5 | 3.7 | 2.4 | 2.3 | 2.4 |
| 26-34 | 8.4 | 7.1 | 8.6 | 5.4 | 4.4 | 5.6 |
| 35-49 | 16.2 | 18.6 | 15.9 | 12.7 | 13.6 | 12.5 |
| 50-64 | 32.2 | 35.0 | 32.0 | 26.7 | 29.9 | 26.5 |
| 65 or Older | 43.0 | * | 43.1 | 33.9 | * | 33.8 |
| FEMALE | 20.4 | 11.5 | 21.0 | 16.7 | 8.9 | 17.2 |
| 12-17 | 1.6 | 2.2 | 1.6 | 1.0 | 1.1 | 1.0 |
| 18-25 | 3.8 | 3.4 | 3.8 | 2.6 | 2.4 | 2.6 |
| 26-34 | 6.1 | 5.5 | 6.2 | 4.0 | 2.7 | 4.1 |
| 35-49 | 14.3 | 18.6 | 14.0 | 11.5 | 13.8 | 11.3 |
| 50-64 | 35.0 | * | 34.9 | 29.1 | * | 28.9 |
| 65 or Older | 48.6 | * | 48.6 | 40.8 | * | 40.8 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.123A Pneumonia in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | PNEUMONIA IN LIFETIME | | | PNEUMONIA IN PAST YEAR | | |
|---------------|-----------------------|--|--------|------------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 15,775 | 996 | 14,779 | 3,680 | 208 | 3,472 |
| 12-17 | 824 | 62 | 762 | 225 | 22 | 203 |
| 18-25 | 1,142 | 291 | 852 | 259 | 73 | 186 |
| 26-34 | 1,501 | 121 | 1,380 | 231 | 6 | 225 |
| 35-49 | 3,755 | 250 | 3,504 | 845 | 53 | 792 |
| 50-64 | 4,950 | 184 | 4,766 | 1,095 | 24 | 1,071 |
| 65 or Older | 3,603 | * | 3,515 | 1,025 | * | 995 |
| MALE | 6,392 | 570 | 5,822 | 1,413 | 141 | 1,272 |
| 12-17 | 361 | 23 | 338 | 104 | 10 | 94 |
| 18-25 | 498 | 161 | 338 | 99 | 38 | 61 |
| 26-34 | 554 | 69 | 485 | 87 | 5 | 82 |
| 35-49 | 1,401 | 141 | 1,260 | 277 | 34 | 243 |
| 50-64 | 2,065 | 109 | 1,957 | 415 | * | 391 |
| 65 or Older | 1,513 | * | 1,445 | 430 | * | 401 |
| FEMALE | 9,383 | 426 | 8,957 | 2,267 | 67 | 2,200 |
| 12-17 | 463 | 39 | 424 | 121 | 12 | 109 |
| 18-25 | 644 | 130 | 514 | 160 | 35 | 125 |
| 26-34 | 947 | 51 | 896 | 144 | 1 | 143 |
| 35-49 | 2,353 | 110 | 2,244 | 568 | 19 | 549 |
| 50-64 | 2,885 | * | 2,809 | 680 | * | 680 |
| 65 or Older | 2,090 | * | 2,070 | 594 | * | 594 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.123B Pneumonia in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | PNEUMONIA IN LIFETIME | | | PNEUMONIA IN PAST YEAR | | |
|---------------|-----------------------|--|------|------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 6.6 | 4.6 | 6.8 | 1.5 | 1.0 | 1.6 |
| 12-17 | 3.4 | 3.2 | 3.4 | 0.9 | 1.1 | 0.9 |
| 18-25 | 3.6 | 4.2 | 3.4 | 0.8 | 1.0 | 0.7 |
| 26-34 | 4.4 | 2.9 | 4.6 | 0.7 | 0.1 | 0.7 |
| 35-49 | 5.8 | 4.4 | 6.0 | 1.3 | 0.9 | 1.3 |
| 50-64 | 10.0 | 7.7 | 10.1 | 2.2 | 1.0 | 2.3 |
| 65 or Older | 10.3 | * | 10.2 | 2.9 | * | 2.9 |
| MALE | 5.5 | 4.1 | 5.7 | 1.2 | 1.0 | 1.3 |
| 12-17 | 3.0 | 2.4 | 3.0 | 0.9 | 1.1 | 0.8 |
| 18-25 | 3.1 | 3.8 | 2.9 | 0.6 | 0.9 | 0.5 |
| 26-34 | 3.3 | 2.4 | 3.4 | 0.5 | 0.2 | 0.6 |
| 35-49 | 4.4 | 3.9 | 4.5 | 0.9 | 0.9 | 0.9 |
| 50-64 | 8.6 | 6.4 | 8.8 | 1.7 | * | 1.8 |
| 65 or Older | 10.1 | * | 10.0 | 2.9 | * | 2.8 |
| FEMALE | 7.6 | 5.4 | 7.7 | 1.8 | 0.8 | 1.9 |
| 12-17 | 3.9 | 3.9 | 3.9 | 1.0 | 1.2 | 1.0 |
| 18-25 | 4.0 | 4.9 | 3.9 | 1.0 | 1.3 | 0.9 |
| 26-34 | 5.5 | 3.8 | 5.6 | 0.8 | 0.1 | 0.9 |
| 35-49 | 7.2 | 5.4 | 7.3 | 1.7 | 0.9 | 1.8 |
| 50-64 | 11.3 | * | 11.3 | 2.7 | * | 2.7 |
| 65 or Older | 10.4 | * | 10.4 | 3.0 | * | 3.0 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.124A Sexually Transmitted Disease (STD) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | STD IN LIFETIME | | | STD IN PAST YEAR | | |
|---------------|-----------------|--|-------|------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 8,205 | 1,475 | 6,730 | 1,994 | 504 | 1,491 |
| 12-17 | 115 | 45 | 69 | 90 | 39 | 51 |
| 18-25 | 1,249 | 426 | 824 | 679 | 259 | 420 |
| 26-34 | 1,930 | 380 | 1,549 | 537 | 89 | 448 |
| 35-49 | 3,006 | 473 | 2,534 | 532 | 105 | 427 |
| 50-64 | 1,688 | 140 | 1,548 | 131 | * | 131 |
| 65 or Older | 217 | * | 206 | 25 | * | 14 |
| MALE | 3,026 | 624 | 2,402 | 569 | 186 | 383 |
| 12-17 | 22 | 8 | 14 | 13 | 7 | 6 |
| 18-25 | 235 | 117 | 118 | 133 | 73 | 60 |
| 26-34 | 605 | 213 | 392 | 183 | 57 | 126 |
| 35-49 | 1,245 | 223 | 1,022 | 210 | 37 | 173 |
| 50-64 | 752 | 51 | 701 | 4 | * | 4 |
| 65 or Older | 167 | * | 156 | 25 | * | 14 |
| FEMALE | 5,179 | 852 | 4,328 | 1,425 | 318 | 1,108 |
| 12-17 | 93 | 37 | 56 | 77 | 32 | 45 |
| 18-25 | 1,014 | 308 | 706 | 545 | 186 | 360 |
| 26-34 | 1,325 | 167 | 1,158 | 354 | 32 | 322 |
| 35-49 | 1,761 | 250 | 1,511 | 322 | 68 | 254 |
| 50-64 | 936 | * | 846 | 127 | * | 127 |
| 65 or Older | 51 | * | 51 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

NOTE: Sexually Transmitted Disease (STD) includes chlamydia, gonorrhea, herpes, and syphilis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.124B Sexually Transmitted Disease (STD) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | STD IN LIFETIME | | | STD IN PAST YEAR | | |
|---------------|-----------------|--|-----|------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 3.4 | 6.8 | 3.1 | 0.8 | 2.3 | 0.7 |
| 12-17 | 0.5 | 2.3 | 0.3 | 0.4 | 2.0 | 0.2 |
| 18-25 | 3.9 | 6.1 | 3.3 | 2.1 | 3.7 | 1.7 |
| 26-34 | 5.6 | 9.0 | 5.2 | 1.6 | 2.1 | 1.5 |
| 35-49 | 4.7 | 8.3 | 4.3 | 0.8 | 1.9 | 0.7 |
| 50-64 | 3.4 | 5.9 | 3.3 | 0.3 | * | 0.3 |
| 65 or Older | 0.6 | * | 0.6 | 0.1 | * | 0.0 |
| MALE | 2.6 | 4.5 | 2.4 | 0.5 | 1.3 | 0.4 |
| 12-17 | 0.2 | 0.9 | 0.1 | 0.1 | 0.8 | 0.1 |
| 18-25 | 1.5 | 2.8 | 1.0 | 0.8 | 1.7 | 0.5 |
| 26-34 | 3.6 | 7.5 | 2.8 | 1.1 | 2.0 | 0.9 |
| 35-49 | 3.9 | 6.1 | 3.6 | 0.7 | 1.0 | 0.6 |
| 50-64 | 3.1 | 3.0 | 3.1 | 0.0 | * | 0.0 |
| 65 or Older | 1.1 | * | 1.1 | 0.2 | * | 0.1 |
| FEMALE | 4.2 | 10.7 | 3.7 | 1.2 | 4.0 | 1.0 |
| 12-17 | 0.8 | 3.7 | 0.5 | 0.6 | 3.2 | 0.4 |
| 18-25 | 6.4 | 11.5 | 5.3 | 3.4 | 6.9 | 2.7 |
| 26-34 | 7.6 | 12.2 | 7.2 | 2.0 | 2.3 | 2.0 |
| 35-49 | 5.4 | 12.4 | 4.9 | 1.0 | 3.4 | 0.8 |
| 50-64 | 3.6 | * | 3.4 | 0.5 | * | 0.5 |
| 65 or Older | 0.3 | * | 0.3 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

NOTE: Sexually Transmitted Disease (STD) includes chlamydia, gonorrhea, herpes, and syphilis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.125A Stroke in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | STROKE IN LIFETIME | | | STROKE IN PAST YEAR | | |
|---------------|--------------------|--|-------|---------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 3,339 | 155 | 3,184 | 1,111 | 65 | 1,046 |
| 12-17 | 31 | 6 | 25 | 5 | 0 | 5 |
| 18-25 | 20 | 8 | 12 | 4 | 3 | * |
| 26-34 | 68 | 4 | 64 | 13 | * | 13 |
| 35-49 | 267 | 40 | 227 | 64 | 5 | 59 |
| 50-64 | 1,155 | 88 | 1,067 | 298 | 56 | 242 |
| 65 or Older | 1,798 | * | 1,788 | 727 | * | 727 |
| MALE | 1,468 | 78 | 1,391 | 546 | 49 | 497 |
| 12-17 | 15 | 2 | 13 | 4 | 0 | 4 |
| 18-25 | 10 | 6 | 4 | 3 | 3 | * |
| 26-34 | 27 | * | 27 | 3 | * | 3 |
| 35-49 | 127 | 11 | 116 | 31 | * | 31 |
| 50-64 | 509 | 58 | 451 | 153 | 46 | 107 |
| 65 or Older | 780 | * | 780 | 351 | * | 351 |
| FEMALE | 1,871 | 78 | 1,793 | 565 | 16 | 549 |
| 12-17 | 16 | 3 | 13 | * | * | 1 |
| 18-25 | 10 | 1 | 9 | * | * | * |
| 26-34 | 41 | 4 | 37 | 10 | * | 10 |
| 35-49 | 140 | 30 | 111 | 33 | 5 | 28 |
| 50-64 | 645 | * | 616 | 145 | * | 134 |
| 65 or Older | 1,019 | * | 1,009 | 376 | * | 376 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.125B Stroke in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | STROKE IN LIFETIME | | | STROKE IN PAST YEAR | | |
|---------------|--------------------|--|-----|---------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 1.4 | 0.7 | 1.5 | 0.5 | 0.3 | 0.5 |
| 12-17 | 0.1 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 |
| 18-25 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | * |
| 26-34 | 0.2 | 0.1 | 0.2 | 0.0 | * | 0.0 |
| 35-49 | 0.4 | 0.7 | 0.4 | 0.1 | 0.1 | 0.1 |
| 50-64 | 2.3 | 3.7 | 2.3 | 0.6 | 2.4 | 0.5 |
| 65 or Older | 5.1 | * | 5.2 | 2.1 | * | 2.1 |
| MALE | 1.3 | 0.6 | 1.4 | 0.5 | 0.4 | 0.5 |
| 12-17 | 0.1 | 0.3 | 0.1 | 0.0 | 0.0 | 0.0 |
| 18-25 | 0.1 | 0.2 | 0.0 | 0.0 | 0.1 | * |
| 26-34 | 0.2 | * | 0.2 | 0.0 | * | 0.0 |
| 35-49 | 0.4 | 0.3 | 0.4 | 0.1 | * | 0.1 |
| 50-64 | 2.1 | 3.4 | 2.0 | 0.6 | 2.7 | 0.5 |
| 65 or Older | 5.2 | * | 5.4 | 2.3 | * | 2.4 |
| FEMALE | 1.5 | 1.0 | 1.5 | 0.5 | 0.2 | 0.5 |
| 12-17 | 0.1 | 0.3 | 0.1 | * | * | 0.0 |
| 18-25 | 0.1 | 0.0 | 0.1 | * | * | * |
| 26-34 | 0.2 | 0.3 | 0.2 | 0.1 | * | 0.1 |
| 35-49 | 0.4 | 1.5 | 0.4 | 0.1 | 0.2 | 0.1 |
| 50-64 | 2.5 | * | 2.5 | 0.6 | * | 0.5 |
| 65 or Older | 5.1 | * | 5.1 | 1.9 | * | 1.9 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.126A Ulcer(s) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Numbers in Thousands, 2005

| Gender/Age | ULCER(S) IN LIFETIME | | | ULCER(S) IN PAST YEAR | | |
|---------------|----------------------|--|-------|-----------------------|--|-------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 8,974 | 841 | 8,133 | 2,827 | 223 | 2,604 |
| 12-17 | 200 | 26 | 174 | 94 | 16 | 78 |
| 18-25 | 666 | 175 | 491 | 342 | 76 | 267 |
| 26-34 | 1,018 | 156 | 863 | 398 | 71 | 327 |
| 35-49 | 2,563 | 284 | 2,279 | 807 | 45 | 761 |
| 50-64 | 2,467 | 161 | 2,307 | 603 | * | 588 |
| 65 or Older | 2,059 | * | 2,020 | 583 | * | 583 |
| MALE | 3,693 | 460 | 3,233 | 969 | 101 | 868 |
| 12-17 | 90 | 13 | 77 | 44 | 8 | 36 |
| 18-25 | 264 | 86 | 179 | 127 | 29 | 98 |
| 26-34 | 408 | 113 | 295 | 164 | 49 | 115 |
| 35-49 | 994 | 156 | 838 | 267 | 15 | 252 |
| 50-64 | 993 | 64 | 929 | 149 | * | 149 |
| 65 or Older | 944 | * | 915 | 218 | * | 218 |
| FEMALE | 5,281 | 381 | 4,900 | 1,858 | 122 | 1,736 |
| 12-17 | 110 | 13 | 97 | 50 | 8 | 42 |
| 18-25 | 402 | 90 | 312 | 216 | 47 | 169 |
| 26-34 | 610 | 43 | 567 | 234 | 22 | 212 |
| 35-49 | 1,569 | 128 | 1,441 | 540 | 31 | 509 |
| 50-64 | 1,475 | * | 1,378 | 454 | * | 439 |
| 65 or Older | 1,115 | * | 1,104 | 365 | * | 365 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.126B Ulcer(s) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Percentages, 2005

| Gender/Age | ULCER(S) IN LIFETIME | | | ULCER(S) IN PAST YEAR | | |
|---------------|----------------------|--|-----|-----------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 3.7 | 3.9 | 3.7 | 1.2 | 1.0 | 1.2 |
| 12-17 | 0.8 | 1.3 | 0.8 | 0.4 | 0.8 | 0.4 |
| 18-25 | 2.1 | 2.5 | 2.0 | 1.1 | 1.1 | 1.1 |
| 26-34 | 3.0 | 3.7 | 2.9 | 1.2 | 1.7 | 1.1 |
| 35-49 | 4.0 | 5.0 | 3.9 | 1.3 | 0.8 | 1.3 |
| 50-64 | 5.0 | 6.7 | 4.9 | 1.2 | * | 1.2 |
| 65 or Older | 5.9 | * | 5.9 | 1.7 | * | 1.7 |
| MALE | 3.2 | 3.3 | 3.2 | 0.8 | 0.7 | 0.9 |
| 12-17 | 0.7 | 1.4 | 0.7 | 0.4 | 0.9 | 0.3 |
| 18-25 | 1.7 | 2.0 | 1.5 | 0.8 | 0.7 | 0.8 |
| 26-34 | 2.4 | 3.9 | 2.1 | 1.0 | 1.7 | 0.8 |
| 35-49 | 3.1 | 4.3 | 3.0 | 0.8 | 0.4 | 0.9 |
| 50-64 | 4.1 | 3.8 | 4.2 | 0.6 | * | 0.7 |
| 65 or Older | 6.3 | * | 6.3 | 1.5 | * | 1.5 |
| FEMALE | 4.3 | 4.8 | 4.2 | 1.5 | 1.5 | 1.5 |
| 12-17 | 0.9 | 1.3 | 0.9 | 0.4 | 0.8 | 0.4 |
| 18-25 | 2.5 | 3.4 | 2.4 | 1.4 | 1.7 | 1.3 |
| 26-34 | 3.5 | 3.1 | 3.5 | 1.3 | 1.6 | 1.3 |
| 35-49 | 4.8 | 6.3 | 4.7 | 1.6 | 1.5 | 1.6 |
| 50-64 | 5.8 | * | 5.5 | 1.8 | * | 1.8 |
| 65 or Older | 5.6 | * | 5.5 | 1.8 | * | 1.8 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.