

**Table 7.127A No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drugs or Alcohol <sup>1</sup>		Illicit Drugs <sup>1</sup>		Alcohol	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	19,734 <sup>b</sup>	20,108	22,540 <sup>b</sup>	22,844	20,779 <sup>b</sup>	21,164
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	231	237	259	258	261	263
Asian or Pacific Islander	954	1,055	1,024	1,120	983	1,082
Asian Only	885	967	945	1,029	910	989
Native Hawaiian or Other Pacific Islander Only	*	*	78	*	*	93
Black or African American	3,350	3,272	3,598	3,580	3,580	3,549
White	14,832 <sup>b</sup>	15,174	17,241 <sup>b</sup>	17,484	15,563 <sup>b</sup>	15,884
Two or More Races	368	370	419	402	392	385
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	3,283 <sup>b</sup>	3,466	3,800 <sup>b</sup>	3,981	3,471 <sup>b</sup>	3,661
Not Hispanic or Latino	16,451	16,642	18,741	18,863	17,307	17,503
Black or African American	3,215	3,154	3,449	3,442	3,432	3,419
White	11,857	12,007	13,781	13,843	12,419	12,542
<b>GENDER</b>						
Male	10,145 <sup>a</sup>	10,338	11,517	11,637	10,661 <sup>b</sup>	10,893
Female	9,589 <sup>a</sup>	9,769	11,024 <sup>b</sup>	11,207	10,117	10,270
<b>FAMILY INCOME</b>						
Less Than \$20,000	3,622	3,572	4,033	3,994	3,894	3,836
\$20,000 - \$49,999	6,741	6,736	7,683	7,536	7,121	7,135
\$50,000 - \$74,999	3,782	3,724	4,355	4,290	3,942	3,896
\$75,000 or More	5,590 <sup>a</sup>	6,076	6,470 <sup>a</sup>	7,024	5,822 <sup>a</sup>	6,297
<b>REGION</b>						
Northeast	3,449	3,551	3,994 <sup>a</sup>	4,086	3,663	3,705
Midwest	4,390	4,422	5,098	5,089	4,622	4,657
South	7,262	7,243	8,178	8,187	7,603	7,635
West	4,633 <sup>b</sup>	4,892	5,270 <sup>b</sup>	5,482	4,891 <sup>b</sup>	5,167

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.127B No Illicit Drug or Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drugs or Alcohol <sup>1</sup>		Illicit Drugs <sup>1</sup>		Alcohol	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	78.3 <sup>a</sup>	79.3	89.4	90.1	82.4 <sup>a</sup>	83.5
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	72.8	77.3	81.7	84.1	82.3	85.9
Asian or Pacific Islander	87.5	90.2	94.0	95.8	90.2	92.6
Asian Only	88.0	90.8	94.0	96.6	90.5	92.8
Native Hawaiian or Other Pacific Islander Only	*	*	93.6	*	*	89.5
Black or African American	84.3 <sup>b</sup>	81.3	90.5	88.9	90.1 <sup>a</sup>	88.2
White	76.6 <sup>b</sup>	78.2	89.1 <sup>a</sup>	90.1	80.4 <sup>b</sup>	81.8
Two or More Races	77.5 <sup>a</sup>	83.2	88.3	90.5	82.5	86.7
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	77.6	78.9	89.8	90.6	82.0	83.3
Not Hispanic or Latino	78.4	79.4	89.3	90.0	82.5 <sup>a</sup>	83.5
Black or African American	84.5 <sup>b</sup>	81.5	90.7	89.0	90.2	88.4
White	76.5 <sup>a</sup>	78.0	88.9 <sup>a</sup>	89.9	80.1 <sup>a</sup>	81.5
<b>GENDER</b>						
Male	78.8	79.8	89.4	89.9	82.8 <sup>a</sup>	84.1
Female	77.7	78.8	89.4	90.3	82.0	82.8
<b>FAMILY INCOME</b>						
Less Than \$20,000	78.4	79.2	87.3	88.6	84.3	85.0
\$20,000 - \$49,999	78.6 <sup>a</sup>	80.5	89.6	90.0	83.1 <sup>b</sup>	85.2
\$50,000 - \$74,999	78.0	78.5	89.8	90.5	81.3	82.2
\$75,000 or More	77.9	78.6	90.2	90.9	81.1	81.4
<b>REGION</b>						
Northeast	76.0	78.1	88.1 <sup>a</sup>	89.9	80.8	81.5
Midwest	77.3	78.0	89.7	89.8	81.4	82.2
South	80.7	79.9	90.9	90.3	84.5	84.2
West	77.3 <sup>b</sup>	80.5	87.9 <sup>b</sup>	90.2	81.6 <sup>b</sup>	85.0

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.128A Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use <sup>1</sup>		Marijuana Use		Binge Alcohol Use <sup>2</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	2,674	2,511	1,909 <sup>a</sup>	1,730	2,794 <sup>b</sup>	2,513
<b>RACE<sup>3</sup></b>						
American Indian or Alaska Native	58	49	36	33	42	32
Asian or Pacific Islander	66	49	46	24	56	39
Asian Only	60	36	43 <sup>a</sup>	16	50	33
Native Hawaiian or Other Pacific Islander Only	5	*	3	*	*	6
Black or African American	376	445	257	293	198	224
White	2,119 <sup>a</sup>	1,926	1,525 <sup>a</sup>	1,350	2,443 <sup>b</sup>	2,179
Two or More Races	56	42	45	30	54	39
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	433	412	284	275	519	467
Not Hispanic or Latino	2,240	2,099	1,624 <sup>a</sup>	1,455	2,275 <sup>b</sup>	2,046
Black or African American	354	427	243	278	186	209
White	1,727 <sup>a</sup>	1,556	1,266 <sup>a</sup>	1,103	1,957 <sup>a</sup>	1,751
<b>GENDER</b>						
Male	1,360	1,313	1,037	967	1,496 <sup>a</sup>	1,341
Female	1,314	1,198	872 <sup>a</sup>	763	1,298	1,172
<b>FAMILY INCOME</b>						
Less Than \$20,000	585	516	406	332	466	427
\$20,000 - \$49,999	890	836	632	551	915 <sup>b</sup>	720
\$50,000 - \$74,999	493	452	351	307	565	506
\$75,000 or More	706	707	520	540	848	859
<b>REGION</b>						
Northeast	542	459	389	360	519	492
Midwest	584	577	417	389	700	641
South	823	878	544	566	845	832
West	726 <sup>a</sup>	597	559 <sup>b</sup>	415	730 <sup>b</sup>	548

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.<sup>3</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.128B Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use <sup>1</sup>		Marijuana Use		Binge Alcohol Use <sup>2</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	10.6	9.9	7.6 <sup>a</sup>	6.8	11.1 <sup>b</sup>	9.9
<b>RACE<sup>3</sup></b>						
American Indian or Alaska Native	18.3	15.9	11.3	10.9	13.4	10.4
Asian or Pacific Islander	6.0	4.2	4.2	2.0	5.2	3.3
Asian Only	6.0	3.4	4.3 <sup>a</sup>	1.5	5.0	3.1
Native Hawaiian or Other Pacific Islander Only	6.4	*	3.9	*	*	6.0
Black or African American	9.5	11.1	6.5	7.3	5.0	5.6
White	10.9 <sup>a</sup>	9.9	7.9 <sup>a</sup>	7.0	12.6 <sup>b</sup>	11.2
Two or More Races	11.7	9.5	9.5	6.8	11.3	8.9
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	10.2	9.4	6.7	6.3	12.3	10.6
Not Hispanic or Latino	10.7	10.0	7.7 <sup>a</sup>	6.9	10.8 <sup>b</sup>	9.8
Black or African American	9.3	11.0	6.4	7.2	4.9	5.4
White	11.1 <sup>a</sup>	10.1	8.2 <sup>a</sup>	7.2	12.6 <sup>b</sup>	11.4
<b>GENDER</b>						
Male	10.6	10.1	8.1	7.5	11.6 <sup>a</sup>	10.4
Female	10.6	9.7	7.1 <sup>a</sup>	6.2	10.5 <sup>a</sup>	9.4
<b>FAMILY INCOME</b>						
Less Than \$20,000	12.7	11.4	8.8	7.4	10.1	9.5
\$20,000 - \$49,999	10.4	10.0	7.4	6.6	10.7 <sup>b</sup>	8.6
\$50,000 - \$74,999	10.2	9.5	7.2	6.5	11.7	10.7
\$75,000 or More	9.8	9.1	7.2	7.0	11.8	11.1
<b>REGION</b>						
Northeast	11.9 <sup>a</sup>	10.1	8.6	7.9	11.4	10.8
Midwest	10.3	10.2	7.3	6.9	12.3	11.3
South	9.1	9.7	6.0	6.2	9.4	9.2
West	12.1 <sup>b</sup>	9.8	9.3 <sup>b</sup>	6.8	12.2 <sup>b</sup>	9.0

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.<sup>3</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.129A Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use <sup>1</sup>		Marijuana Use		Binge Alcohol Use <sup>2</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	16,397	17,209	12,667	12,896	51,931	52,577
<b>RACE<sup>3</sup></b>						
American Indian or Alaska Native	163	189	117	138	540	644
Asian or Pacific Islander	369	354	245	202	1,402	1,469
Asian Only	255	285	156	152	1,202	1,262
Native Hawaiian or Other Pacific Islander Only	*	68	*	51	*	208
Black or African American	2,156	2,423	1,797	1,978	5,061 <sup>a</sup>	5,734
White	13,403	13,938	10,297	10,303	44,282	44,114
Two or More Races	306	305	211	275	646	616
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	1,806	2,028	1,258	1,353	6,929	7,144
Not Hispanic or Latino	14,591	15,181	11,408	11,543	45,002	45,433
Black or African American	2,075	2,341	1,725	1,908	4,936	5,604
White	11,756	12,098	9,158	9,100	37,844	37,487
<b>GENDER</b>						
Male	10,132	10,768	8,236	8,706	34,699	34,684
Female	6,265	6,441	4,431	4,190	17,232	17,893
<b>FAMILY INCOME</b>						
Less Than \$20,000	4,594	4,752	3,723	3,501	10,373	10,410
\$20,000 - \$49,999	5,617 <sup>a</sup>	6,490	4,277 <sup>a</sup>	4,968	18,245	18,217
\$50,000 - \$74,999	2,721	2,638	2,038	1,916	9,292	9,378
\$75,000 or More	3,465	3,329	2,629	2,512	14,021	14,571
<b>EDUCATION</b>						
< High School	3,131	3,507	2,399	2,660	8,407	8,277
High School Graduate	5,344	5,868	4,057 <sup>a</sup>	4,621	17,144	17,221
Some College/College Graduate	7,922	7,834	6,211	5,615	26,381	27,079

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>3</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.129B Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use <sup>1</sup>		Marijuana Use		Binge Alcohol Use <sup>2</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	7.6	7.9	5.9	5.9	24.1	24.1
<b>RACE<sup>3</sup></b>						
American Indian or Alaska Native	8.5	9.6	6.1	7.0	28.2	32.6
Asian or Pacific Islander	3.8	3.6	2.5	2.0	14.5	14.8
Asian Only	2.8	3.1	1.7	1.7	13.3	13.8
Native Hawaiian or Other Pacific Islander Only	*	9.0	*	6.6	*	27.2
Black or African American	8.7	9.6	7.2	7.8	20.4	22.7
White	7.6	7.8	5.8	5.8	25.1	24.8
Two or More Races	12.4	11.7	8.6	10.6	26.2	23.7
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	6.7	7.3	4.7	4.9	25.9	25.8
Not Hispanic or Latino	7.7	8.0	6.1	6.1	23.9	23.9
Black or African American	8.6	9.5	7.1	7.7	20.4	22.7
White	7.8	7.9	6.0	6.0	25.0	24.6
<b>GENDER</b>						
Male	9.8	10.3	7.9	8.3	33.5	33.0
Female	5.6	5.7	4.0	3.7	15.4	15.8
<b>FAMILY INCOME</b>						
Less Than \$20,000	10.7	11.4	8.6	8.4	24.1	24.9
\$20,000 - \$49,999	7.2 <sup>b</sup>	8.6	5.5 <sup>b</sup>	6.6	23.4	24.0
\$50,000 - \$74,999	7.1	6.7	5.3	4.9	24.3	23.8
\$75,000 or More	6.2	5.5	4.7	4.1	25.1	23.9
<b>EDUCATION</b>						
< High School	8.6 <sup>a</sup>	9.8	6.6	7.4	23.0	23.2
High School Graduate	7.8	8.6	6.0 <sup>a</sup>	6.7	25.2	25.1
Some College/College Graduate	7.2	6.9	5.6 <sup>a</sup>	4.9	23.8	23.8

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>3</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.130A Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2004	2005
<b>TOTAL</b>	1,168	1,144
<b>RACE<sup>1</sup></b>		
American Indian or Alaska Native	18	17
Asian or Pacific Islander	49	28
Asian Only	47	20
Native Hawaiian or Other Pacific Islander Only	2	*
Black or African American	107	146
White	966	928
Two or More Races	29	25
<b>HISPANIC ORIGIN AND RACE</b>		
Hispanic or Latino	202	195
Not Hispanic or Latino	967	949
Black or African American	98	142
White	783	750
<b>GENDER</b>		
Male	587	538
Female	581	606
<b>FAMILY INCOME</b>		
Less Than \$20,000	191	180
\$20,000 - \$49,999	376	392
\$50,000 - \$74,999	258	236
\$75,000 or More	343	336
<b>REGION</b>		
Northeast	227	207
Midwest	260	245
South	373	394
West	308	297

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.130B Inhalant Use in Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2004	2005
<b>TOTAL</b>	4.6	4.5
<b>RACE<sup>1</sup></b>		
American Indian or Alaska Native	5.7	5.4
Asian or Pacific Islander	4.5	2.4
Asian Only	4.7	1.9
Native Hawaiian or Other Pacific Islander Only	2.2	*
Black or African American	2.7	3.6
White	5.0	4.8
Two or More Races	6.0	5.7
<b>HISPANIC ORIGIN AND RACE</b>		
Hispanic or Latino	4.8	4.4
Not Hispanic or Latino	4.6	4.5
Black or African American	2.6	3.7
White	5.1	4.9
<b>GENDER</b>		
Male	4.6	4.2
Female	4.7	4.9
<b>FAMILY INCOME</b>		
Less Than \$20,000	4.1	4.0
\$20,000 - \$49,999	4.4	4.7
\$50,000 - \$74,999	5.3	5.0
\$75,000 or More	4.8	4.3
<b>REGION</b>		
Northeast	5.0	4.6
Midwest	4.6	4.3
South	4.1	4.3
West	5.1	4.9

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 7.131A Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	305	279	1,410	1,294	930	823
<b>RACE<sup>1</sup></b>						
American Indian or Alaska Native	7	12	20	*	18	25
Asian or Pacific Islander	*	10	43	32	20	12
Asian Only	*	*	41	24	13	9
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	54	42	226	208	129	128
White	231	209	1,082	1,000	735	639
Two or More Races	8	7	39	29	27	20
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	63	61	258	266	131	150
Not Hispanic or Latino	242	218	1,152 <sup>a</sup>	1,027	799 <sup>a</sup>	673
Black or African American	54	42	209	206	125	116
White	170	152	852	756	614 <sup>a</sup>	518
<b>GENDER</b>						
Male	179	151	744	660	471	414
Female	126	128	666	633	458	409
<b>FAMILY INCOME</b>						
Less Than \$20,000	99	77	305	260	231	196
\$20,000 - \$49,999	104	101	518	487	365	363
\$50,000 - \$74,999	38	45	252	236	161	136
\$75,000 or More	64	55	335	311	173 <sup>a</sup>	128
<b>REGION</b>						
Northeast	45	37	263	200	137	126
Midwest	74 <sup>a</sup>	43	344	320	246 <sup>a</sup>	188
South	106	105	485	489	352	350
West	80	95	318	284	195	159

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.131B Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	3.6	3.4	16.9	15.7	11.1	10.0
<b>RACE<sup>1</sup></b>						
American Indian or Alaska Native	6.3	10.2	18.2	*	16.1	22.4
Asian or Pacific Islander	*	2.7	12.3	9.0	5.7	3.4
Asian Only	*	*	12.7	7.4	4.1	2.9
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	3.9	3.3	16.2	16.0	9.3	9.8
White	3.7	3.3	17.1	15.9	11.6 <sup>a</sup>	10.2
Two or More Races	4.6	4.0	21.2	17.1	14.9	11.5
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	4.2	3.9	17.5	17.3	8.9	9.7
Not Hispanic or Latino	3.5	3.3	16.7	15.4	11.6 <sup>a</sup>	10.1
Black or African American	4.0	3.4	15.7	16.5	9.4	9.3
White	3.4	3.1	17.1	15.5	12.3 <sup>a</sup>	10.6
<b>GENDER</b>						
Male	4.2	3.6	17.5	15.9	11.1	10.0
Female	3.0	3.1	16.2	15.5	11.1	10.0
<b>FAMILY INCOME</b>						
Less Than \$20,000	6.2	5.1	19.2	17.3	14.5	13.0
\$20,000 - \$49,999	3.6	3.5	17.8	17.0	12.5	12.7
\$50,000 - \$74,999	2.4	3.2	16.1	16.4	10.2	9.5
\$75,000 or More	2.8	2.3	14.6	12.8	7.6 <sup>a</sup>	5.3
<b>REGION</b>						
Northeast	3.0	2.5	17.6 <sup>a</sup>	13.7	9.2	8.6
Midwest	3.9 <sup>a</sup>	2.3	18.2	17.6	13.0 <sup>a</sup>	10.3
South	3.5	3.6	16.1	16.7	11.7	11.9
West	4.0	4.7	16.1	14.2	9.9	7.9

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.132A Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month <sup>1</sup>		Use Cocaine Once a Month <sup>1</sup>		Have Five or More Drinks Once or Twice a Week <sup>1</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	8,699	8,487	12,320	12,151	9,514	9,636
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	109	90	154	135	133	117
Asian or Pacific Islander	401	436	468	507	507	591
Asian Only	374	397	423	451	474	538
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	1,395	1,355	2,134	2,161	1,922	1,932
White	6,642	6,486	9,327	9,141	6,777	6,827
Two or More Races	153	119	237	206	175	169
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	1,553	1,545	1,874 <sup>a</sup>	2,024	1,647	1,694
Not Hispanic or Latino	7,146	6,942	10,446 <sup>a</sup>	10,126	7,868	7,942
Black or African American	1,329	1,299	2,043	2,076	1,850	1,863
White	5,233	5,078	7,640	7,301	5,284	5,284
<b>GENDER</b>						
Male	4,130	3,975	6,337	6,248	4,336	4,475
Female	4,569	4,511	5,984	5,902	5,178	5,161
<b>FAMILY INCOME</b>						
Less Than \$20,000	1,609	1,534	2,265	2,279	1,912	1,917
\$20,000 - \$49,999	3,030	2,985	4,220	4,123	3,307	3,290
\$50,000 - \$74,999	1,695 <sup>a</sup>	1,497	2,376 <sup>b</sup>	2,108	1,769	1,718
\$75,000 or More	2,366	2,471	3,460	3,641	2,527	2,711
<b>REGION</b>						
Northeast	1,391	1,378	2,131	2,107	1,655	1,650
Midwest	1,892	1,949	2,733	2,727	1,984	2,040
South	3,436 <sup>a</sup>	3,249	4,657	4,503	3,616	3,649
West	1,981	1,911	2,799	2,813	2,260	2,297

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents with unknown data were excluded.

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.132B Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month <sup>1</sup>		Use Cocaine Once a Month <sup>1</sup>		Have Five or More Drinks Once or Twice a Week <sup>1</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	35.0	34.0	49.6	48.8	38.1	38.4
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	34.9	30.0	49.6	44.5	42.8	38.5
Asian or Pacific Islander	37.9	38.0	43.9	44.2	47.2	51.1
Asian Only	38.4	38.0	43.0	43.1	47.8	51.2
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	35.6	34.2	54.5	54.8	48.8	48.9
White	34.7	33.9	48.9	48.0	35.3	35.5
Two or More Races	32.6	27.3	50.9	47.6	37.4	38.4
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	37.3	35.9	45.2	47.2	39.3	39.2
Not Hispanic or Latino	34.5	33.6	50.5 <sup>a</sup>	49.2	37.8	38.2
Black or African American	35.4	34.1	54.5	54.8	49.1	49.1
White	34.1	33.4	50.0 <sup>a</sup>	48.2	34.3	34.5
<b>GENDER</b>						
Male	32.6	31.1	50.1	49.2	34.0	34.9
Female	37.4	36.9	49.2	48.4	42.3	42.0
<b>FAMILY INCOME</b>						
Less Than \$20,000	35.3	34.8	50.0	51.7	41.9	43.2
\$20,000 - \$49,999	35.9	36.2	49.9	50.2	38.9	39.8
\$50,000 - \$74,999	35.3 <sup>a</sup>	31.9	49.8 <sup>b</sup>	45.2	36.7	36.4
\$75,000 or More	33.4	32.3	48.9	47.9	35.5	35.3
<b>REGION</b>						
Northeast	31.0	30.8	47.7	47.3	36.9	36.7
Midwest	33.7	34.7	48.8	48.9	35.2	36.3
South	38.7 <sup>a</sup>	36.4	52.6	50.6	40.5	40.7
West	33.4	32.0	47.4	47.3	38.0	38.2

\*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Respondents with unknown data were excluded.

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.133A Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Numbers in Thousands, 2004 and 2005**

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drugs		Alcohol		Either Illicit Drugs or Alcohol <sup>1</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	1,427	1,280	1,535	1,626	2,327	2,308
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	8	16	*	26	*	32
Asian or Pacific Islander	*	3	*	5	*	8
Asian Only	*	2	*	4	*	5
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	336	345	285	368	445	508
White	983	892	1,158	1,211	1,733	1,720
Two or More Races	34	24	32	16	41	40
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	142	182	234	252	314	373
Not Hispanic or Latino	1,285	1,098	1,301	1,374	2,013	1,934
Black or African American	334	343	285	368	443	507
White	845	722	928	969	1,425	1,372
<b>GENDER</b>						
Male	914	748	1,149	1,047	1,571	1,519
Female	513	532	386 <sup>a</sup>	579	755	789
<b>FAMILY INCOME</b>						
Less Than \$20,000	650	582	703	721	1,031	982
\$20,000 - \$49,999	452	322	481	480	779	681
\$50,000 - \$74,999	174	163	146	151	249	285
\$75,000 or More	151	214	206	274	268	360
<b>AGE GROUP</b>						
12-17	134	142	126	119	185	181
18 or Older	1,293	1,139	1,409	1,508	2,142	2,127

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.133B Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Percentages, 2004 and 2005**

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALITY FACILITY					
	Illicit Drugs		Alcohol		Either Illicit Drugs or Alcohol <sup>1</sup>	
	2004	2005	2004	2005	2004	2005
<b>TOTAL</b>	6.1	5.5	6.5	7.0	9.9	10.0
<b>RACE<sup>2</sup></b>						
American Indian or Alaska Native	2.2	4.3	*	6.8	*	8.4
Asian or Pacific Islander	*	0.6	*	0.9	*	1.5
Asian Only	*	0.5	*	0.9	*	1.1
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	12.6	12.3	10.7	13.2	16.7	18.2
White	5.1	4.7	6.0	6.3	8.9	9.0
Two or More Races	9.2	7.3	8.7	5.0	10.9	12.2
<b>HISPANIC ORIGIN AND RACE</b>						
Hispanic or Latino	4.4	5.7	7.3	7.9	9.7	11.7
Not Hispanic or Latino	6.3	5.5	6.4	6.9	9.9	9.7
Black or African American	13.0	12.5	11.1	13.3	17.3	18.4
White	5.1	4.5	5.6	6.0	8.6	8.5
<b>GENDER</b>						
Male	5.9	5.0	7.5	7.0	10.2	10.2
Female	6.4	6.4	4.8 <sup>a</sup>	7.0	9.4	9.5
<b>FAMILY INCOME</b>						
Less Than \$20,000	10.4	10.0	11.2	12.3	16.5	16.8
\$20,000 - \$49,999	5.8	4.1	6.1	6.1	10.0	8.6
\$50,000 - \$74,999	4.4	4.6	3.7	4.3	6.3	8.1
\$75,000 or More	2.8	3.6	3.8	4.7	4.9	6.1
<b>AGE GROUP</b>						
12-17	5.9	6.8	5.5	5.7	8.1	8.6
18 or Older	6.1	5.4	6.7	7.2	10.1	10.1

\*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

<sup>2</sup> These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.