

Table 7.17A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	5,381	5,127	6,272	6,299	3,803	4,213	3,615	4,080
Marijuana and Hashish	4,508	4,107	5,005	5,025	2,948	3,210	2,114	2,285
Cocaine	948	1,122	735	678	167 ^a	303	170	293
Crack	154	243	216	190	31 ^a	94	65 ^a	155
Heroin	56	29	40	37	12	28	58	41
Hallucinogens	385	397	311	350	111	156	123	184
LSD	67	38	40	35	12	17	22	14
PCP	23	20	9	13	*	4	15	11
Ecstasy	165	232	178	145	73	80	34	45
Inhalants	102	116	190	144	84	77	261	274
Nonmedical Use of Psychotherapeutics ³	1,680	1,647	1,786	1,830	1,132	1,240	1,410	1,689
Pain Relievers	1,271	1,204	1,296	1,383	740	839	1,096	1,232
OxyContin [®]	135	132	106	101	41	69	43	32
Tranquilizers	486	599	481	459	337	367	313	393
Stimulants	438	336	379	345	181	188	192	198
Methamphetamine	206	152	151	155	115	112	110	93
Sedatives	45	64	62	43	81	99	77	65
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,447	2,547	2,558	2,585	1,349	1,567	1,893 ^a	2,264

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.17B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS²	32.2	32.0	16.5	16.1	5.7	5.9	3.0 ^a	3.5
Marijuana and Hashish	27.0	25.6	13.2	12.9	4.5	4.5	1.8	1.9
Cocaine	5.7	7.0	1.9	1.7	0.3	0.4	0.1 ^a	0.3
Crack	0.9	1.5	0.6	0.5	0.0 ^a	0.1	0.1 ^a	0.1
Heroin	0.3	0.2	0.1	0.1	0.0	0.0	0.0	0.0
Hallucinogens	2.3	2.5	0.8	0.9	0.2	0.2	0.1	0.2
LSD	0.4	0.2	0.1	0.1	0.0	0.0	0.0	0.0
PCP	0.1	0.1	0.0	0.0	*	0.0	0.0	0.0
Ecstasy	1.0 ^a	1.4	0.5	0.4	0.1	0.1	0.0	0.0
Inhalants	0.6	0.7	0.5	0.4	0.1	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ³	10.1	10.3	4.7	4.7	1.7	1.7	1.2	1.4
Pain Relievers	7.6	7.5	3.4	3.5	1.1	1.2	0.9	1.1
OxyContin [®]	0.8	0.8	0.3	0.3	0.1	0.1	0.0	0.0
Tranquilizers	2.9	3.7	1.3	1.2	0.5	0.5	0.3	0.3
Stimulants	2.6	2.1	1.0	0.9	0.3	0.3	0.2	0.2
Methamphetamine	1.2	0.9	0.4	0.4	0.2	0.2	0.1	0.1
Sedatives	0.3	0.4	0.2	0.1	0.1	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	14.7	15.9	6.7	6.6	2.0	2.2	1.6 ^a	1.9

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.18A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	441	366	815	751	374	337	1,044	1,056
Marijuana and Hashish	385	324	683	626	284	255	557	525
Cocaine	53	56	54	44	11	13	7 ^a	29
Crack	11	10	2	8	4	3	*	5
Heroin	9	6	4	1	*	3	2	6
Hallucinogens	77	65	69	54	19	25	46	60
LSD	11	12	13	8	6	9	9	3
PCP	7	7	3	2	1	4	*	4
Ecstasy	32	31	29	24	5	5	11	14
Inhalants	32	20	67	65	46	38	168	190
Nonmedical Use of Psychotherapeutics ³	176	132	236	217	106	92	396	398
Pain Relievers	154 ^a	112	179	165	89	74	328	335
OxyContin [®]	32 ^b	10	18	16	8	2	8	7
Tranquilizers	40	35	47	57	19	16	56	53
Stimulants	46	45	71	61	12	17	49	48
Methamphetamine	11	21	27	18	3	8	15	19
Sedatives	3	2	4	2	5	10	13	13
ILLCIT DRUGS OTHER THAN MARIJUANA²	247 ^a	184	341	308	156	138	584	622

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.18B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	65.6	59.9	38.4	39.5	22.8	20.1	5.0	5.0
Marijuana and Hashish	57.3	53.1	32.2	32.9	17.3	15.2	2.7	2.5
Cocaine	8.0	9.2	2.6	2.3	0.7	0.7	0.0 ^a	0.1
Crack	1.6	1.6	0.1	0.4	0.2	0.2	*	0.0
Heroin	1.4	0.9	0.2	0.0	*	0.2	0.0	0.0
Hallucinogens	11.5	10.6	3.3	2.9	1.1	1.5	0.2	0.3
LSD	1.6	2.0	0.6	0.4	0.4	0.6	0.0	0.0
PCP	1.0	1.2	0.1	0.1	0.1	0.2	*	0.0
Ecstasy	4.7	5.1	1.4	1.3	0.3	0.3	0.1	0.1
Inhalants	4.7	3.4	3.1	3.4	2.8	2.2	0.8	0.9
Nonmedical Use of Psychotherapeutics ³	26.1	21.6	11.1	11.4	6.4	5.5	1.9	1.9
Pain Relievers	23.0	18.4	8.5	8.7	5.4	4.4	1.6	1.6
OxyContin [®]	4.8 ^b	1.6	0.9	0.8	0.5	0.1	0.0	0.0
Tranquilizers	5.9	5.7	2.2	3.0	1.2	0.9	0.3	0.2
Stimulants	6.9	7.4	3.3	3.2	0.7	1.0	0.2	0.2
Methamphetamine	1.7	3.5	1.3	1.0	0.2	0.5	0.1	0.1
Sedatives	0.5	0.3	0.2	0.1	0.3	0.6	0.1	0.1
ILLCIT DRUGS OTHER THAN MARIJUANA²	36.7	30.2	16.1	16.2	9.5	8.2	2.8	2.9

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.19A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	2,217	2,293	2,263	2,379	949	1,015	809	856
Marijuana and Hashish	1,960	1,909	1,883	2,038	805	857	548	585
Cocaine	408	485	207	245	29	54	47	48
Crack	46	45	19	26	5	13	11	20
Heroin	16	14	16	20	1 ^a	10	10	14
Hallucinogens	228	214	173	188	40	43	36	58
LSD	50	26	24	17	5	8	2	9
PCP	9	3	7	8	*	*	1	*
Ecstasy	109	123	84	93	22	27	16	24
Inhalants	59	60	55	43	4	8	25	35
Nonmedical Use of Psychotherapeutics ³	763	787	654	649	244	294	298	305
Pain Relievers	567	591	509	510	186	205	242	228
OxyContin [®]	63	63	52	47	18	28	9	8
Tranquilizers	253	284	190	170	75	88	47	64
Stimulants	226	197	126	139	40	62	54	36
Methamphetamine	84	76	54	66	18	31	30	21
Sedatives	21	29	14	13	4	12	10	21
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,076	1,165	893	932	285	358	365	394

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.19B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	45.7	46.2	26.9	27.6	15.3	16.4	6.4	6.7
Marijuana and Hashish	40.4	38.4	22.4	23.6	13.0	13.9	4.3	4.6
Cocaine	8.4	9.8	2.5	2.8	0.5	0.9	0.4	0.4
Crack	0.9	0.9	0.2	0.3	0.1	0.2	0.1	0.2
Heroin	0.3	0.3	0.2	0.2	0.0 ^a	0.2	0.1	0.1
Hallucinogens	4.7	4.3	2.1	2.2	0.6	0.7	0.3	0.5
LSD	1.0	0.5	0.3	0.2	0.1	0.1	0.0	0.1
PCP	0.2	0.1	0.1	0.1	*	*	0.0	*
Ecstasy	2.3	2.5	1.0	1.1	0.4	0.4	0.1	0.2
Inhalants	1.2	1.2	0.7	0.5	0.1	0.1	0.2	0.3
Nonmedical Use of Psychotherapeutics ³	15.7	15.8	7.8	7.5	3.9	4.7	2.3	2.4
Pain Relievers	11.7	11.9	6.0	5.9	3.0	3.3	1.9	1.8
OxyContin [®]	1.3	1.3	0.6	0.5	0.3	0.5	0.1	0.1
Tranquilizers	5.2	5.7	2.3	2.0	1.2	1.4	0.4	0.5
Stimulants	4.7	4.0	1.5	1.6	0.6	1.0	0.4	0.3
Methamphetamine	1.7	1.5	0.6	0.8	0.3	0.5	0.2	0.2
Sedatives	0.4	0.6	0.2	0.1	0.1	0.2	0.1	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA²	22.2	23.5	10.6	10.8	4.6	5.8	2.9	3.1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.20A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	2,724	2,468	3,194	3,170	2,480	2,860	1,762	2,169
Marijuana and Hashish	2,164	1,874	2,439	2,361	1,859	2,098	1,009	1,175
Cocaine	486	581	474	389	127	237	116	217
Crack	98	188	196	157	23	77	54	131
Heroin	31	10	21	17	11	15	46	21
Hallucinogens	80	118	68	108	52	88	42	66
LSD	5	*	2	10	*	*	11	*
PCP	8	9	*	3	*	*	13	7
Ecstasy	24 ^a	78	65	28	46	48	7	7
Inhalants	12	36	68	36	34	31	67	49
Nonmedical Use of Psychotherapeutics ³	742	728	895	964	782	854	715	986
Pain Relievers	550	501	608	708	465	560	526	669
OxyContin [®]	40	60	36	38	15	39	27	16
Tranquilizers	193	280	244	232	242	263	210	276
Stimulants	166	93	181	145	129	108	88	114
Methamphetamine	111	55	70	71	94	73	64	54
Sedatives	21	33	44	28	72	78	53	32
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,125	1,197	1,323	1,344	909	1,071	945	1,248

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.20B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS²	24.4	23.6	11.6	11.1	4.2	4.5	2.0 ^a	2.6
Marijuana and Hashish	19.4	17.9	8.9	8.3	3.2	3.3	1.2	1.4
Cocaine	4.4	5.6	1.7	1.4	0.2	0.4	0.1	0.3
Crack	0.9	1.8	0.7	0.5	0.0	0.1	0.1	0.2
Heroin	0.3	0.1	0.1	0.1	0.0	0.0	0.1	0.0
Hallucinogens	0.7	1.1	0.2	0.4	0.1	0.1	0.0	0.1
LSD	0.0	*	0.0	0.0	*	*	0.0	*
PCP	0.1	0.1	*	0.0	*	*	0.0	0.0
Ecstasy	0.2 ^a	0.7	0.2	0.1	0.1	0.1	0.0	0.0
Inhalants	0.1	0.3	0.2	0.1	0.1	0.0	0.1	0.1
Nonmedical Use of Psychotherapeutics ³	6.6	7.0	3.3	3.4	1.3	1.4	0.8 ^a	1.2
Pain Relievers	4.9	4.8	2.2	2.5	0.8	0.9	0.6	0.8
OxyContin [®]	0.4	0.6	0.1	0.1	0.0	0.1	0.0	0.0
Tranquilizers	1.7	2.7	0.9	0.8	0.4	0.4	0.2	0.3
Stimulants	1.5	0.9	0.7	0.5	0.2	0.2	0.1	0.1
Methamphetamine	1.0	0.5	0.3	0.2	0.2	0.1	0.1	0.1
Sedatives	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA²	10.1	11.4	4.8	4.7	1.6	1.7	1.1	1.5

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.