

Table 7.21A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	10,117	10,207	1,866	2,008	5,340	5,432	1,749	2,072
Marijuana and Hashish	8,429	8,472	1,339	1,456	4,033	3,869	775	829
Cocaine	1,476	1,732	128	217	375	371	42	76
Crack	369	426	59	129	33	101	6	26
Heroin	90	83	55	33	18	12	*	8
Hallucinogens	602	672	66	113	204	232	57	71
LSD	100	62	17	9	18	29	5	5
PCP	32	32	*	10	*	4	14	*
Ecstasy	287	349	24	28	129	108	10	17
Inhalants	233	218	28	49	143	119	232	225
Nonmedical Use of Psychotherapeutics ²	3,048	2,966	638	635	1,550	1,751	771 ^a	1,054
Pain Relievers	2,270	2,215	457	419	1,037	1,211	639	813
OxyContin [®]	252	243	30	27	30	59	13	*
Tranquilizers	850	949	201	216	453	476	111	176
Stimulants	701	626	116	112	297	243	75	86
Methamphetamine	374	345	71	64	99	74	39	29
Sedatives	129	123	36	21	60	83	41	45
ILLCIT DRUGS OTHER THAN MARIJUANA¹	4,354	4,485	831	891	2,000	2,214	1,062 ^a	1,373

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.21B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	25.0	24.9	9.6	10.3	6.6	6.4	1.7 ^a	2.1
Marijuana and Hashish	20.8	20.7	6.9	7.4	5.0	4.5	0.8	0.8
Cocaine	3.6	4.2	0.7	1.1	0.5	0.4	0.0	0.1
Crack	0.9	1.0	0.3	0.7	0.0	0.1	0.0	0.0
Heroin	0.2	0.2	0.3	0.2	0.0	0.0	*	0.0
Hallucinogens	1.5	1.6	0.3	0.6	0.3	0.3	0.1	0.1
LSD	0.2	0.2	0.1	0.0	0.0	0.0	0.0	0.0
PCP	0.1	0.1	*	0.0	*	0.0	0.0	*
Ecstasy	0.7	0.9	0.1	0.1	0.2	0.1	0.0	0.0
Inhalants	0.6	0.5	0.1	0.3	0.2	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ²	7.5	7.2	3.3	3.2	1.9	2.1	0.8 ^a	1.1
Pain Relievers	5.6	5.4	2.3	2.1	1.3	1.4	0.6	0.8
OxyContin [®]	0.6	0.6	0.2	0.1	0.0	0.1	0.0	*
Tranquilizers	2.1	2.3	1.0	1.1	0.6	0.6	0.1	0.2
Stimulants	1.7	1.5	0.6	0.6	0.4	0.3	0.1	0.1
Methamphetamine	0.9	0.8	0.4	0.3	0.1	0.1	0.0	0.0
Sedatives	0.3	0.3	0.2	0.1	0.1	0.1	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	10.8	11.0	4.3	4.5	2.5	2.6	1.1 ^a	1.4

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.22A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	1,067 ^a	931	356	345	563	524	688	711
Marijuana and Hashish	922	837	293	258	430	368	264	267
Cocaine	106	94	7	16	13	18	*	13
Crack	12	17	*	5	4	4	*	*
Heroin	8	8	1	1	5	1	2	4
Hallucinogens	135	107	25	32	30	37	21	28
LSD	26	18	3	1	3	11	5	2
PCP	9	11	1	3	2	1	*	1
Ecstasy	57	49	9	8	9	11	2	6
Inhalants	99	71	24	38	45	52	144	152
Nonmedical Use of Psychotherapeutics ²	355	299	84	83	163	142	312	315
Pain Relievers	295	247	60	69	128	104	268	266
OxyContin [®]	54 ^a	26	5	4	4	1	2	3
Tranquilizers	76	97	25	18	29 ^a	11	31	35
Stimulants	98	93	14	15	31	30	35	33
Methamphetamine	39	37	2	4	2	11	13	14
Sedatives	9	8	1	6	4	6	13	7
ILLCIT DRUGS OTHER THAN MARIJUANA¹	514 ^a	415	121	142	230	216	463	481

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.22B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS¹	58.8	57.7	30.1	30.8	21.5	20.3	3.5	3.5
Marijuana and Hashish	50.8	51.9	24.7	23.0	16.4	14.3	1.3	1.3
Cocaine	5.8	5.8	0.6	1.5	0.5	0.7	*	0.1
Crack	0.7	1.0	*	0.4	0.1	0.2	*	*
Heroin	0.4	0.5	0.1	0.1	0.2	0.0	0.0	0.0
Hallucinogens	7.5	6.7	2.1	2.9	1.1	1.4	0.1	0.1
LSD	1.4	1.1	0.3	0.1	0.1	0.4	0.0	0.0
PCP	0.5	0.7	0.1	0.2	0.1	0.1	*	0.0
Ecstasy	3.1	3.0	0.7	0.7	0.4	0.4	0.0	0.0
Inhalants	5.4	4.4	2.0	3.4	1.7	2.0	0.7	0.8
Nonmedical Use of Psychotherapeutics ²	19.5	18.5	7.1	7.4	6.2	5.5	1.6	1.6
Pain Relievers	16.3	15.3	5.0	6.2	4.9	4.0	1.4	1.3
OxyContin [®]	3.0 ^a	1.6	0.5	0.4	0.1	0.0	0.0	0.0
Tranquilizers	4.2	6.0	2.1	1.6	1.1 ^a	0.4	0.2	0.2
Stimulants	5.4	5.8	1.2	1.3	1.2	1.2	0.2	0.2
Methamphetamine	2.2	2.3	0.2	0.4	0.1	0.4	0.1	0.1
Sedatives	0.5	0.5	0.1	0.5	0.2	0.2	0.1	0.0
ILLICIT DRUGS OTHER THAN MARIJUANA¹	28.3	25.7	10.2	12.6	8.8	8.4	2.4	2.4

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.23A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS¹	3,968	4,092	442	473	1,462	1,595	367	383
Marijuana and Hashish	3,474	3,506	362	387	1,174	1,297	185	198
Cocaine	558	660	40	38	87	124	7	10
Crack	62	70	11	15	8	14	*	5
Heroin	31	44	8	11	2	1	2	3
Hallucinogens	352	338	18	32	89	107	18	26
LSD	66	37	2	7	15	14	*	2
PCP	15	11	*	*	*	*	1	*
Ecstasy	178	192	8	13	37	51	8	11
Inhalants	94	88	4	12	24	23	21	23
Nonmedical Use of Psychotherapeutics ²	1,274	1,282	131	136	387	448	168	169
Pain Relievers	980	984	96	95	282	322	146	133
OxyContin [®]	123	113	7	8	10	25	1	*
Tranquilizers	394	415	27	42	124	127	20	22
Stimulants	326	311	40	22	66	87	14	14
Methamphetamine	137	148	25	21	19	25	5	*
Sedatives	28	44	4	11	11	9	6	10
ILLICIT DRUGS OTHER THAN MARIJUANA¹	1,735	1,860	167	185	519	596	198	210

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.23B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS¹	39.8	41.0	16.0	17.6	15.4	16.3	3.7	3.8
Marijuana and Hashish	34.9	35.2	13.1	14.3	12.3	13.2	1.9	2.0
Cocaine	5.6	6.6	1.5	1.4	0.9	1.3	0.1	0.1
Crack	0.6	0.7	0.4	0.6	0.1	0.1	*	0.0
Heroin	0.3	0.4	0.3	0.4	0.0	0.0	0.0	0.0
Hallucinogens	3.5	3.4	0.6	1.2	0.9	1.1	0.2	0.3
LSD	0.7	0.4	0.1	0.3	0.2	0.1	*	0.0
PCP	0.2	0.1	*	*	*	*	0.0	*
Ecstasy	1.8	1.9	0.3	0.5	0.4	0.5	0.1	0.1
Inhalants	0.9	0.9	0.2	0.4	0.3	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ²	12.8	12.9	4.7	5.0	4.1	4.6	1.7	1.7
Pain Relievers	9.8	9.9	3.5	3.5	3.0	3.3	1.5	1.3
OxyContin [®]	1.2	1.1	0.3	0.3	0.1	0.3	0.0	*
Tranquilizers	4.0	4.2	1.0	1.6	1.3	1.3	0.2	0.2
Stimulants	3.3	3.1	1.4	0.8	0.7	0.9	0.1	0.1
Methamphetamine	1.4	1.5	0.9	0.8	0.2	0.3	0.0	*
Sedatives	0.3	0.4	0.1	0.4	0.1	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA¹	17.4	18.6	6.0	6.9	5.5	6.1	2.0	2.1

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.24A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	5,082	5,184	1,068	1,191	3,315	3,313	694	978
Marijuana and Hashish	4,033	4,129	684	812	2,429	2,205	325	363
Cocaine	813	978	81	163	275	229	35	53
Crack	295	339	49	109	21	83	6	21
Heroin	51	31	46	21	11	10	*	*
Hallucinogens	115	226	23	49	85	88	18	17
LSD	8	7	11	*	*	*	*	*
PCP	8	9	*	7	*	*	13	*
Ecstasy	52	108	7	7	83	46	*	*
Inhalants	40	59	*	*	74	44	67	49
Nonmedical Use of Psychotherapeutics ²	1,419	1,385	423	416	1,000	1,161	292 ^a	569
Pain Relievers	996	984	301	255	627	784	225	414
OxyContin [®]	75	104	17	14	16	34	9	*
Tranquilizers	380	437	149	156	299	337	61	120
Stimulants	277	221	62	75	200	126	26	39
Methamphetamine	198	160	43	39	78	39	21	15
Sedatives	92	71	32	4	45	69	22	28
ILLCIT DRUGS OTHER THAN MARIJUANA¹	2,105	2,210	544	565	1,251	1,403	401 ^a	682

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.24B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	17.7	17.7	6.9	7.5	4.8	4.6	1.0 ^a	1.4
Marijuana and Hashish	14.1	14.1	4.4	5.1	3.6	3.0	0.5	0.5
Cocaine	2.8	3.3	0.5	1.0	0.4	0.3	0.0	0.1
Crack	1.0	1.2	0.3	0.7	0.0	0.1	0.0	0.0
Heroin	0.2	0.1	0.3	0.1	0.0	0.0	*	*
Hallucinogens	0.4	0.8	0.2	0.3	0.1	0.1	0.0	0.0
LSD	0.0	0.0	0.1	*	*	*	*	*
PCP	0.0	0.0	*	0.0	*	*	0.0	*
Ecstasy	0.2	0.4	0.0	0.0	0.1	0.1	*	*
Inhalants	0.1	0.2	*	*	0.1	0.1	0.1	0.1
Nonmedical Use of Psychotherapeutics ²	5.0	4.7	2.7	2.6	1.5	1.6	0.4 ^b	0.8
Pain Relievers	3.5	3.4	1.9	1.6	0.9	1.1	0.3 ^a	0.6
OxyContin [®]	0.3	0.4	0.1	0.1	0.0	0.0	0.0	*
Tranquilizers	1.3	1.5	1.0	1.0	0.4	0.5	0.1	0.2
Stimulants	1.0	0.8	0.4	0.5	0.3	0.2	0.0	0.1
Methamphetamine	0.7	0.5	0.3	0.2	0.1	0.1	0.0	0.0
Sedatives	0.3	0.2	0.2	0.0	0.1	0.1	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA¹	7.3	7.5	3.5	3.6	1.8	1.9	0.6 ^a	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.25A Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Numbers in Thousands, 2004 and 2005

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	19,071	19,720	2,674	2,511	6,239	6,543	10,159	10,666
Heavy Alcohol Use ¹	5,381	5,127	441	366	2,217	2,293	2,724	2,468
Binge Alcohol Use But Not Heavy Use ¹	6,272	6,299	815	751	2,263	2,379	3,194	3,170
Alcohol Use But Not Binge Use ¹	3,803	4,213	374	337	949	1,015	2,480	2,860
No Alcohol Use	3,615	4,080	1,044	1,056	809	856	1,762	2,169
PAST MONTH CIGARETTE USE	11,983	12,216	1,423	1,276	4,410	4,565	6,150	6,375
Heavy Alcohol Use ¹	4,076	3,789	342	297	1,810	1,793	1,923	1,699
Binge Alcohol Use But Not Heavy Use ¹	4,025	4,174	540	480	1,582	1,699	1,903	1,994
Alcohol Use But Not Binge Use ¹	2,015	2,244	184	154	576	600	1,256	1,490
No Alcohol Use	1,866	2,008	356	345	442	473	1,068	1,191
NO PAST MONTH CIGARETTE USE	7,088	7,504	1,251	1,235	1,829	1,978	4,009	4,291
Heavy Alcohol Use ¹	1,305	1,338	98	70	407	500	800	768
Binge Alcohol Use But Not Heavy Use ¹	2,247	2,126	275	271	682	680	1,291	1,175
Alcohol Use But Not Binge Use ¹	1,787	1,968	190	183	374	415	1,224	1,370
No Alcohol Use	1,749	2,072	688	711	367	383	694	978

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.25B Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Percentages, 2004 and 2005

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	7.9	8.1	10.6	9.9	19.4	20.1	5.5	5.8
Heavy Alcohol Use ¹	32.2	32.0	65.6	59.9	45.7	46.2	24.4	23.6
Binge Alcohol Use But Not Heavy Use ¹	16.5	16.1	38.4	39.5	26.9	27.6	11.6	11.1
Alcohol Use But Not Binge Use ¹	5.7	5.9	22.8	20.1	15.3	16.4	4.2	4.5
No Alcohol Use	3.0 ^a	3.5	5.0	5.0	6.4	6.7	2.0 ^a	2.6
PAST MONTH CIGARETTE USE	20.0	20.2	47.5	46.7	34.6	36.0	13.9	14.1
Heavy Alcohol Use ¹	39.9	39.0	70.1	70.9	51.8	52.3	30.9	28.9
Binge Alcohol Use But Not Heavy Use ¹	24.8	24.9	57.0	56.7	35.5	37.2	17.5	17.6
Alcohol Use But Not Binge Use ¹	14.4	15.5	48.5	44.2	28.5	30.3	10.9	12.3
No Alcohol Use	9.6	10.3	30.1	30.8	16.0	17.6	6.9	7.5
NO PAST MONTH CIGARETTE USE	3.9	4.1	5.6	5.5	9.4	10.0	2.9	3.1
Heavy Alcohol Use ¹	20.1	21.2	53.8 ^b	36.1	30.1	32.5	16.2	16.7
Binge Alcohol Use But Not Heavy Use ¹	10.3	9.5	23.4	25.7	17.2	16.7	7.8	6.8
Alcohol Use But Not Binge Use ¹	3.4	3.5	15.0	13.8	8.9	9.9	2.6	2.7
No Alcohol Use	1.7 ^a	2.1	3.5	3.5	3.7	3.8	1.0 ^a	1.4

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.