

Table 7.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	10,361	10,987
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	2,876	3,005	4,278	4,677
Cigars	7,779	7,617	5,948	6,023
Pipe Tobacco	962	1,105	873	1,085
ALCOHOL	40,437	40,938	80,497 ^a	85,090
Binge Alcohol Use ²	26,471	26,486	28,254	28,604
Heavy Alcohol Use ²	10,211	9,715	6,478	6,320

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	5.7	6.0
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	4.8	5.0	2.4	2.6
Cigars	13.0	12.6	3.3	3.3
Pipe Tobacco	1.6	1.8	0.5	0.6
ALCOHOL	67.5	67.6	44.6 ^b	46.6
Binge Alcohol Use ²	44.2	43.8	15.6	15.7
Heavy Alcohol Use ²	17.0	16.1	3.6	3.5

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.27A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	637	591
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	307	266	264	258
Cigars	814	720	387	344
Pipe Tobacco	134	112	44	46
ALCOHOL	1,815 ^a	1,614	2,621	2,577
Binge Alcohol Use ²	1,436 ^a	1,265	1,358	1,248
Heavy Alcohol Use ²	489	418	183	193

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.27B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	2.9	2.6
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	10.3	9.7	1.2	1.1
Cigars	27.2	26.3	1.7	1.5
Pipe Tobacco	4.5	4.1	0.2	0.2
ALCOHOL	60.5	59.1	11.8	11.4
Binge Alcohol Use ²	47.9	46.3	6.1	5.5
Heavy Alcohol Use ²	16.3	15.3	0.8	0.9

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.28A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	1,628	1,716
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,115	1,077	474 ^a	586
Cigars	2,866	2,756	1,237	1,139
Pipe Tobacco	324	336	72 ^b	135
ALCOHOL	9,964	9,972	9,510	9,811
Binge Alcohol Use ²	7,947	7,990	5,324	5,607
Heavy Alcohol Use ²	3,495	3,427	1,353	1,540

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.28B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	8.4	8.7
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	8.8	8.5	2.4 ^a	3.0
Cigars	22.5	21.8	6.4	5.7
Pipe Tobacco	2.5	2.7	0.4 ^a	0.7
ALCOHOL	78.3	78.7	48.9	49.5
Binge Alcohol Use ²	62.4	63.1	27.4	28.3
Heavy Alcohol Use ²	27.5	27.1	7.0	7.8

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.29A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	8,097	8,680
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,454	1,662	3,540	3,833
Cigars	4,099	4,141	4,324	4,540
Pipe Tobacco	505	656	756	904
ALCOHOL	28,657	29,352	68,366 ^a	72,702
Binge Alcohol Use ²	17,089	17,230	21,572	21,750
Heavy Alcohol Use ²	6,228	5,871	4,942	4,587

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.29B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2004 and 2005

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
TOBACCO PRODUCTS¹	N/A	N/A	5.8	6.2
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3.3	3.7	2.5	2.7
Cigars	9.3	9.2	3.1	3.2
Pipe Tobacco	1.1	1.5	0.5	0.6
ALCOHOL	64.9	65.0	49.2 ^b	51.8
Binge Alcohol Use ²	38.7	38.2	15.5	15.5
Heavy Alcohol Use ²	14.1	13.0	3.6	3.3

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.