

**Table 7.30A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	11,703	11,533	19,180	19,761	17,023	17,634	22,351	22,591
Cigarettes	10,211	9,715	16,260	16,770	13,965	14,452	19,459	19,594
Smokeless Tobacco	1,728	1,903	1,900	2,173	1,383	1,376	2,144	2,230
Cigars	3,112	3,076	4,530	4,447	3,731	3,536	2,354	2,581
Pipe Tobacco	292	332	475	602	412	443	656	813

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.30B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	70.1	71.9	50.4	50.6	25.7	24.9	18.7	19.3
Cigarettes	61.2	60.6	42.7	42.9	21.1	20.4	16.3	16.7
Smokeless Tobacco	10.4	11.9	5.0	5.6	2.1	1.9	1.8	1.9
Cigars	18.6	19.2	11.9	11.4	5.6	5.0	2.0	2.2
Pipe Tobacco	1.7	2.1	1.2	1.5	0.6	0.6	0.5	0.7

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.31A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	564 <sup>a</sup>	474	1,136	1,023	476	430	1,458	1,398
Cigarettes	489	418	947	847	379	349	1,184	1,119
Smokeless Tobacco	125	108	193	172	39	28	213	215
Cigars	242	226	434 <sup>a</sup>	360	160	120	366	358
Pipe Tobacco	37	37	55	50	17	23	68	49

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.31B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	84.0 <sup>a</sup>	77.5	53.5	53.8	29.0	25.6	7.0	6.6
Cigarettes	72.8	68.4	44.6	44.5	23.1	20.8	5.7	5.3
Smokeless Tobacco	18.6	17.7	9.1	9.0	2.4	1.7	1.0	1.0
Cigars	36.0	37.0	20.4	18.9	9.7 <sup>a</sup>	7.2	1.8	1.7
Pipe Tobacco	5.5	6.0	2.6	2.6	1.0	1.3	0.3	0.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.32A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	3,900	3,882	5,022	5,256	2,315	2,256	3,122	2,988
Cigarettes	3,495	3,427	4,452	4,564	2,017	1,982	2,767	2,694
Smokeless Tobacco	720	745	496	570	144	118	228	231
Cigars	1,375	1,215	1,470	1,508	601	538	657	633
Pipe Tobacco	164	141	116 <sup>a</sup>	181	56	65	60	85

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.32B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	80.4	78.2	59.6	60.9	37.3	36.5	24.5	23.5
Cigarettes	72.1 <sup>a</sup>	69.0	52.9	52.9	32.5	32.0	21.8	21.2
Smokeless Tobacco	14.9	15.0	5.9	6.6	2.3	1.9	1.8	1.8
Cigars	28.4 <sup>b</sup>	24.5	17.5	17.5	9.7	8.7	5.2	5.0
Pipe Tobacco	3.4	2.8	1.4 <sup>a</sup>	2.1	0.9	1.0	0.5	0.7

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.33A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	7,238	7,177	13,021	13,482	14,232	14,948	17,771	18,205
Cigarettes	6,228	5,871	10,861	11,360	11,569	12,122	15,509	15,781
Smokeless Tobacco	883	1,049	1,210	1,431	1,199	1,230	1,702	1,784
Cigars	1,496	1,634	2,627	2,579	2,970	2,878	1,330	1,590
Pipe Tobacco	91	154	303	371	339	356	528	679

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.33B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2004 and 2005**

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>2</sup></b>	64.8	68.6	47.4	47.3	24.4	23.7	20.6	21.8
Cigarettes	55.8	56.1	39.5	39.8	19.8	19.2	18.0	18.9
Smokeless Tobacco	7.9	10.0	4.4	5.0	2.1	2.0	2.0	2.1
Cigars	13.4	15.6	9.6	9.0	5.1	4.6	1.5	1.9
Pipe Tobacco	0.8	1.5	1.1	1.3	0.6	0.6	0.6	0.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.