

Table 7.53A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2005

| Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹ | Total | AGE GROUP | | |
|--|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 1,840 | 265 | 532 | 1,044 |
| From More Than One Doctor | 182 | 38 | 53 | 90 |
| Wrote Fake Prescription | 21 | 2 | 13 | 6 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 40 | 8 | 9 | 24 |
| From Friend or Relative for Free | 6,533 | 685 | 2,223 | 3,626 |
| Bought from Friend or Relative | 911 | 144 | 444 | 322 |
| Took from Friend or Relative without Asking | 459 | 148 | 159 | 152 |
| Bought from Drug Dealer or Other Stranger | 469 | 65 | 200 | 204 |
| Bought on the Internet | 84 | 2 | 15 | 67 |
| Some Other Way | 390 | 92 | 124 | 175 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.53B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2005

| Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹ | Total | AGE GROUP | | |
|--|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 16.8 | 18.3 | 14.1 | 18.3 |
| From More Than One Doctor | 1.7 | 2.6 | 1.4 | 1.6 |
| Wrote Fake Prescription | 0.2 | 0.1 | 0.3 | 0.1 |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.4 | 0.5 | 0.2 | 0.4 |
| From Friend or Relative for Free | 59.8 | 47.3 | 58.9 | 63.5 |
| Bought from Friend or Relative | 8.3 | 10.0 | 11.8 | 5.6 |
| Took from Friend or Relative without Asking | 4.2 | 10.2 | 4.2 | 2.7 |
| Bought from Drug Dealer or Other Stranger | 4.3 | 4.5 | 5.3 | 3.6 |
| Bought on the Internet | 0.8 | 0.1 | 0.4 | 1.2 |
| Some Other Way | 3.6 | 6.3 | 3.3 | 3.1 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.54A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2005

| Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 451 | 29 | 108 | 313 |
| From More Than One Doctor | 81 | 3 | 14 | 63 |
| Wrote Fake Prescription | 2 | 0 | 1 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 15 | 7 | 5 | 3 |
| From Friend or Relative for Free | 3,265 | 207 | 988 | 2,070 |
| Bought from Friend or Relative | 514 | 78 | 269 | 167 |
| Took from Friend or Relative without Asking | 171 | 37 | 77 | 57 |
| Bought from Drug Dealer or Other Stranger | 311 | 49 | 159 | 103 |
| Bought on the Internet | 39 | * | 10 | 29 |
| Some Other Way | 102 | 16 | 39 | 47 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.54B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2005

| Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 9.1 | 6.8 | 6.5 | 11.0 |
| From More Than One Doctor | 1.6 | 0.8 | 0.8 | 2.2 |
| Wrote Fake Prescription | 0.0 | 0.1 | 0.1 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.3 | 1.6 | 0.3 | 0.1 |
| From Friend or Relative for Free | 66.0 | 48.5 | 59.1 | 72.6 |
| Bought from Friend or Relative | 10.4 | 18.2 | 16.1 | 5.9 |
| Took from Friend or Relative without Asking | 3.5 | 8.7 | 4.6 | 2.0 |
| Bought from Drug Dealer or Other Stranger | 6.3 | 11.6 | 9.5 | 3.6 |
| Bought on the Internet | 0.8 | * | 0.6 | 1.0 |
| Some Other Way | 2.1 | 3.8 | 2.3 | 1.6 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.55A Source Where Prescription Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2005

| Source for Most Recent Nonmedical Use among Past Year Prescription Stimulant Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 99 | 27 | 32 | 40 |
| From More Than One Doctor | 37 | 7 | 6 | 24 |
| Wrote Fake Prescription | 9 | * | 7 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 10 | * | 7 | * |
| From Friend or Relative for Free | 1,108 | 199 | 482 | 427 |
| Bought from Friend or Relative | 250 | 38 | 179 | 32 |
| Took from Friend or Relative without Asking | 99 | 37 | 31 | 32 |
| Bought from Drug Dealer or Other Stranger | 126 | 20 | 53 | 53 |
| Bought on the Internet | 139 | 7 | 14 | * |
| Some Other Way | 49 | 10 | 28 | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Past Year Prescription Stimulant Use refers only to nonmedical prescription stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.55B Source Where Prescription Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2005

| Source for Most Recent Nonmedical Use among Past Year Prescription Stimulant Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 5.1 | 7.7 | 3.8 | 5.4 |
| From More Than One Doctor | 1.9 | 2.0 | 0.7 | 3.2 |
| Wrote Fake Prescription | 0.5 | * | 0.8 | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 0.5 | * | 0.8 | * |
| From Friend or Relative for Free | 57.6 | 57.0 | 57.5 | 57.9 |
| Bought from Friend or Relative | 13.0 | 10.9 | 21.4 | 4.3 |
| Took from Friend or Relative without Asking | 5.2 | 10.5 | 3.6 | 4.4 |
| Bought from Drug Dealer or Other Stranger | 6.5 | 5.7 | 6.3 | 7.1 |
| Bought on the Internet | 7.2 | 2.0 | 1.7 | * |
| Some Other Way | 2.5 | 2.8 | 3.3 | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Past Year Prescription Stimulant Use refers only to nonmedical prescription stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.56A Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2005

| Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹ | Total | AGE GROUP | | |
|--|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From Friend or Relative for Free | 589 | 92 | 239 | 257 |
| Bought from Friend or Relative | 348 | 22 | 97 | 229 |
| Took from Friend or Relative without Asking | 32 | 6 | 6 | * |
| Bought from Drug Dealer or Other Stranger | 210 | * | 92 | 98 |
| Bought on the Internet | 19 | * | 9 | * |
| Some Other Way | 36 | 7 | 10 | 19 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.56B Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2005

| Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹ | Total | AGE GROUP | | |
|--|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From Friend or Relative for Free | 47.7 | 62.2 | 52.8 | 40.6 |
| Bought from Friend or Relative | 28.2 | 14.7 | 21.4 | 36.2 |
| Took from Friend or Relative without Asking | 2.6 | 3.9 | 1.4 | * |
| Bought from Drug Dealer or Other Stranger | 17.1 | * | 20.2 | 15.4 |
| Bought on the Internet | 1.5 | * | 2.0 | * |
| Some Other Way | 2.9 | 5.0 | 2.2 | 3.0 |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.57A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Numbers in Thousands, 2005**

| Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 73 | * | 18 | * |
| From More Than One Doctor | 18 | * | * | * |
| Wrote Fake Prescription | 1 | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 10 | * | 8 | * |
| From Friend or Relative for Free | 462 | * | 80 | * |
| Bought from Friend or Relative | 40 | * | 20 | * |
| Took from Friend or Relative without Asking | 38 | * | * | * |
| Bought from Drug Dealer or Other Stranger | 39 | * | 16 | * |
| Bought on the Internet | * | * | * | * |
| Some Other Way | 16 | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.57B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2005

| Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹ | Total | AGE GROUP | | |
|---|-------|-----------|-------|-------------|
| | | 12-17 | 18-25 | 26 or Older |
| From One Doctor | 10.5 | * | 10.5 | * |
| From More Than One Doctor | 2.5 | * | * | * |
| Wrote Fake Prescription | 0.1 | * | * | * |
| Stole from Doctor's Office, Clinic, Hospital, or Pharmacy | 1.4 | * | 4.6 | * |
| From Friend or Relative for Free | 66.3 | * | 46.9 | * |
| Bought from Friend or Relative | 5.7 | * | 11.7 | * |
| Took from Friend or Relative without Asking | 5.5 | * | * | * |
| Bought from Drug Dealer or Other Stranger | 5.5 | * | 9.5 | * |
| Bought on the Internet | * | * | * | * |
| Some Other Way | 2.4 | * | * | * |

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.