

Table 7.68A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
ILLCIT DRUGS²	6,300	5,963	105	97	6,152	5,836
Marijuana and Hashish	4,575	4,272	86	71	4,452	4,179
Cocaine	601	652	8	7	587	640
Crack	116	167	1	2	115	165
Heroin	34	38	1	3	32	35
Hallucinogens	354 ^a	274	6	6	344 ^a	263
LSD	35	29	0	3	35	26
PCP	15	9	2	*	14	8
Ecstasy	223 ^a	156	4	3	217 ^a	150
Inhalants	131	125	1	2	130	121
Nonmedical Use of Psychotherapeutics ³	2,329	2,348	30	33	2,285	2,302
Pain Relievers	1,630	1,674	21	29	1,596	1,634
OxyContin [®]	--	126	--	1	--	124
Tranquilizers	692	678	5	6	684	668
Stimulants	492	520	7	3	483	513
Methamphetamine	207	230	1	1	205	228
Sedatives	124	101	2	1	123	100
ILLCIT DRUGS OTHER THAN MARIJUANA²	2,961	2,929	35	40	2,907	2,872

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.68B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
ILLCIT DRUGS²	10.2	9.6	4.3	3.9	10.4	9.9
Marijuana and Hashish	7.4 ^a	6.9	3.5	2.8	7.6	7.1
Cocaine	1.0	1.1	0.3	0.3	1.0	1.1
Crack	0.2	0.3	0.0	0.1	0.2	0.3
Heroin	0.1	0.1	0.0	0.1	0.1	0.1
Hallucinogens	0.6 ^a	0.4	0.3	0.2	0.6 ^a	0.4
LSD	0.1	0.0	0.0	0.1	0.1	0.0
PCP	0.0	0.0	0.1	*	0.0	0.0
Ecstasy	0.4 ^a	0.3	0.1	0.1	0.4 ^a	0.3
Inhalants	0.2	0.2	0.0	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ³	3.8	3.8	1.2	1.3	3.9	3.9
Pain Relievers	2.6	2.7	0.9	1.2	2.7	2.8
OxyContin [®]	--	0.2	--	0.1	--	0.2
Tranquilizers	1.1	1.1	0.2	0.3	1.2	1.1
Stimulants	0.8	0.8	0.3	0.1	0.8	0.9
Methamphetamine	0.3	0.4	0.0	0.1	0.3	0.4
Sedatives	0.2	0.2	0.1	0.0	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA²	4.8	4.7	1.4	1.6	4.9	4.9

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.69A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	6,300	5,963	105	97	6,152	5,836
AGE						
15-17	994 ^a	908	12	9	975 ^a	884
18-25	2,573	2,499	71	63	2,476	2,425
26-44	2,733	2,556	22	24	2,701	2,526
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5,632 ^a	5,220	90	82	5,502 ^a	5,110
White	4,534 ^a	4,185	62	58	4,447 ^a	4,105
Black or African American	780	750	24	22	746	723
American Indian or Alaska Native	60	50	*	*	59	49
Native Hawaiian or Other Pacific Islander	24	*	*	*	23	*
Asian	131 ^a	80	*	*	128	79
Two or More Races	102	129	*	*	99	127
Hispanic or Latino	668	743	15	15	651	725
TRIMESTER²						
First	N/A	N/A	54	50	N/A	N/A
Second	N/A	N/A	31	26	N/A	N/A
Third	N/A	N/A	18	21	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.69B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	10.2	9.6	4.3	3.9	10.4	9.9
AGE						
15-17	16.5 ^b	14.8	12.8	12.3	16.5 ^b	14.7
18-25	16.4	15.5	7.5	7.0	16.9	16.1
26-44	6.8	6.4	1.6	1.6	7.0	6.7
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	10.8 ^a	10.0	4.7	4.1	11.0 ^a	10.2
White	11.4	10.7	4.4	3.9	11.6	10.9
Black or African American	9.5	8.8	8.0	6.8	9.4	8.9
American Indian or Alaska Native	15.4	12.9	*	*	16.3	13.0
Native Hawaiian or Other Pacific Islander	12.6	*	*	*	12.8	*
Asian	4.3 ^a	2.5	*	*	4.4 ^a	2.6
Two or More Races	15.4	19.5	*	*	15.9	19.8
Hispanic or Latino	7.2	7.7	3.0	3.0	7.4	7.9
TRIMESTER²						
First	N/A	N/A	7.7	7.0	N/A	N/A
Second	N/A	N/A	3.2	3.2	N/A	N/A
Third	N/A	N/A	2.3	2.3	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.70A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	4,575	4,272	86	71	4,452	4,179
AGE						
15-17	724	679	9	8	709	662
18-25	2,086 ^a	1,937	60	45	2,004	1,883
26-44	1,765	1,657	17	18	1,739	1,634
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	4,171 ^a	3,818	74	60	4,061 ^a	3,738
White	3,374 ^a	3,074	52	46	3,299 ^a	3,013
Black or African American	594	547	18	12	565	531
American Indian or Alaska Native	44	40	*	*	43	39
Native Hawaiian or Other Pacific Islander	10	9	*	*	10	9
Asian	78	57	*	*	74	57
Two or More Races	72	90	*	*	70	89
Hispanic or Latino	404	454	11	11	391	441
TRIMESTER²						
First	N/A	N/A	43	36	N/A	N/A
Second	N/A	N/A	25	22	N/A	N/A
Third	N/A	N/A	16	13	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.70B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	7.4 ^a	6.9	3.5	2.8	7.6	7.1
AGE						
15-17	12.0	11.0	9.5	10.7	12.0	11.0
18-25	13.3 ^b	12.0	6.3	4.9	13.7 ^b	12.5
26-44	4.4	4.2	1.2	1.2	4.5	4.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	8.0 ^a	7.3	3.9	3.0	8.1 ^a	7.5
White	8.5	7.8	3.7	3.1	8.6	8.0
Black or African American	7.2	6.4	6.2	3.8	7.1	6.5
American Indian or Alaska Native	11.3	10.3	*	*	12.0	10.4
Native Hawaiian or Other Pacific Islander	5.2	4.3	*	*	5.4	4.3
Asian	2.5	1.8	*	*	2.6	1.9
Two or More Races	11.0	13.7	*	*	11.2	13.9
Hispanic or Latino	4.3	4.7	2.2	2.2	4.5	4.8
TRIMESTER²						
First	N/A	N/A	6.1	5.0	N/A	N/A
Second	N/A	N/A	2.5	2.6	N/A	N/A
Third	N/A	N/A	2.1	1.5	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.71A Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOBACCO PRODUCTS²	19,288	18,627	459	423	18,707	18,124
Cigarettes	18,655	17,948	438	413	18,110	17,455
Smokeless Tobacco	184	184	7	4	178	179
Cigars	1,876	1,926	35	28	1,815	1,889
Pipe Tobacco	160	206	8	3	152	203
ALCOHOL	31,647	31,824	238	300	31,246	31,376
Binge Alcohol Use ³	13,849	13,939	99	96	13,676	13,782
Heavy Alcohol Use ³	3,166	3,158	16	18	3,140	3,132

*Low precision; no estimate reported.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.71B Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOBACCO PRODUCTS²	31.3 ^a	30.1	18.9	17.0	31.7 ^a	30.7
Cigarettes	30.2 ^a	29.0	18.0	16.6	30.7 ^a	29.6
Smokeless Tobacco	0.3	0.3	0.3	0.2	0.3	0.3
Cigars	3.0	3.1	1.4	1.1	3.1	3.2
Pipe Tobacco	0.3	0.3	0.3	0.1	0.3	0.3
ALCOHOL	51.3	51.4	9.8	12.1	53.0	53.1
Binge Alcohol Use ³	22.4	22.5	4.1	3.9	23.2	23.3
Heavy Alcohol Use ³	5.1	5.1	0.7	0.7	5.3	5.3

*Low precision; no estimate reported.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.72A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	18,655	17,948	438	413	18,110	17,455
AGE						
15-17	1,257 ^a	1,149	25	17	1,223 ^a	1,117
18-25	5,739	5,675	263	240	5,433	5,400
26-44	11,659	11,123	149	157	11,454	10,939
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	16,915 ^a	16,050	403	377	16,420 ^a	15,606
White	14,035 ^b	13,241	356	319	13,604 ^a	12,868
Black or African American	1,994	2,030	28 ^a	48	1,953	1,972
American Indian or Alaska Native	191	156	*	*	187	152
Native Hawaiian or Other Pacific Islander	69	*	*	*	67	*
Asian	372	289	*	*	368	286
Two or More Races	253	262	*	*	241	257
Hispanic or Latino	1,740	1,898	35	36	1,690	1,849
TRIMESTER²						
First	N/A	N/A	155	170	N/A	N/A
Second	N/A	N/A	150 ^a	105	N/A	N/A
Third	N/A	N/A	130	126	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.72B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	30.2 ^a	29.0	18.0	16.6	30.7 ^a	29.6
AGE						
15-17	20.9 ^b	18.7	25.9	22.3	20.7 ^b	18.5
18-25	36.7 ^a	35.3	27.8	26.4	37.2 ^a	35.8
26-44	29.1	28.1	10.8	10.4	29.8	28.8
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	32.3 ^b	30.8	21.0	19.0	32.7 ^a	31.3
White	35.2 ^a	33.8	25.0	21.5	35.6 ^a	34.3
Black or African American	24.2	23.8	9.3	15.0	24.7	24.2
American Indian or Alaska Native	49.2	40.6	*	*	52.0	40.6
Native Hawaiian or Other Pacific Islander	36.5	*	*	*	36.7	*
Asian	12.2	9.1	*	*	12.8	9.5
Two or More Races	38.4	39.6	*	*	38.6	40.0
Hispanic or Latino	18.7	19.5	6.8	7.2	19.3	20.2
TRIMESTER²						
First	N/A	N/A	22.0	23.7	N/A	N/A
Second	N/A	N/A	15.5	12.9	N/A	N/A
Third	N/A	N/A	17.3	13.7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.73A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	31,647	31,824	238	300	31,246	31,376
AGE						
15-17	1,717	1,699	14	10	1,694	1,672
18-25	8,722 ^a	8,963	100	88	8,577	8,826
26-44	21,208	21,163	124 ^a	202	20,975	20,879
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	28,113	28,192	194 ^a	266	27,783	27,801
White	23,042	23,095	154	204	22,802	22,790
Black or African American	3,384	3,453	19 ^a	43	3,336	3,388
American Indian or Alaska Native	194	170	*	*	186	169
Native Hawaiian or Other Pacific Islander	78	*	*	*	77	*
Asian	1,063	994	*	*	1,036	983
Two or More Races	352	400	*	*	346	397
Hispanic or Latino	3,534	3,632	44	34	3,464	3,575
TRIMESTER²						
First	N/A	N/A	138	147	N/A	N/A
Second	N/A	N/A	59	84	N/A	N/A
Third	N/A	N/A	35	61	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.73B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	51.3	51.4	9.8	12.1	53.0	53.1
AGE						
15-17	28.5	27.6	14.5	13.9	28.7	27.7
18-25	55.7	55.7	10.5	9.7	58.7	58.5
26-44	53.0	53.4	8.9 ^a	13.5	54.6	55.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	53.7	54.1	10.1 ^a	13.4	55.3	55.7
White	57.8	58.9	10.8	13.8	59.6	60.7
Black or African American	41.0	40.5	6.4 ^a	13.4	42.2	41.6
American Indian or Alaska Native	49.8	44.3	*	*	51.7	45.0
Native Hawaiian or Other Pacific Islander	40.9	*	*	*	42.1	*
Asian	34.9	31.2	*	*	36.0	32.4
Two or More Races	53.3	60.5	*	*	55.6	61.8
Hispanic or Latino	37.9	37.4	8.6	6.8	39.6	39.1
TRIMESTER²						
First	N/A	N/A	19.6	20.6	N/A	N/A
Second	N/A	N/A	6.1	10.2	N/A	N/A
Third	N/A	N/A	4.7	6.7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.74A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	13,849	13,939	99	96	13,676	13,782
AGE						
15-17	999	1,023	7	6	985	1,003
18-25	4,968	5,162	46	45	4,893	5,089
26-44	7,882	7,754	46	45	7,798	7,690
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	12,194	12,178	84	78	12,050	12,051
White	10,065	10,002	57	47	9,975	9,915
Black or African American	1,513	1,485	9 ^a	28	1,481	1,447
American Indian or Alaska Native	134	121	*	*	126	120
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	274	345	*	*	263	344
Two or More Races	156	179	*	*	154	177
Hispanic or Latino	1,655	1,761	15	19	1,626	1,732
TRIMESTER²						
First	N/A	N/A	77	54	N/A	N/A
Second	N/A	N/A	13	21	N/A	N/A
Third	N/A	N/A	5	15	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.74B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	22.4	22.5	4.1	3.9	23.2	23.3
AGE						
15-17	16.6	16.6	7.0	8.4	16.7	16.6
18-25	31.7	32.1	4.8	4.9	33.5	33.8
26-44	19.7	19.6	3.3	3.0	20.3	20.3
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	23.3	23.3	4.4	3.9	24.0	24.1
White	25.3	25.5	4.0	3.2	26.1	26.4
Black or African American	18.3	17.4	3.0 ^a	8.8	18.7	17.8
American Indian or Alaska Native	34.3	31.6	*	*	35.1	32.1
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	9.0	10.8	*	*	9.1	11.3
Two or More Races	23.7	27.1	*	*	24.7	27.6
Hispanic or Latino	17.8	18.1	2.9	3.7	18.6	18.9
TRIMESTER²						
First	N/A	N/A	10.9	7.5	N/A	N/A
Second	N/A	N/A	1.4	2.6	N/A	N/A
Third	N/A	N/A	0.7	1.6	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.75A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	3,166	3,158	16	18	3,140	3,132
AGE						
15-17	218	216	0	2	216	211
18-25	1,380	1,418	7	7	1,365	1,408
26-44	1,568	1,524	9	9	1,559	1,513
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	2,945	2,904	15	13	2,924	2,883
White	2,568	2,552	11	12	2,553	2,533
Black or African American	237	235	1	1	233	234
American Indian or Alaska Native	27	21	*	*	25	21
Native Hawaiian or Other Pacific Islander	12	10	*	*	11	10
Asian	66	40	*	*	66	40
Two or More Races	36	45	*	*	35	44
Hispanic or Latino	221	254	1	*	217	249
TRIMESTER²						
First	N/A	N/A	12	12	N/A	N/A
Second	N/A	N/A	*	6	N/A	N/A
Third	N/A	N/A	1	1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.75B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	5.1	5.1	0.7	0.7	5.3	5.3
AGE						
15-17	3.6	3.5	0.1	2.9	3.7	3.5
18-25	8.8	8.8	0.7	0.8	9.3	9.3
26-44	3.9	3.8	0.6	0.6	4.1	4.0
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	5.6	5.6	0.8	0.7	5.8	5.8
White	6.4	6.5	0.8	0.8	6.7	6.8
Black or African American	2.9	2.8	0.5	0.3	2.9	2.9
American Indian or Alaska Native	6.8	5.6	*	*	7.0	5.7
Native Hawaiian or Other Pacific Islander	6.2	4.7	*	*	6.1	4.7
Asian	2.2	1.3	*	*	2.3	1.3
Two or More Races	5.4	6.8	*	*	5.6	6.9
Hispanic or Latino	2.4	2.6	0.2	*	2.5	2.7
TRIMESTER²						
First	N/A	N/A	1.6	1.7	N/A	N/A
Second	N/A	N/A	*	0.7	N/A	N/A
Third	N/A	N/A	0.1	0.1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

^a Difference between estimate and 2004-2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2004-2005 estimate is statistically significant at the 0.01 level.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.