

Table 7.117C Asthma in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | ASTHMA IN LIFETIME | | | ASTHMA IN PAST YEAR | | |
|---------------|--------------------|--|-----|---------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 539 | 139 | 562 | 423 | 96 | 426 |
| 12-17 | 82 | 25 | 93 | 63 | 20 | 68 |
| 18-25 | 95 | 47 | 97 | 73 | 33 | 69 |
| 26-34 | 181 | 68 | 185 | 137 | 50 | 135 |
| 35-49 | 262 | 79 | 267 | 190 | 55 | 187 |
| 50-64 | 304 | 72 | 322 | 263 | 37 | 272 |
| 65 or Older | 282 | * | 298 | 214 | * | 222 |
| MALE | 319 | 100 | 312 | 245 | 73 | 240 |
| 12-17 | 59 | 18 | 65 | 45 | 14 | 46 |
| 18-25 | 69 | 37 | 65 | 50 | 26 | 44 |
| 26-34 | 137 | 58 | 133 | 99 | 41 | 94 |
| 35-49 | 159 | 49 | 155 | 105 | 32 | 99 |
| 50-64 | 174 | 53 | 172 | 133 | 34 | 136 |
| 65 or Older | 140 | * | 139 | 127 | * | 125 |
| FEMALE | 419 | 104 | 445 | 342 | 63 | 352 |
| 12-17 | 57 | 15 | 63 | 46 | 13 | 48 |
| 18-25 | 65 | 31 | 68 | 52 | 21 | 51 |
| 26-34 | 127 | 37 | 133 | 104 | 30 | 104 |
| 35-49 | 199 | 62 | 206 | 156 | 44 | 157 |
| 50-64 | 257 | * | 269 | 223 | * | 234 |
| 65 or Older | 246 | * | 264 | 179 | * | 188 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.117D Asthma in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | ASTHMA IN LIFETIME | | | ASTHMA IN PAST YEAR | | |
|---------------|--------------------|--|------|---------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.22 | 0.61 | 0.24 | 0.18 | 0.43 | 0.19 |
| 12-17 | 0.34 | 1.15 | 0.35 | 0.26 | 0.96 | 0.27 |
| 18-25 | 0.30 | 0.63 | 0.33 | 0.23 | 0.47 | 0.25 |
| 26-34 | 0.53 | 1.52 | 0.56 | 0.40 | 1.14 | 0.43 |
| 35-49 | 0.41 | 1.31 | 0.43 | 0.29 | 0.92 | 0.31 |
| 50-64 | 0.61 | 2.83 | 0.65 | 0.53 | 1.52 | 0.55 |
| 65 or Older | 0.81 | * | 0.80 | 0.61 | * | 0.62 |
| MALE | 0.28 | 0.70 | 0.30 | 0.21 | 0.51 | 0.23 |
| 12-17 | 0.48 | 1.71 | 0.50 | 0.37 | 1.42 | 0.38 |
| 18-25 | 0.43 | 0.82 | 0.51 | 0.32 | 0.59 | 0.37 |
| 26-34 | 0.81 | 1.94 | 0.89 | 0.59 | 1.37 | 0.65 |
| 35-49 | 0.50 | 1.29 | 0.54 | 0.33 | 0.85 | 0.35 |
| 50-64 | 0.72 | 2.95 | 0.75 | 0.55 | 1.93 | 0.60 |
| 65 or Older | 0.94 | * | 0.94 | 0.85 | * | 0.85 |
| FEMALE | 0.34 | 1.19 | 0.35 | 0.28 | 0.76 | 0.29 |
| 12-17 | 0.48 | 1.44 | 0.50 | 0.38 | 1.27 | 0.40 |
| 18-25 | 0.41 | 1.05 | 0.44 | 0.33 | 0.78 | 0.35 |
| 26-34 | 0.73 | 2.52 | 0.76 | 0.60 | 2.12 | 0.62 |
| 35-49 | 0.61 | 2.74 | 0.61 | 0.47 | 2.03 | 0.48 |
| 50-64 | 1.00 | * | 1.01 | 0.87 | * | 0.89 |
| 65 or Older | 1.23 | * | 1.22 | 0.89 | * | 0.90 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.118C Bronchitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | BRONCHITIS IN LIFETIME | | | BRONCHITIS IN PAST YEAR | | |
|---------------|------------------------|--|-----|-------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 528 | 152 | 572 | 374 | 92 | 379 |
| 12-17 | 59 | 15 | 60 | 39 | 11 | 37 |
| 18-25 | 79 | 48 | 75 | 47 | 30 | 40 |
| 26-34 | 173 | 62 | 171 | 102 | 33 | 96 |
| 35-49 | 263 | 74 | 274 | 148 | 38 | 146 |
| 50-64 | 346 | 94 | 367 | 257 | 57 | 258 |
| 65 or Older | 259 | * | 275 | 190 | * | 191 |
| MALE | 330 | 107 | 330 | 226 | 71 | 220 |
| 12-17 | 41 | 10 | 42 | 26 | 7 | 26 |
| 18-25 | 56 | 33 | 48 | 30 | 21 | 24 |
| 26-34 | 115 | 52 | 109 | 73 | 29 | 68 |
| 35-49 | 166 | 56 | 162 | 88 | 29 | 83 |
| 50-64 | 212 | 66 | 208 | 159 | 53 | 154 |
| 65 or Older | 143 | * | 145 | 115 | * | 114 |
| FEMALE | 412 | 105 | 448 | 289 | 58 | 296 |
| 12-17 | 40 | 11 | 40 | 27 | 8 | 26 |
| 18-25 | 56 | 30 | 55 | 36 | 21 | 32 |
| 26-34 | 124 | 34 | 128 | 68 | 15 | 68 |
| 35-49 | 198 | 48 | 212 | 120 | 24 | 123 |
| 50-64 | 270 | * | 286 | 199 | * | 204 |
| 65 or Older | 218 | * | 235 | 154 | * | 154 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.118D Bronchitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | BRONCHITIS IN LIFETIME | | | BRONCHITIS IN PAST YEAR | | |
|---------------|------------------------|--|------|-------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.22 | 0.64 | 0.24 | 0.16 | 0.41 | 0.17 |
| 12-17 | 0.24 | 0.77 | 0.26 | 0.16 | 0.55 | 0.16 |
| 18-25 | 0.25 | 0.62 | 0.27 | 0.15 | 0.42 | 0.15 |
| 26-34 | 0.51 | 1.40 | 0.54 | 0.30 | 0.76 | 0.31 |
| 35-49 | 0.41 | 1.23 | 0.43 | 0.23 | 0.65 | 0.24 |
| 50-64 | 0.70 | 3.50 | 0.72 | 0.52 | 2.30 | 0.53 |
| 65 or Older | 0.74 | * | 0.75 | 0.54 | * | 0.54 |
| MALE | 0.29 | 0.71 | 0.31 | 0.20 | 0.50 | 0.21 |
| 12-17 | 0.33 | 1.06 | 0.36 | 0.21 | 0.75 | 0.22 |
| 18-25 | 0.35 | 0.74 | 0.39 | 0.19 | 0.49 | 0.21 |
| 26-34 | 0.68 | 1.74 | 0.75 | 0.43 | 1.01 | 0.47 |
| 35-49 | 0.52 | 1.47 | 0.56 | 0.28 | 0.78 | 0.29 |
| 50-64 | 0.88 | 3.56 | 0.89 | 0.66 | 2.97 | 0.67 |
| 65 or Older | 0.96 | * | 0.96 | 0.77 | * | 0.77 |
| FEMALE | 0.33 | 1.23 | 0.35 | 0.23 | 0.70 | 0.24 |
| 12-17 | 0.33 | 1.07 | 0.35 | 0.22 | 0.76 | 0.23 |
| 18-25 | 0.35 | 1.00 | 0.38 | 0.22 | 0.74 | 0.23 |
| 26-34 | 0.71 | 2.37 | 0.76 | 0.39 | 1.06 | 0.41 |
| 35-49 | 0.60 | 2.25 | 0.62 | 0.36 | 1.18 | 0.38 |
| 50-64 | 1.05 | * | 1.05 | 0.78 | * | 0.79 |
| 65 or Older | 1.09 | * | 1.08 | 0.77 | * | 0.75 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.119C Diabetes in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | DIABETES IN LIFETIME | | | DIABETES IN PAST YEAR | | |
|---------------|----------------------|--|-----|-----------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 585 | 82 | 620 | 542 | 75 | 568 |
| 12-17 | 17 | 6 | 16 | 14 | 5 | 13 |
| 18-25 | 28 | 15 | 25 | 25 | 13 | 21 |
| 26-34 | 86 | 36 | 82 | 82 | 33 | 76 |
| 35-49 | 180 | 52 | 179 | 166 | 51 | 163 |
| 50-64 | 384 | 44 | 419 | 358 | 35 | 387 |
| 65 or Older | 346 | * | 380 | 328 | * | 357 |
| MALE | 386 | 61 | 397 | 359 | 60 | 367 |
| 12-17 | 11 | 5 | 10 | 8 | 5 | 6 |
| 18-25 | 20 | 13 | 16 | 18 | 11 | 14 |
| 26-34 | 52 | 19 | 50 | 50 | 16 | 48 |
| 35-49 | 138 | 50 | 133 | 127 | 50 | 120 |
| 50-64 | 263 | 25 | 285 | 247 | 25 | 265 |
| 65 or Older | 207 | * | 222 | 197 | * | 209 |
| FEMALE | 420 | 54 | 445 | 389 | 44 | 409 |
| 12-17 | 13 | 2 | 13 | 11 | 1 | 11 |
| 18-25 | 20 | 7 | 18 | 17 | 6 | 16 |
| 26-34 | 68 | 30 | 64 | 64 | 29 | 58 |
| 35-49 | 117 | 15 | 119 | 109 | 13 | 111 |
| 50-64 | 278 | * | 305 | 259 | * | 281 |
| 65 or Older | 263 | * | 292 | 248 | * | 272 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.119D Diabetes in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | DIABETES IN LIFETIME | | | DIABETES IN PAST YEAR | | |
|---------------|----------------------|--|------|-----------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.24 | 0.36 | 0.26 | 0.23 | 0.33 | 0.24 |
| 12-17 | 0.07 | 0.29 | 0.07 | 0.06 | 0.28 | 0.06 |
| 18-25 | 0.09 | 0.21 | 0.10 | 0.08 | 0.19 | 0.09 |
| 26-34 | 0.25 | 0.83 | 0.27 | 0.24 | 0.77 | 0.25 |
| 35-49 | 0.28 | 0.89 | 0.29 | 0.26 | 0.88 | 0.27 |
| 50-64 | 0.77 | 1.80 | 0.80 | 0.72 | 1.44 | 0.75 |
| 65 or Older | 0.99 | * | 1.00 | 0.94 | * | 0.95 |
| MALE | 0.33 | 0.43 | 0.37 | 0.31 | 0.42 | 0.35 |
| 12-17 | 0.09 | 0.57 | 0.08 | 0.07 | 0.57 | 0.05 |
| 18-25 | 0.13 | 0.30 | 0.14 | 0.11 | 0.27 | 0.12 |
| 26-34 | 0.31 | 0.67 | 0.36 | 0.30 | 0.56 | 0.34 |
| 35-49 | 0.44 | 1.33 | 0.46 | 0.40 | 1.32 | 0.42 |
| 50-64 | 1.10 | 1.48 | 1.17 | 1.03 | 1.48 | 1.10 |
| 65 or Older | 1.39 | * | 1.42 | 1.32 | * | 1.35 |
| FEMALE | 0.34 | 0.66 | 0.36 | 0.31 | 0.55 | 0.33 |
| 12-17 | 0.11 | 0.16 | 0.11 | 0.09 | 0.07 | 0.10 |
| 18-25 | 0.12 | 0.26 | 0.14 | 0.11 | 0.24 | 0.12 |
| 26-34 | 0.39 | 2.13 | 0.39 | 0.37 | 2.05 | 0.36 |
| 35-49 | 0.36 | 0.76 | 0.38 | 0.33 | 0.66 | 0.35 |
| 50-64 | 1.08 | * | 1.10 | 1.01 | * | 1.03 |
| 65 or Older | 1.31 | * | 1.32 | 1.24 | * | 1.24 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.120C Heart Disease in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | HEART DISEASE IN LIFETIME | | | HEART DISEASE IN PAST YEAR | | |
|---------------|---------------------------|--|-----|----------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 515 | 88 | 535 | 444 | 66 | 454 |
| 12-17 | 15 | 3 | 15 | 9 | 3 | 8 |
| 18-25 | 17 | 8 | 15 | 11 | 4 | 11 |
| 26-34 | 49 | 35 | 35 | 43 | 31 | 30 |
| 35-49 | 115 | 33 | 114 | 98 | 23 | 96 |
| 50-64 | 310 | 67 | 321 | 270 | 51 | 275 |
| 65 or Older | 342 | * | 404 | 302 | * | 345 |
| MALE | 374 | 75 | 382 | 307 | 56 | 309 |
| 12-17 | 10 | 2 | 10 | 8 | 2 | 7 |
| 18-25 | 10 | 5 | 9 | 5 | 2 | 5 |
| 26-34 | 24 | 8 | 23 | 18 | 3 | 17 |
| 35-49 | 88 | 25 | 86 | 70 | 18 | 69 |
| 50-64 | 229 | 65 | 233 | 190 | 49 | 189 |
| 65 or Older | 245 | * | 282 | 214 | * | 235 |
| FEMALE | 348 | 44 | 357 | 320 | 36 | 326 |
| 12-17 | 10 | 2 | 10 | 5 | 2 | 4 |
| 18-25 | 14 | 6 | 12 | 10 | 4 | 9 |
| 26-34 | 43 | 34 | 27 | 39 | * | 25 |
| 35-49 | 78 | 21 | 76 | 68 | 15 | 66 |
| 50-64 | 215 | * | 223 | 200 | * | 207 |
| 65 or Older | 245 | * | 274 | 221 | * | 241 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.120D Heart Disease in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | HEART DISEASE IN LIFETIME | | | HEART DISEASE IN PAST YEAR | | |
|---------------|---------------------------|--|------|----------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.22 | 0.40 | 0.23 | 0.19 | 0.30 | 0.20 |
| 12-17 | 0.06 | 0.15 | 0.07 | 0.04 | 0.15 | 0.04 |
| 18-25 | 0.05 | 0.11 | 0.06 | 0.04 | 0.06 | 0.04 |
| 26-34 | 0.14 | 0.82 | 0.12 | 0.13 | 0.73 | 0.10 |
| 35-49 | 0.18 | 0.57 | 0.19 | 0.15 | 0.41 | 0.16 |
| 50-64 | 0.62 | 2.71 | 0.64 | 0.54 | 2.09 | 0.56 |
| 65 or Older | 0.98 | * | 0.99 | 0.86 | * | 0.87 |
| MALE | 0.32 | 0.54 | 0.36 | 0.27 | 0.40 | 0.30 |
| 12-17 | 0.08 | 0.25 | 0.09 | 0.06 | 0.25 | 0.06 |
| 18-25 | 0.06 | 0.11 | 0.07 | 0.03 | 0.04 | 0.04 |
| 26-34 | 0.14 | 0.27 | 0.16 | 0.11 | 0.12 | 0.12 |
| 35-49 | 0.28 | 0.69 | 0.30 | 0.22 | 0.49 | 0.25 |
| 50-64 | 0.95 | 3.54 | 0.99 | 0.79 | 2.78 | 0.82 |
| 65 or Older | 1.64 | * | 1.67 | 1.44 | * | 1.47 |
| FEMALE | 0.28 | 0.55 | 0.30 | 0.26 | 0.45 | 0.27 |
| 12-17 | 0.08 | 0.18 | 0.09 | 0.04 | 0.18 | 0.04 |
| 18-25 | 0.09 | 0.23 | 0.09 | 0.06 | 0.14 | 0.07 |
| 26-34 | 0.25 | 2.41 | 0.17 | 0.23 | * | 0.15 |
| 35-49 | 0.24 | 1.01 | 0.25 | 0.21 | 0.74 | 0.21 |
| 50-64 | 0.84 | * | 0.86 | 0.78 | * | 0.80 |
| 65 or Older | 1.22 | * | 1.23 | 1.10 | * | 1.11 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.121C Hepatitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | HEPATITIS IN LIFETIME | | | HEPATITIS IN PAST YEAR | | |
|---------------|-----------------------|--|-----|------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 218 | 60 | 212 | 127 | 42 | 120 |
| 12-17 | 9 | 1 | 9 | 3 | 0 | 3 |
| 18-25 | 19 | 9 | 17 | 14 | 6 | 12 |
| 26-34 | 61 | 18 | 60 | 45 | 15 | 42 |
| 35-49 | 106 | 52 | 95 | 64 | 34 | 54 |
| 50-64 | 158 | 22 | 159 | 83 | 19 | 81 |
| 65 or Older | 103 | * | 104 | 52 | * | 52 |
| MALE | 162 | 53 | 155 | 93 | 35 | 86 |
| 12-17 | 7 | 1 | 7 | 1 | 0 | 1 |
| 18-25 | 15 | 8 | 13 | 12 | 5 | 11 |
| 26-34 | 41 | 18 | 37 | 18 | 15 | 11 |
| 35-49 | 81 | 44 | 70 | 43 | 25 | 35 |
| 50-64 | 122 | 21 | 122 | 74 | 19 | 72 |
| 65 or Older | 66 | * | 67 | 30 | * | 30 |
| FEMALE | 150 | 29 | 149 | 85 | 23 | 82 |
| 12-17 | 5 | * | 6 | 3 | * | 3 |
| 18-25 | 13 | 5 | 12 | 6 | 3 | 5 |
| 26-34 | 47 | 1 | 47 | 41 | * | 41 |
| 35-49 | 69 | 28 | 63 | 48 | 23 | 42 |
| 50-64 | 93 | * | 94 | 39 | * | 38 |
| 65 or Older | 79 | * | 80 | 42 | * | 42 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.121D Hepatitis in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | HEPATITIS IN LIFETIME | | | HEPATITIS IN PAST YEAR | | |
|---------------|-----------------------|--|------|------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.09 | 0.27 | 0.10 | 0.05 | 0.19 | 0.05 |
| 12-17 | 0.04 | 0.06 | 0.04 | 0.01 | 0.01 | 0.02 |
| 18-25 | 0.06 | 0.13 | 0.07 | 0.04 | 0.09 | 0.05 |
| 26-34 | 0.18 | 0.43 | 0.20 | 0.13 | 0.36 | 0.14 |
| 35-49 | 0.16 | 0.90 | 0.16 | 0.10 | 0.60 | 0.09 |
| 50-64 | 0.32 | 0.92 | 0.33 | 0.17 | 0.77 | 0.17 |
| 65 or Older | 0.29 | * | 0.30 | 0.15 | * | 0.15 |
| MALE | 0.14 | 0.38 | 0.15 | 0.08 | 0.25 | 0.08 |
| 12-17 | 0.06 | 0.12 | 0.06 | 0.01 | 0.02 | 0.01 |
| 18-25 | 0.09 | 0.18 | 0.11 | 0.08 | 0.13 | 0.09 |
| 26-34 | 0.24 | 0.63 | 0.26 | 0.11 | 0.53 | 0.08 |
| 35-49 | 0.26 | 1.18 | 0.25 | 0.13 | 0.68 | 0.12 |
| 50-64 | 0.51 | 1.24 | 0.54 | 0.31 | 1.09 | 0.32 |
| 65 or Older | 0.44 | * | 0.46 | 0.20 | * | 0.21 |
| FEMALE | 0.12 | 0.37 | 0.13 | 0.07 | 0.29 | 0.07 |
| 12-17 | 0.05 | * | 0.05 | 0.03 | * | 0.03 |
| 18-25 | 0.08 | 0.19 | 0.09 | 0.04 | 0.12 | 0.04 |
| 26-34 | 0.27 | 0.07 | 0.29 | 0.23 | * | 0.25 |
| 35-49 | 0.21 | 1.37 | 0.20 | 0.15 | 1.13 | 0.14 |
| 50-64 | 0.36 | * | 0.37 | 0.15 | * | 0.15 |
| 65 or Older | 0.39 | * | 0.40 | 0.21 | * | 0.21 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.122C High Blood Pressure in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | HIGH BLOOD PRESSURE IN LIFETIME | | | HIGH BLOOD PRESSURE IN PAST YEAR | | |
|---------------|---------------------------------|--|-----|----------------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 809 | 179 | 966 | 752 | 166 | 861 |
| 12-17 | 29 | 8 | 28 | 22 | 5 | 22 |
| 18-25 | 51 | 23 | 46 | 42 | 19 | 37 |
| 26-34 | 151 | 48 | 152 | 122 | 32 | 124 |
| 35-49 | 292 | 101 | 313 | 269 | 91 | 275 |
| 50-64 | 496 | 121 | 593 | 462 | 115 | 543 |
| 65 or Older | 453 | * | 682 | 431 | * | 606 |
| MALE | 539 | 151 | 606 | 482 | 138 | 518 |
| 12-17 | 22 | 5 | 21 | 16 | 3 | 16 |
| 18-25 | 38 | 18 | 34 | 31 | 15 | 27 |
| 26-34 | 121 | 42 | 120 | 97 | 28 | 97 |
| 35-49 | 221 | 81 | 226 | 202 | 71 | 198 |
| 50-64 | 333 | 100 | 372 | 310 | 94 | 340 |
| 65 or Older | 291 | * | 391 | 271 | * | 326 |
| FEMALE | 564 | 109 | 677 | 536 | 101 | 631 |
| 12-17 | 19 | 6 | 18 | 15 | 4 | 15 |
| 18-25 | 35 | 15 | 33 | 29 | 12 | 27 |
| 26-34 | 88 | 23 | 87 | 74 | 13 | 74 |
| 35-49 | 194 | 65 | 207 | 177 | 58 | 187 |
| 50-64 | 342 | * | 420 | 327 | * | 394 |
| 65 or Older | 334 | * | 489 | 326 | * | 455 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.122D High Blood Pressure in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | HIGH BLOOD PRESSURE IN LIFETIME | | | HIGH BLOOD PRESSURE IN PAST YEAR | | |
|---------------|---------------------------------|--|------|----------------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.34 | 0.74 | 0.36 | 0.31 | 0.71 | 0.33 |
| 12-17 | 0.12 | 0.39 | 0.12 | 0.09 | 0.26 | 0.10 |
| 18-25 | 0.16 | 0.32 | 0.18 | 0.13 | 0.28 | 0.15 |
| 26-34 | 0.44 | 1.10 | 0.48 | 0.36 | 0.74 | 0.40 |
| 35-49 | 0.45 | 1.63 | 0.47 | 0.42 | 1.50 | 0.43 |
| 50-64 | 1.00 | 4.15 | 1.02 | 0.93 | 3.98 | 0.96 |
| 65 or Older | 1.29 | * | 1.29 | 1.23 | * | 1.22 |
| MALE | 0.47 | 0.99 | 0.51 | 0.42 | 0.92 | 0.45 |
| 12-17 | 0.18 | 0.55 | 0.19 | 0.13 | 0.33 | 0.14 |
| 18-25 | 0.24 | 0.41 | 0.28 | 0.20 | 0.35 | 0.23 |
| 26-34 | 0.72 | 1.42 | 0.82 | 0.58 | 0.97 | 0.67 |
| 35-49 | 0.70 | 2.02 | 0.73 | 0.64 | 1.80 | 0.66 |
| 50-64 | 1.38 | 4.84 | 1.43 | 1.29 | 4.58 | 1.35 |
| 65 or Older | 1.95 | * | 1.98 | 1.82 | * | 1.83 |
| FEMALE | 0.46 | 1.25 | 0.48 | 0.43 | 1.19 | 0.46 |
| 12-17 | 0.15 | 0.56 | 0.16 | 0.13 | 0.39 | 0.13 |
| 18-25 | 0.22 | 0.53 | 0.24 | 0.18 | 0.45 | 0.20 |
| 26-34 | 0.50 | 1.62 | 0.53 | 0.42 | 0.94 | 0.45 |
| 35-49 | 0.59 | 2.86 | 0.60 | 0.54 | 2.61 | 0.55 |
| 50-64 | 1.33 | * | 1.34 | 1.28 | * | 1.30 |
| 65 or Older | 1.67 | * | 1.67 | 1.63 | * | 1.63 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.123C Pneumonia in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | PNEUMONIA IN LIFETIME | | | PNEUMONIA IN PAST YEAR | | |
|---------------|-----------------------|--|-----|------------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 480 | 102 | 505 | 237 | 47 | 238 |
| 12-17 | 37 | 9 | 38 | 20 | 6 | 19 |
| 18-25 | 50 | 27 | 46 | 24 | 13 | 21 |
| 26-34 | 113 | 32 | 114 | 43 | 5 | 43 |
| 35-49 | 177 | 50 | 185 | 90 | 27 | 92 |
| 50-64 | 330 | 64 | 356 | 151 | 19 | 152 |
| 65 or Older | 257 | * | 266 | 155 | * | 152 |
| MALE | 307 | 75 | 316 | 155 | 43 | 154 |
| 12-17 | 23 | 5 | 23 | 12 | 4 | 12 |
| 18-25 | 34 | 20 | 29 | 15 | 9 | 12 |
| 26-34 | 67 | 18 | 66 | 32 | 5 | 32 |
| 35-49 | 117 | 40 | 117 | 52 | 23 | 56 |
| 50-64 | 212 | 43 | 221 | 103 | * | 102 |
| 65 or Older | 167 | * | 174 | 99 | * | 96 |
| FEMALE | 372 | 68 | 386 | 181 | 17 | 183 |
| 12-17 | 29 | 8 | 29 | 15 | 5 | 14 |
| 18-25 | 37 | 17 | 35 | 18 | 9 | 17 |
| 26-34 | 89 | 27 | 89 | 29 | 1 | 29 |
| 35-49 | 134 | 31 | 140 | 71 | 14 | 71 |
| 50-64 | 264 | * | 279 | 121 | * | 122 |
| 65 or Older | 189 | * | 196 | 113 | * | 115 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.123D Pneumonia in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | PNEUMONIA IN LIFETIME | | | PNEUMONIA IN PAST YEAR | | |
|---------------|-----------------------|--|------|------------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.20 | 0.45 | 0.22 | 0.10 | 0.21 | 0.11 |
| 12-17 | 0.15 | 0.46 | 0.16 | 0.08 | 0.31 | 0.08 |
| 18-25 | 0.16 | 0.38 | 0.18 | 0.07 | 0.19 | 0.08 |
| 26-34 | 0.33 | 0.74 | 0.36 | 0.13 | 0.12 | 0.14 |
| 35-49 | 0.27 | 0.86 | 0.29 | 0.14 | 0.46 | 0.15 |
| 50-64 | 0.67 | 2.56 | 0.70 | 0.30 | 0.79 | 0.31 |
| 65 or Older | 0.73 | * | 0.73 | 0.44 | * | 0.44 |
| MALE | 0.27 | 0.53 | 0.29 | 0.13 | 0.31 | 0.15 |
| 12-17 | 0.19 | 0.54 | 0.20 | 0.10 | 0.40 | 0.11 |
| 18-25 | 0.21 | 0.45 | 0.24 | 0.10 | 0.21 | 0.11 |
| 26-34 | 0.40 | 0.61 | 0.46 | 0.19 | 0.17 | 0.23 |
| 35-49 | 0.37 | 1.06 | 0.41 | 0.17 | 0.61 | 0.20 |
| 50-64 | 0.88 | 2.42 | 0.93 | 0.43 | * | 0.45 |
| 65 or Older | 1.12 | * | 1.13 | 0.66 | * | 0.66 |
| FEMALE | 0.30 | 0.83 | 0.32 | 0.15 | 0.22 | 0.15 |
| 12-17 | 0.24 | 0.75 | 0.25 | 0.12 | 0.48 | 0.12 |
| 18-25 | 0.23 | 0.64 | 0.26 | 0.12 | 0.34 | 0.13 |
| 26-34 | 0.51 | 1.89 | 0.54 | 0.17 | 0.11 | 0.18 |
| 35-49 | 0.41 | 1.50 | 0.43 | 0.22 | 0.68 | 0.23 |
| 50-64 | 1.03 | * | 1.04 | 0.47 | * | 0.48 |
| 65 or Older | 0.94 | * | 0.95 | 0.57 | * | 0.57 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.124C Sexually Transmitted Disease (STD) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | STD IN LIFETIME | | | STD IN PAST YEAR | | |
|---------------|-----------------|--|-----|------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 296 | 115 | 288 | 119 | 48 | 110 |
| 12-17 | 13 | 9 | 10 | 12 | 8 | 8 |
| 18-25 | 55 | 32 | 45 | 40 | 25 | 30 |
| 26-34 | 134 | 63 | 121 | 74 | 26 | 70 |
| 35-49 | 173 | 66 | 167 | 69 | 28 | 62 |
| 50-64 | 183 | 57 | 180 | 46 | * | 46 |
| 65 or Older | 69 | * | 68 | 18 | * | 14 |
| MALE | 195 | 78 | 190 | 71 | 34 | 62 |
| 12-17 | 7 | 5 | 5 | 6 | 5 | 3 |
| 18-25 | 24 | 16 | 18 | 18 | 13 | 12 |
| 26-34 | 85 | 52 | 67 | 50 | 24 | 44 |
| 35-49 | 120 | 47 | 114 | 43 | 17 | 40 |
| 50-64 | 118 | 28 | 121 | 4 | * | 4 |
| 65 or Older | 60 | * | 59 | 18 | * | 14 |
| FEMALE | 224 | 80 | 215 | 95 | 34 | 90 |
| 12-17 | 11 | 7 | 9 | 10 | 7 | 8 |
| 18-25 | 48 | 26 | 40 | 35 | 21 | 27 |
| 26-34 | 104 | 36 | 101 | 55 | 12 | 54 |
| 35-49 | 126 | 48 | 121 | 52 | 23 | 48 |
| 50-64 | 138 | * | 131 | 46 | * | 46 |
| 65 or Older | 33 | * | 34 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

NOTE: Sexually Transmitted Disease (STD) includes chlamydia, gonorrhea, herpes, and syphilis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.124D Sexually Transmitted Disease (STD) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | STD IN LIFETIME | | | STD IN PAST YEAR | | |
|---------------|-----------------|--|------|------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.12 | 0.51 | 0.13 | 0.05 | 0.22 | 0.05 |
| 12-17 | 0.06 | 0.45 | 0.05 | 0.05 | 0.42 | 0.04 |
| 18-25 | 0.17 | 0.44 | 0.18 | 0.13 | 0.35 | 0.12 |
| 26-34 | 0.39 | 1.42 | 0.39 | 0.22 | 0.63 | 0.23 |
| 35-49 | 0.27 | 1.11 | 0.27 | 0.11 | 0.50 | 0.10 |
| 50-64 | 0.37 | 2.28 | 0.37 | 0.09 | * | 0.10 |
| 65 or Older | 0.20 | * | 0.20 | 0.05 | * | 0.04 |
| MALE | 0.17 | 0.55 | 0.18 | 0.06 | 0.25 | 0.06 |
| 12-17 | 0.06 | 0.54 | 0.05 | 0.05 | 0.52 | 0.03 |
| 18-25 | 0.15 | 0.37 | 0.15 | 0.11 | 0.31 | 0.10 |
| 26-34 | 0.50 | 1.74 | 0.47 | 0.30 | 0.83 | 0.31 |
| 35-49 | 0.38 | 1.25 | 0.40 | 0.14 | 0.46 | 0.14 |
| 50-64 | 0.49 | 1.59 | 0.52 | 0.02 | * | 0.02 |
| 65 or Older | 0.40 | * | 0.41 | 0.12 | * | 0.10 |
| FEMALE | 0.18 | 0.97 | 0.18 | 0.08 | 0.43 | 0.08 |
| 12-17 | 0.09 | 0.71 | 0.08 | 0.09 | 0.66 | 0.07 |
| 18-25 | 0.30 | 0.91 | 0.30 | 0.22 | 0.74 | 0.20 |
| 26-34 | 0.60 | 2.49 | 0.61 | 0.32 | 0.89 | 0.34 |
| 35-49 | 0.38 | 2.17 | 0.38 | 0.16 | 1.11 | 0.15 |
| 50-64 | 0.54 | * | 0.52 | 0.18 | * | 0.18 |
| 65 or Older | 0.17 | * | 0.17 | * | * | * |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

NOTE: Sexually Transmitted Disease (STD) includes chlamydia, gonorrhea, herpes, and syphilis.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.125C Stroke in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | STROKE IN LIFETIME | | | STROKE IN PAST YEAR | | |
|---------------|--------------------|--|-----|---------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 300 | 45 | 299 | 132 | 31 | 129 |
| 12-17 | 8 | 4 | 7 | 2 | 0 | 2 |
| 18-25 | 6 | 3 | 5 | 2 | 2 | * |
| 26-34 | 20 | 4 | 20 | 7 | * | 7 |
| 35-49 | 46 | 22 | 41 | 19 | 5 | 19 |
| 50-64 | 209 | 38 | 208 | 69 | 31 | 61 |
| 65 or Older | 197 | * | 203 | 107 | * | 110 |
| MALE | 169 | 33 | 167 | 91 | 29 | 87 |
| 12-17 | 5 | 2 | 4 | 2 | 0 | 2 |
| 18-25 | 4 | 3 | 3 | 2 | 2 | * |
| 26-34 | 15 | * | 15 | 3 | * | 3 |
| 35-49 | 32 | 9 | 31 | 13 | * | 13 |
| 50-64 | 104 | 31 | 101 | 50 | 29 | 41 |
| 65 or Older | 126 | * | 128 | 75 | * | 75 |
| FEMALE | 235 | 31 | 237 | 98 | 12 | 98 |
| 12-17 | 6 | 3 | 5 | * | * | 1 |
| 18-25 | 4 | 1 | 4 | * | * | * |
| 26-34 | 14 | 4 | 13 | 6 | * | 6 |
| 35-49 | 34 | 20 | 27 | 14 | 5 | 13 |
| 50-64 | 179 | * | 181 | 47 | * | 46 |
| 65 or Older | 141 | * | 146 | 83 | * | 85 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.125D Stroke in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | STROKE IN LIFETIME | | | STROKE IN PAST YEAR | | |
|---------------|--------------------|--|------|---------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.13 | 0.20 | 0.14 | 0.06 | 0.14 | 0.06 |
| 12-17 | 0.03 | 0.19 | 0.03 | 0.01 | 0.01 | 0.01 |
| 18-25 | 0.02 | 0.05 | 0.02 | 0.01 | 0.03 | * |
| 26-34 | 0.06 | 0.09 | 0.07 | 0.02 | * | 0.02 |
| 35-49 | 0.07 | 0.38 | 0.07 | 0.03 | 0.09 | 0.03 |
| 50-64 | 0.42 | 1.53 | 0.43 | 0.14 | 1.27 | 0.13 |
| 65 or Older | 0.56 | * | 0.57 | 0.31 | * | 0.31 |
| MALE | 0.15 | 0.23 | 0.16 | 0.08 | 0.21 | 0.09 |
| 12-17 | 0.04 | 0.18 | 0.04 | 0.02 | 0.02 | 0.02 |
| 18-25 | 0.02 | 0.07 | 0.02 | 0.01 | 0.06 | * |
| 26-34 | 0.09 | * | 0.11 | 0.02 | * | 0.02 |
| 35-49 | 0.10 | 0.24 | 0.11 | 0.04 | * | 0.05 |
| 50-64 | 0.43 | 1.81 | 0.45 | 0.21 | 1.67 | 0.18 |
| 65 or Older | 0.84 | * | 0.86 | 0.50 | * | 0.51 |
| FEMALE | 0.19 | 0.39 | 0.20 | 0.08 | 0.15 | 0.08 |
| 12-17 | 0.05 | 0.32 | 0.05 | * | * | 0.01 |
| 18-25 | 0.03 | 0.05 | 0.03 | * | * | * |
| 26-34 | 0.08 | 0.28 | 0.08 | 0.03 | * | 0.04 |
| 35-49 | 0.10 | 0.98 | 0.09 | 0.04 | 0.24 | 0.04 |
| 50-64 | 0.70 | * | 0.71 | 0.18 | * | 0.18 |
| 65 or Older | 0.70 | * | 0.71 | 0.41 | * | 0.42 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.126C Ulcer(s) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Numbers in Thousands, 2005

| Gender/Age | ULCER(S) IN LIFETIME | | | ULCER(S) IN PAST YEAR | | |
|---------------|----------------------|--|-----|-----------------------|--|-----|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 356 | 97 | 349 | 193 | 39 | 188 |
| 12-17 | 18 | 7 | 17 | 12 | 6 | 11 |
| 18-25 | 40 | 20 | 35 | 29 | 13 | 26 |
| 26-34 | 105 | 38 | 96 | 69 | 29 | 64 |
| 35-49 | 145 | 54 | 139 | 80 | 17 | 79 |
| 50-64 | 224 | 62 | 216 | 113 | * | 112 |
| 65 or Older | 199 | * | 205 | 105 | * | 105 |
| MALE | 239 | 69 | 229 | 117 | 30 | 114 |
| 12-17 | 13 | 5 | 12 | 9 | 5 | 8 |
| 18-25 | 25 | 14 | 21 | 18 | 8 | 17 |
| 26-34 | 78 | 35 | 65 | 53 | 27 | 46 |
| 35-49 | 96 | 41 | 88 | 47 | 9 | 46 |
| 50-64 | 139 | 35 | 136 | 51 | * | 51 |
| 65 or Older | 131 | * | 134 | 72 | * | 73 |
| FEMALE | 262 | 66 | 257 | 147 | 25 | 146 |
| 12-17 | 13 | 5 | 12 | 9 | 3 | 8 |
| 18-25 | 29 | 14 | 27 | 21 | 11 | 19 |
| 26-34 | 69 | 14 | 69 | 45 | 9 | 45 |
| 35-49 | 106 | 34 | 107 | 65 | 14 | 65 |
| 50-64 | 169 | * | 162 | 100 | * | 99 |
| 65 or Older | 146 | * | 147 | 76 | * | 76 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

Table 7.126D Ulcer(s) in Lifetime and Past Year among Persons Aged 12 or Older, by Past Year Illicit Drug or Alcohol Dependence or Abuse, Gender, and Age Group: Standard Errors of Percentages, 2005

| Gender/Age | ULCER(S) IN LIFETIME | | | ULCER(S) IN PAST YEAR | | |
|---------------|----------------------|--|------|-----------------------|--|------|
| | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | | Total | Illicit Drug or Alcohol Dependence or Abuse ^{1,2} | |
| | | Yes | No | | Yes | No |
| TOTAL | 0.15 | 0.44 | 0.16 | 0.08 | 0.18 | 0.09 |
| 12-17 | 0.08 | 0.36 | 0.08 | 0.05 | 0.30 | 0.05 |
| 18-25 | 0.13 | 0.28 | 0.14 | 0.09 | 0.19 | 0.10 |
| 26-34 | 0.31 | 0.89 | 0.31 | 0.20 | 0.67 | 0.21 |
| 35-49 | 0.22 | 0.91 | 0.23 | 0.12 | 0.29 | 0.13 |
| 50-64 | 0.45 | 2.49 | 0.45 | 0.23 | * | 0.24 |
| 65 or Older | 0.57 | * | 0.58 | 0.30 | * | 0.30 |
| MALE | 0.21 | 0.49 | 0.22 | 0.10 | 0.22 | 0.11 |
| 12-17 | 0.11 | 0.58 | 0.11 | 0.08 | 0.52 | 0.07 |
| 18-25 | 0.16 | 0.32 | 0.18 | 0.12 | 0.18 | 0.14 |
| 26-34 | 0.46 | 1.21 | 0.46 | 0.31 | 0.94 | 0.32 |
| 35-49 | 0.30 | 1.09 | 0.31 | 0.15 | 0.25 | 0.16 |
| 50-64 | 0.58 | 2.01 | 0.60 | 0.21 | * | 0.23 |
| 65 or Older | 0.88 | * | 0.90 | 0.48 | * | 0.50 |
| FEMALE | 0.21 | 0.81 | 0.22 | 0.12 | 0.32 | 0.13 |
| 12-17 | 0.11 | 0.45 | 0.11 | 0.07 | 0.34 | 0.07 |
| 18-25 | 0.18 | 0.51 | 0.20 | 0.13 | 0.39 | 0.14 |
| 26-34 | 0.40 | 1.01 | 0.42 | 0.26 | 0.66 | 0.28 |
| 35-49 | 0.32 | 1.58 | 0.34 | 0.20 | 0.67 | 0.21 |
| 50-64 | 0.66 | * | 0.64 | 0.39 | * | 0.39 |
| 65 or Older | 0.73 | * | 0.73 | 0.38 | * | 0.38 |

*Low precision; no estimate reported.

NOTE: Respondents were asked to indicate whether a doctor or other medical professional had told them that they had any of 20 listed health conditions in their lifetime or in the past year. Respondents with unknown specific health condition information (e.g., on asthma, bronchitis, diabetes) were excluded.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Dependence or abuse is based on definitions found in the 4th edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.