

**Table 7.17C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	199	196	230	218	203	214	149	186
Marijuana and Hashish	178	172	200	184	184	186	106	133
Cocaine	81	91	76	70	35	59	33	55
Crack	33	48	50	45	14	26	21	40
Heroin	18	9	15	12	11	12	21	13
Hallucinogens	35	37	33	38	22	48	21	41
LSD	14	9	9	10	5	7	12	6
PCP	10	8	4	6	*	3	13	7
Ecstasy	24	29	29	20	21	24	11	13
Inhalants	15	27	31	21	21	22	49	39
Nonmedical Use of Psychotherapeutics <sup>3</sup>	96	100	108	115	97	110	102	125
Pain Relievers	79	80	83	98	70	92	89	108
OxyContin <sup>®</sup>	22	22	18	20	15	21	15	10
Tranquilizers	52	62	51	52	50	58	50	62
Stimulants	46	35	59	46	39	34	35	35
Methamphetamine	34	25	37	33	33	28	29	26
Sedatives	14	17	19	15	34	34	25	23
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	124	130	136	138	105	128	120	145

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.17D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	1.04	0.99	0.55	0.53	0.30	0.29	0.12	0.16
Marijuana and Hashish	0.96	0.90	0.49	0.45	0.28	0.26	0.09	0.11
Cocaine	0.46	0.55	0.19	0.18	0.05	0.08	0.03	0.05
Crack	0.19	0.30	0.13	0.12	0.02	0.04	0.02	0.03
Heroin	0.11	0.06	0.04	0.03	0.02	0.02	0.02	0.01
Hallucinogens	0.21	0.23	0.09	0.10	0.03	0.07	0.02	0.04
LSD	0.08	0.05	0.02	0.03	0.01	0.01	0.01	0.00
PCP	0.06	0.05	0.01	0.01	*	0.00	0.01	0.01
Ecstasy	0.15	0.18	0.08	0.05	0.03	0.03	0.01	0.01
Inhalants	0.09	0.17	0.08	0.05	0.03	0.03	0.04	0.03
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.55	0.59	0.28	0.29	0.15	0.15	0.08	0.11
Pain Relievers	0.46	0.48	0.22	0.24	0.11	0.13	0.07	0.09
OxyContin <sup>®</sup>	0.13	0.14	0.05	0.05	0.02	0.03	0.01	0.01
Tranquilizers	0.31	0.38	0.13	0.13	0.08	0.08	0.04	0.05
Stimulants	0.27	0.21	0.15	0.12	0.06	0.05	0.03	0.03
Methamphetamine	0.20	0.16	0.10	0.08	0.05	0.04	0.02	0.02
Sedatives	0.08	0.11	0.05	0.04	0.05	0.05	0.02	0.02
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.69	0.75	0.34	0.34	0.16	0.18	0.10	0.12

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.17P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS <sup>2</sup>	0.8522		0.6349		0.6454		0.0225	
Marijuana and Hashish	0.2858		0.6594		0.8482		0.2058	
Cocaine	0.0653		0.4517		0.0757		0.0472	
Crack	0.0936		0.6380		0.0429		0.0443	
Heroin	0.2114		0.8282		0.3518		0.5164	
Hallucinogens	0.5888		0.5364		0.4784		0.1730	
LSD	0.1030		0.6669		0.6101		0.5365	
PCP	0.8250		0.6272		*		0.8191	
Ecstasy	0.0480		0.2900		0.9516		0.4843	
Inhalants	0.5665		0.1783		0.6715		0.7628	
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.7979		0.9811		0.8597		0.0549	
Pain Relievers	0.8714		0.6822		0.7012		0.2530	
OxyContin®	0.9443		0.7782		0.3363		0.5379	
Tranquilizers	0.0930		0.6329		0.9316		0.2762	
Stimulants	0.1220		0.5675		0.9107		0.8414	
Methamphetamine	0.2610		0.9956		0.7928		0.7018	
Sedatives	0.3366		0.4040		0.7999		0.7660	
ILLCIT DRUGS OTHER THAN MARIJUANA <sup>2</sup>	0.2311		0.8256		0.4795		0.0280	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.18C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	30	24	40	37	27	25	48	47
Marijuana and Hashish	28	23	35	34	24	22	33	33
Cocaine	9	10	11	9	4	5	4	9
Crack	4	4	1	4	2	3	*	4
Heroin	4	2	2	1	*	3	2	3
Hallucinogens	11	11	10	9	5	7	10	11
LSD	4	4	5	3	3	5	3	2
PCP	4	4	2	2	1	3	*	2
Ecstasy	8	7	7	6	3	3	6	5
Inhalants	8	5	11	10	8	9	20	19
Nonmedical Use of Psychotherapeutics <sup>3</sup>	18	15	23	20	14	12	29	28
Pain Relievers	16	14	19	17	13	11	27	26
OxyContin <sup>®</sup>	7	3	5	6	3	2	5	3
Tranquilizers	8	7	8	11	5	5	11	10
Stimulants	10	9	13	12	4	6	9	9
Methamphetamine	5	6	9	7	1	5	6	6
Sedatives	1	1	2	1	3	4	4	5
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	21	18	26	23	16	15	36	34

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.18D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	2.65	2.66	1.42	1.49	1.47	1.31	0.22	0.21
Marijuana and Hashish	2.67	2.64	1.30	1.45	1.36	1.19	0.16	0.16
Cocaine	1.31	1.57	0.49	0.45	0.26	0.29	0.02	0.04
Crack	0.64	0.60	0.04	0.21	0.15	0.19	*	0.02
Heroin	0.59	0.40	0.10	0.04	*	0.15	0.01	0.01
Hallucinogens	1.57	1.72	0.46	0.46	0.33	0.44	0.05	0.05
LSD	0.66	0.72	0.23	0.16	0.18	0.32	0.02	0.01
PCP	0.64	0.62	0.10	0.09	0.07	0.19	*	0.01
Ecstasy	1.16	1.18	0.34	0.32	0.19	0.16	0.03	0.02
Inhalants	1.12	0.77	0.49	0.52	0.51	0.50	0.10	0.09
Nonmedical Use of Psychotherapeutics <sup>3</sup>	2.25	2.14	0.99	0.98	0.84	0.68	0.14	0.13
Pain Relievers	2.13	2.02	0.86	0.85	0.80	0.63	0.13	0.12
OxyContin <sup>®</sup>	0.99	0.50	0.25	0.30	0.21	0.11	0.02	0.01
Tranquilizers	1.11	1.19	0.39	0.56	0.32	0.28	0.05	0.05
Stimulants	1.40	1.36	0.58	0.64	0.24	0.37	0.04	0.04
Methamphetamine	0.73	1.00	0.41	0.38	0.08	0.28	0.03	0.03
Sedatives	0.22	0.16	0.11	0.06	0.15	0.25	0.02	0.02
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	2.46	2.51	1.09	1.11	0.95	0.84	0.17	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.18P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH					
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>	
	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	0.1296		0.6008		0.1760	0.9081
Marijuana and Hashish	0.2594		0.7179		0.2460	0.3666
Cocaine	0.5480		0.7032		0.8360	0.0200
Crack	0.9890		0.1327		0.9246	*
Heroin	0.5088		0.2083		*	0.3437
Hallucinogens	0.7090		0.5390		0.5001	0.3298
LSD	0.7113		0.4938		0.5732	0.1595
PCP	0.8245		0.8201		0.4545	*
Ecstasy	0.8183		0.7781		0.9468	0.7223
Inhalants	0.3211		0.7201		0.4524	0.5021
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.1455		0.8533		0.3681	0.8910
Pain Relievers	0.1179		0.8560		0.3072	0.9746
OxyContin <sup>®</sup>	0.0043		0.9615		0.1407	0.8572
Tranquilizers	0.8848		0.2334		0.6189	0.7767
Stimulants	0.7995		0.8790		0.4986	0.8554
Methamphetamine	0.1523		0.5786		0.2810	0.7459
Sedatives	0.4367		0.5193		0.3542	0.8890
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.0615		0.9336		0.3262	0.5739

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.19C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	97	91	76	84	50	55	48	49
Marijuana and Hashish	90	80	66	79	47	51	37	40
Cocaine	35	36	21	26	7	12	10	11
Crack	10	10	6	7	4	6	5	8
Heroin	6	5	5	6	1	4	4	5
Hallucinogens	24	22	18	20	9	11	9	15
LSD	12	7	7	6	3	5	2	5
PCP	4	3	4	4	*	*	1	*
Ecstasy	18	17	13	15	7	8	6	10
Inhalants	11	12	13	11	2	4	7	10
Nonmedical Use of Psychotherapeutics <sup>3</sup>	45	47	40	40	27	29	29	29
Pain Relievers	36	39	36	35	24	24	26	25
OxyContin <sup>®</sup>	12	11	10	9	10	10	4	4
Tranquilizers	26	28	21	19	14	16	9	12
Stimulants	24	23	18	19	8	14	12	10
Methamphetamine	15	16	12	14	5	10	9	8
Sedatives	7	8	5	5	2	6	4	11
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	58	58	48	50	28	33	31	35

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.19D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	1.21	1.10	0.75	0.78	0.72	0.78	0.35	0.37
Marijuana and Hashish	1.18	1.06	0.69	0.75	0.69	0.74	0.28	0.31
Cocaine	0.64	0.67	0.25	0.30	0.11	0.20	0.08	0.09
Crack	0.20	0.21	0.07	0.08	0.06	0.10	0.04	0.06
Heroin	0.12	0.10	0.06	0.07	0.02	0.07	0.03	0.04
Hallucinogens	0.48	0.43	0.22	0.23	0.14	0.18	0.07	0.12
LSD	0.24	0.14	0.08	0.07	0.05	0.07	0.01	0.04
PCP	0.08	0.05	0.04	0.05	*	*	0.01	*
Ecstasy	0.36	0.33	0.15	0.17	0.12	0.13	0.05	0.08
Inhalants	0.22	0.23	0.15	0.13	0.03	0.06	0.05	0.08
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.77	0.82	0.45	0.45	0.41	0.45	0.22	0.23
Pain Relievers	0.66	0.71	0.41	0.40	0.37	0.38	0.20	0.19
OxyContin <sup>®</sup>	0.24	0.22	0.12	0.11	0.15	0.15	0.03	0.03
Tranquilizers	0.49	0.53	0.25	0.21	0.23	0.26	0.07	0.10
Stimulants	0.46	0.44	0.21	0.22	0.13	0.22	0.09	0.08
Methamphetamine	0.30	0.31	0.14	0.16	0.09	0.16	0.07	0.06
Sedatives	0.14	0.17	0.06	0.06	0.04	0.09	0.03	0.08
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.92	0.92	0.52	0.55	0.43	0.51	0.24	0.27

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 7.19P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS <sup>2</sup>	0.7841		0.5238		0.2968		0.4681	
Marijuana and Hashish	0.2081		0.2153		0.3932		0.4824	
Cocaine	0.1459		0.3357		0.0735		0.9641	
Crack	0.9279		0.5184		0.2475		0.3409	
Heroin	0.7786		0.6710		0.0476		0.4679	
Hallucinogens	0.5406		0.7101		0.8140		0.2053	
LSD	0.0661		0.3879		0.6407		0.2018	
PCP	0.2149		0.8229		*		*	
Ecstasy	0.6525		0.7396		0.6067		0.4935	
Inhalants	0.9538		0.4473		0.3628		0.4386	
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.9226		0.6985		0.1797		0.8628	
Pain Relievers	0.8279		0.8213		0.5519		0.6973	
OxyContin®	0.9093		0.6440		0.4530		0.9427	
Tranquilizers	0.4909		0.3822		0.5419		0.2797	
Stimulants	0.2842		0.7097		0.1592		0.2281	
Methamphetamine	0.6468		0.5813		0.2753		0.4381	
Sedatives	0.4947		0.7972		0.1745		0.3391	
ILLCIT DRUGS OTHER THAN MARIJUANA <sup>2</sup>	0.3280		0.7960		0.0734		0.5116	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.20C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	172	174	210	196	189	202	135	174
Marijuana and Hashish	148	152	184	160	170	175	93	124
Cocaine	72	80	73	64	34	57	31	52
Crack	30	46	50	45	13	25	21	40
Heroin	17	7	14	10	11	11	20	12
Hallucinogens	26	28	26	30	19	46	20	37
LSD	5	*	2	7	*	*	11	*
PCP	8	7	*	3	*	*	13	7
Ecstasy	14	22	26	13	19	22	7	7
Inhalants	7	24	26	16	19	20	45	33
Nonmedical Use of Psychotherapeutics <sup>3</sup>	81	88	96	107	94	103	90	118
Pain Relievers	69	68	71	88	68	85	78	101
OxyContin <sup>®</sup>	16	19	14	16	11	15	14	8
Tranquilizers	43	56	47	46	48	56	48	60
Stimulants	37	24	52	40	38	31	31	33
Methamphetamine	30	19	30	29	32	26	27	24
Sedatives	12	15	19	13	33	33	25	19
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	108	114	123	128	101	120	108	136

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.20D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
<b>ILLCIT DRUGS<sup>2</sup></b>	1.39	1.42	0.70	0.65	0.32	0.31	0.15	0.21
Marijuana and Hashish	1.24	1.27	0.63	0.54	0.29	0.27	0.11	0.15
Cocaine	0.63	0.74	0.26	0.22	0.06	0.09	0.04	0.06
Crack	0.27	0.44	0.18	0.16	0.02	0.04	0.02	0.05
Heroin	0.15	0.07	0.05	0.04	0.02	0.02	0.02	0.01
Hallucinogens	0.23	0.27	0.09	0.11	0.03	0.07	0.02	0.04
LSD	0.05	*	0.01	0.02	*	*	0.01	*
PCP	0.07	0.06	*	0.01	*	*	0.01	0.01
Ecstasy	0.13	0.21	0.09	0.05	0.03	0.03	0.01	0.01
Inhalants	0.06	0.23	0.09	0.06	0.03	0.03	0.05	0.04
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.71	0.81	0.34	0.36	0.16	0.16	0.10	0.14
Pain Relievers	0.60	0.64	0.26	0.30	0.12	0.13	0.09	0.12
OxyContin <sup>®</sup>	0.15	0.18	0.05	0.06	0.02	0.02	0.02	0.01
Tranquilizers	0.38	0.53	0.17	0.16	0.08	0.09	0.06	0.07
Stimulants	0.33	0.23	0.19	0.14	0.06	0.05	0.04	0.04
Methamphetamine	0.27	0.18	0.11	0.10	0.06	0.04	0.03	0.03
Sedatives	0.11	0.14	0.07	0.05	0.06	0.05	0.03	0.02
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	0.92	1.03	0.43	0.43	0.17	0.19	0.12	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.20P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use <sup>1</sup>		Binge Use But Not Heavy Use <sup>1</sup>		Use But Not Binge Use <sup>1</sup>		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS <sup>2</sup>	0.6920		0.5964		0.5220		0.0320	
Marijuana and Hashish	0.4128		0.4725		0.7207		0.1936	
Cocaine	0.2169		0.2942		0.1462		0.0826	
Crack	0.0739		0.4940		0.0655		0.0768	
Heroin	0.2625		0.7867		0.8380		0.3046	
Hallucinogens	0.2415		0.3527		0.5288		0.5424	
LSD	*		0.2997		*		*	
PCP	0.8442		*		*		0.7078	
Ecstasy	0.0304		0.1855		0.9499		0.9610	
Inhalants	0.3182		0.2611		0.8318		0.7717	
Nonmedical Use of Psychotherapeutics <sup>3</sup>	0.7634		0.8027		0.9524		0.0465	
Pain Relievers	0.8748		0.4923		0.6086		0.2055	
OxyContin®	0.3708		0.9565		0.2436		0.5414	
Tranquilizers	0.1488		0.7429		0.9839		0.3338	
Stimulants	0.1390		0.5218		0.5414		0.5222	
Methamphetamine	0.1413		0.9672		0.5028		0.8018	
Sedatives	0.4519		0.4558		0.9963		0.5154	
ILLCIT DRUGS OTHER THAN MARIJUANA <sup>2</sup>	0.3185		0.8693		0.5826		0.0519	

\*Low precision; no estimate reported.

<sup>1</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.