

Table 7.21C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	293	282	111	132	223	244	100	134
Marijuana and Hashish	252	250	85	105	187	199	63	76
Cocaine	103	116	28	48	59	49	17	25
Crack	60	59	20	38	13	34	6	14
Heroin	23	16	21	12	11	10	*	4
Hallucinogens	43	62	17	38	32	34	13	17
LSD	17	12	12	5	7	9	2	3
PCP	11	10	*	7	*	3	13	*
Ecstasy	32	34	10	9	29	21	5	9
Inhalants	26	29	7	12	30	29	49	37
Nonmedical Use of Psychotherapeutics ²	132	138	75	73	119	134	65	102
Pain Relievers	109	114	65	56	89	108	56	93
OxyContin [®]	30	32	12	10	13	16	10	*
Tranquilizers	70	75	41	51	60	69	28	34
Stimulants	58	54	29	27	58	37	19	23
Methamphetamine	49	43	22	21	32	21	18	16
Sedatives	35	26	20	8	21	31	16	21
ILLCIT DRUGS OTHER THAN MARIJUANA¹	175	179	85	92	135	147	83	112

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.21D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	0.63	0.61	0.57	0.65	0.27	0.28	0.10	0.14
Marijuana and Hashish	0.56	0.56	0.44	0.52	0.23	0.23	0.06	0.08
Cocaine	0.24	0.28	0.14	0.24	0.07	0.06	0.02	0.03
Crack	0.14	0.14	0.10	0.19	0.02	0.04	0.01	0.01
Heroin	0.06	0.04	0.11	0.06	0.01	0.01	*	0.00
Hallucinogens	0.11	0.15	0.09	0.19	0.04	0.04	0.01	0.02
LSD	0.04	0.03	0.06	0.02	0.01	0.01	0.00	0.00
PCP	0.03	0.02	*	0.04	*	0.00	0.01	*
Ecstasy	0.08	0.08	0.05	0.05	0.04	0.02	0.00	0.01
Inhalants	0.06	0.07	0.03	0.06	0.04	0.03	0.05	0.04
Nonmedical Use of Psychotherapeutics ²	0.31	0.32	0.38	0.37	0.15	0.16	0.06	0.11
Pain Relievers	0.25	0.27	0.33	0.28	0.11	0.13	0.06	0.09
OxyContin [®]	0.07	0.08	0.06	0.05	0.02	0.02	0.01	*
Tranquilizers	0.17	0.18	0.21	0.26	0.07	0.08	0.03	0.03
Stimulants	0.14	0.13	0.15	0.14	0.07	0.04	0.02	0.02
Methamphetamine	0.12	0.10	0.11	0.10	0.04	0.02	0.02	0.02
Sedatives	0.09	0.06	0.10	0.04	0.03	0.04	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.40	0.41	0.43	0.46	0.16	0.17	0.08	0.12

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.21P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	0.9233		0.4427		0.5175		0.0276	
Marijuana and Hashish	0.8495		0.4195		0.1530		0.4521	
Cocaine	0.1139		0.1100		0.7499		0.2334	
Crack	0.5247		0.1084		0.0726		0.1610	
Heroin	0.7773		0.3775		0.6229		*	
Hallucinogens	0.4148		0.2628		0.7298		0.4634	
LSD	0.0571		0.4959		0.4096		0.9683	
PCP	0.9818		*		*		*	
Ecstasy	0.2212		0.7361		0.4413		0.5201	
Inhalants	0.6368		0.1238		0.4520		0.9759	
Nonmedical Use of Psychotherapeutics ²	0.5102		0.9417		0.5352		0.0124	
Pain Relievers	0.5769		0.6274		0.4221		0.0776	
OxyContin [®]	0.7808		0.8176		0.1760		*	
Tranquilizers	0.3836		0.8389		0.9736		0.1209	
Stimulants	0.2870		0.8968		0.3298		0.6850	
Methamphetamine	0.6001		0.8151		0.4494		0.6958	
Sedatives	0.8745		0.4632		0.6032		0.8449	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.7405		0.6580		0.6206		0.0150	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.22C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	47	41	29	26	33	29	37	39
Marijuana and Hashish	43	39	25	23	28	25	23	23
Cocaine	14	12	4	6	5	6	*	6
Crack	4	5	*	4	2	3	*	*
Heroin	4	4	1	1	3	1	1	3
Hallucinogens	15	13	7	8	7	8	6	7
LSD	7	6	2	1	2	5	2	2
PCP	5	5	1	1	1	1	*	1
Ecstasy	11	9	5	3	4	4	1	4
Inhalants	14	11	6	10	9	9	20	17
Nonmedical Use of Psychotherapeutics ²	27	23	14	12	17	14	24	26
Pain Relievers	25	20	12	11	16	12	23	23
OxyContin [®]	9	7	5	3	2	1	2	1
Tranquilizers	11	13	8	5	6	4	7	8
Stimulants	14	13	5	4	8	8	8	8
Methamphetamine	10	9	2	2	1	5	6	6
Sedatives	3	4	1	3	2	2	4	4
ILLCIT DRUGS OTHER THAN MARIJUANA¹	33	26	17	16	20	17	31	31

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.22D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	1.51	1.58	2.07	1.86	1.17	1.01	0.19	0.19
Marijuana and Hashish	1.54	1.64	1.79	1.74	1.01	0.90	0.12	0.12
Cocaine	0.76	0.75	0.33	0.53	0.18	0.23	*	0.03
Crack	0.25	0.33	*	0.32	0.09	0.13	*	*
Heroin	0.20	0.23	0.05	0.11	0.10	0.03	0.01	0.01
Hallucinogens	0.77	0.84	0.62	0.68	0.26	0.31	0.03	0.04
LSD	0.37	0.34	0.20	0.11	0.09	0.20	0.01	0.01
PCP	0.26	0.32	0.06	0.13	0.05	0.04	*	0.01
Ecstasy	0.56	0.55	0.45	0.31	0.17	0.16	0.01	0.02
Inhalants	0.70	0.66	0.51	0.83	0.33	0.34	0.10	0.08
Nonmedical Use of Psychotherapeutics ²	1.28	1.24	1.14	1.00	0.63	0.52	0.12	0.13
Pain Relievers	1.21	1.14	0.97	0.94	0.58	0.47	0.12	0.12
OxyContin [®]	0.49	0.41	0.39	0.23	0.09	0.02	0.01	0.01
Tranquilizers	0.58	0.78	0.66	0.43	0.24	0.16	0.04	0.04
Stimulants	0.73	0.80	0.40	0.36	0.31	0.32	0.04	0.04
Methamphetamine	0.55	0.57	0.13	0.19	0.04	0.21	0.03	0.03
Sedatives	0.18	0.24	0.08	0.25	0.07	0.08	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.43	1.43	1.30	1.35	0.74	0.64	0.16	0.15

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.22P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	0.6245		0.8028		0.4492		0.8842	
Marijuana and Hashish	0.6405		0.4952		0.1179		0.9331	
Cocaine	0.9900		0.1514		0.4634		*	
Crack	0.3653		*		0.9595		*	
Heroin	0.7871		0.6283		0.1513		0.4033	
Hallucinogens	0.4856		0.4057		0.4564		0.4453	
LSD	0.4946		0.4381		0.1389		0.2483	
PCP	0.5624		0.2085		0.7667		*	
Ecstasy	0.9079		0.9445		0.7814		0.3313	
Inhalants	0.2707		0.1665		0.5326		0.8547	
Nonmedical Use of Psychotherapeutics ²	0.5621		0.8505		0.3823		0.9123	
Pain Relievers	0.5585		0.4052		0.2710		0.8027	
OxyContin [®]	0.0368		0.8772		0.3127		0.9475	
Tranquilizers	0.0650		0.4655		0.0205		0.7097	
Stimulants	0.7578		0.8099		0.9786		0.7943	
Methamphetamine	0.8852		0.4424		0.1040		0.9083	
Sedatives	0.8858		0.1058		0.6357		0.3325	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2033		0.1966		0.6747		0.8721	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.23C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS¹	122	121	34	35	63	72	33	34
Marijuana and Hashish	112	112	31	31	54	65	23	24
Cocaine	39	43	9	10	16	18	3	5
Crack	12	13	5	6	4	6	*	4
Heroin	8	9	4	4	2	1	1	2
Hallucinogens	29	28	6	8	14	17	7	12
LSD	13	8	2	5	7	7	*	2
PCP	5	5	*	*	*	*	1	*
Ecstasy	21	21	4	5	9	12	5	8
Inhalants	14	16	3	7	10	6	6	8
Nonmedical Use of Psychotherapeutics ²	59	62	16	19	32	34	23	22
Pain Relievers	50	51	14	16	27	29	22	19
OxyContin [®]	19	16	4	4	4	8	1	*
Tranquilizers	32	35	7	10	18	18	6	7
Stimulants	28	31	10	8	12	14	5	6
Methamphetamine	19	24	8	8	7	8	3	*
Sedatives	7	11	2	6	5	4	3	9
ILLICIT DRUGS OTHER THAN MARIJUANA¹	72	77	18	22	40	39	24	26

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.23D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	0.86	0.81	1.15	1.18	0.58	0.65	0.32	0.33
Marijuana and Hashish	0.83	0.81	1.07	1.10	0.52	0.60	0.22	0.24
Cocaine	0.37	0.41	0.34	0.36	0.16	0.18	0.03	0.05
Crack	0.12	0.13	0.19	0.24	0.04	0.06	*	0.04
Heroin	0.08	0.09	0.13	0.17	0.02	0.01	0.01	0.02
Hallucinogens	0.29	0.27	0.22	0.29	0.15	0.17	0.07	0.12
LSD	0.13	0.08	0.06	0.17	0.07	0.07	*	0.02
PCP	0.05	0.05	*	*	*	*	0.01	*
Ecstasy	0.21	0.21	0.15	0.17	0.09	0.12	0.05	0.08
Inhalants	0.14	0.16	0.10	0.26	0.10	0.06	0.06	0.08
Nonmedical Use of Psychotherapeutics ²	0.51	0.55	0.57	0.67	0.33	0.34	0.22	0.22
Pain Relievers	0.45	0.47	0.48	0.58	0.28	0.29	0.22	0.19
OxyContin [®]	0.18	0.16	0.13	0.15	0.04	0.09	0.01	*
Tranquilizers	0.30	0.34	0.25	0.38	0.19	0.18	0.06	0.07
Stimulants	0.27	0.30	0.36	0.29	0.12	0.15	0.05	0.06
Methamphetamine	0.19	0.23	0.29	0.28	0.08	0.08	0.03	*
Sedatives	0.07	0.10	0.08	0.24	0.06	0.04	0.03	0.09
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.60	0.63	0.64	0.78	0.39	0.38	0.24	0.25

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.23P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH						
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes
	2004	2005	2004	2005	2004	2005	2004 2005
ILLCIT DRUGS¹	0.3028		0.3361		0.3062		0.7712
Marijuana and Hashish	0.7985		0.4140		0.2700		0.7255
Cocaine	0.0630		0.8947		0.1555		0.5793
Crack	0.6243		0.5808		0.4180		*
Heroin	0.2778		0.5565		0.4809		0.6000
Hallucinogens	0.7171		0.1407		0.5119		0.5841
LSD	0.0535		0.3098		0.9442		*
PCP	0.5588		*		*		*
Ecstasy	0.6510		0.3677		0.3753		0.7904
Inhalants	0.7767		0.3239		0.8634		0.8260
Nonmedical Use of Psychotherapeutics ²	0.9243		0.7166		0.2944		0.9882
Pain Relievers	0.9497		0.9425		0.4315		0.6270
OxyContin [®]	0.6673		0.8285		0.1070		*
Tranquilizers	0.6516		0.1897		0.9719		0.8952
Stimulants	0.7188		0.1733		0.3111		0.9389
Methamphetamine	0.7222		0.7377		0.6472		*
Sedatives	0.2090		0.2596		0.7918		0.7255
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.1582		0.4133		0.2619		0.7593

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.24C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLICIT DRUGS¹	256	248	102	124	212	229	85	123
Marijuana and Hashish	215	217	76	98	177	186	52	69
Cocaine	94	106	26	46	57	47	17	23
Crack	58	57	20	37	12	33	6	13
Heroin	22	13	20	12	11	10	*	*
Hallucinogens	31	54	14	36	27	28	14	9
LSD	6	6	11	*	*	*	*	*
PCP	8	7	*	7	*	*	13	*
Ecstasy	22	26	7	7	27	17	*	*
Inhalants	19	22	*	*	28	27	45	33
Nonmedical Use of Psychotherapeutics ²	115	122	71	69	114	126	53	96
Pain Relievers	93	101	61	52	83	101	46	86
OxyContin [®]	21	26	10	8	12	13	9	*
Tranquilizers	58	65	40	50	57	66	27	32
Stimulants	48	41	26	25	56	34	17	21
Methamphetamine	43	34	21	19	31	19	17	15
Sedatives	34	23	20	4	20	32	15	19
ILLICIT DRUGS OTHER THAN MARIJUANA¹	154	160	79	87	126	138	72	104

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.24D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS¹	0.81	0.76	0.65	0.76	0.30	0.31	0.12	0.18
Marijuana and Hashish	0.69	0.69	0.48	0.61	0.25	0.25	0.07	0.10
Cocaine	0.32	0.35	0.17	0.29	0.08	0.06	0.02	0.03
Crack	0.20	0.19	0.13	0.23	0.02	0.05	0.01	0.02
Heroin	0.08	0.05	0.13	0.07	0.02	0.01	*	*
Hallucinogens	0.11	0.18	0.09	0.23	0.04	0.04	0.02	0.01
LSD	0.02	0.02	0.07	*	*	*	*	*
PCP	0.03	0.02	*	0.04	*	*	0.02	*
Ecstasy	0.08	0.09	0.04	0.04	0.04	0.02	*	*
Inhalants	0.06	0.08	*	*	0.04	0.04	0.06	0.05
Nonmedical Use of Psychotherapeutics ²	0.39	0.40	0.45	0.43	0.16	0.17	0.07	0.14
Pain Relievers	0.31	0.33	0.39	0.33	0.12	0.14	0.06	0.13
OxyContin [®]	0.07	0.09	0.07	0.05	0.02	0.02	0.01	*
Tranquilizers	0.20	0.22	0.26	0.32	0.08	0.09	0.04	0.05
Stimulants	0.17	0.14	0.17	0.16	0.08	0.05	0.02	0.03
Methamphetamine	0.15	0.12	0.13	0.12	0.05	0.03	0.02	0.02
Sedatives	0.12	0.08	0.13	0.02	0.03	0.04	0.02	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.51	0.52	0.50	0.54	0.18	0.19	0.10	0.16

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.24P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2004	2005	2004	2005	2004	2005	2004	2005
ILLCIT DRUGS ¹	0.9485		0.5086		0.4944		0.0342	
Marijuana and Hashish	0.9951		0.3457		0.1451		0.5398	
Cocaine	0.2940		0.1288		0.4060		0.4850	
Crack	0.6472		0.1564		0.0930		0.2513	
Heroin	0.4093		0.2788		0.9150		*	
Hallucinogens	0.0839		0.5185		0.9582		0.9774	
LSD	0.8883		*		*		*	
PCP	0.9040		*		*		*	
Ecstasy	0.1145		0.9891		0.2078		*	
Inhalants	0.5528		*		0.3837		0.7798	
Nonmedical Use of Psychotherapeutics ²	0.6756		0.8849		0.5714		0.0078	
Pain Relievers	0.7878		0.5217		0.3784		0.0408	
OxyContin®	0.4271		0.7909		0.3826		*	
Tranquilizers	0.5825		0.9428		0.8349		0.1341	
Stimulants	0.3297		0.7467		0.2052		0.5989	
Methamphetamine	0.4447		0.8623		0.2461		0.7882	
Sedatives	0.5782		0.1650		0.5906		0.7641	
ILLCIT DRUGS OTHER THAN MARIJUANA ¹	0.8005		0.9171		0.7037		0.0175	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.25C Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Standard Errors of Numbers in Thousands, 2004 and 2005

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	415	432	67	64	130	130	366	395
Heavy Alcohol Use ¹	199	196	30	24	97	91	172	174
Binge Alcohol Use But Not Heavy Use ¹	230	218	40	37	76	84	210	196
Alcohol Use But Not Binge Use ¹	203	214	27	25	50	55	189	202
No Alcohol Use	149	186	48	47	48	49	135	174
PAST MONTH CIGARETTE USE	324	310	56	49	128	127	278	279
Heavy Alcohol Use ¹	162	149	26	22	81	76	141	128
Binge Alcohol Use But Not Heavy Use ¹	172	169	33	30	63	70	149	149
Alcohol Use But Not Binge Use ¹	129	151	18	17	40	41	119	142
No Alcohol Use	111	132	29	26	34	35	102	124
NO PAST MONTH CIGARETTE USE	248	277	52	49	71	78	228	262
Heavy Alcohol Use ¹	115	116	15	10	32	40	108	110
Binge Alcohol Use But Not Heavy Use ¹	149	129	22	21	41	42	143	119
Alcohol Use But Not Binge Use ¹	147	143	20	18	31	34	138	136
No Alcohol Use	100	134	37	39	33	34	85	123

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.25D Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Standard Errors of Percentages, 2004 and 2005

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	0.17	0.18	0.27	0.25	0.40	0.40	0.20	0.21
Heavy Alcohol Use ¹	1.04	0.99	2.65	2.66	1.21	1.10	1.39	1.42
Binge Alcohol Use But Not Heavy Use ¹	0.55	0.53	1.42	1.49	0.75	0.78	0.70	0.65
Alcohol Use But Not Binge Use ¹	0.30	0.29	1.47	1.31	0.72	0.78	0.32	0.31
No Alcohol Use	0.12	0.16	0.22	0.21	0.35	0.37	0.15	0.21
PAST MONTH CIGARETTE USE	0.48	0.47	1.25	1.25	0.74	0.71	0.58	0.57
Heavy Alcohol Use ¹	1.33	1.30	3.08	3.02	1.39	1.36	1.96	1.90
Binge Alcohol Use But Not Heavy Use ¹	0.95	0.91	2.17	2.22	1.12	1.13	1.25	1.20
Alcohol Use But Not Binge Use ¹	0.87	0.99	3.31	3.63	1.65	1.69	0.98	1.11
No Alcohol Use	0.57	0.65	2.07	1.86	1.15	1.18	0.65	0.76
NO PAST MONTH CIGARETTE USE	0.14	0.15	0.22	0.21	0.33	0.36	0.16	0.18
Heavy Alcohol Use ¹	1.61	1.61	5.10	4.50	1.89	1.99	2.00	2.13
Binge Alcohol Use But Not Heavy Use ¹	0.65	0.57	1.66	1.77	0.95	0.95	0.81	0.68
Alcohol Use But Not Binge Use ¹	0.28	0.25	1.46	1.24	0.68	0.76	0.29	0.26
No Alcohol Use	0.10	0.14	0.19	0.19	0.32	0.33	0.12	0.18

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.25P Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2004	2005	2004	2005	2004	2005	2004	2005
TOTAL	0.4711		0.0563		0.1803		0.4816	
Heavy Alcohol Use ¹	0.8522		0.1296		0.7841		0.6920	
Binge Alcohol Use But Not Heavy Use ¹	0.6349		0.6008		0.5238		0.5964	
Alcohol Use But Not Binge Use ¹	0.6454		0.1760		0.2968		0.5220	
No Alcohol Use	0.0225		0.9081		0.4681		0.0320	
PAST MONTH CIGARETTE USE	0.7927		0.6622		0.1715		0.8046	
Heavy Alcohol Use ¹	0.6236		0.8401		0.7820		0.4779	
Binge Alcohol Use But Not Heavy Use ¹	0.9196		0.9201		0.2866		0.9832	
Alcohol Use But Not Binge Use ¹	0.4070		0.3755		0.4629		0.3330	
No Alcohol Use	0.4427		0.8028		0.3361		0.5086	
NO PAST MONTH CIGARETTE USE	0.3648		0.5720		0.2384		0.4767	
Heavy Alcohol Use ¹	0.6523		0.0095		0.3766		0.8497	
Binge Alcohol Use But Not Heavy Use ¹	0.3655		0.3535		0.7406		0.3889	
Alcohol Use But Not Binge Use ¹	0.8655		0.5163		0.3499		0.8528	
No Alcohol Use	0.0276		0.8842		0.7712		0.0342	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.