

**Table 7.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	356	352
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	133	141	225	227
Cigars	253	237	239	262
Pipe Tobacco	91	94	116	110
<b>ALCOHOL</b>	692	687	1,272	1,251
Binge Alcohol Use <sup>2</sup>	520	504	578	591
Heavy Alcohol Use <sup>2</sup>	284	287	272	253

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.18	0.19
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.22	0.23	0.12	0.12
Cigars	0.38	0.38	0.13	0.14
Pipe Tobacco	0.15	0.15	0.06	0.06
<b>ALCOHOL</b>	0.66	0.63	0.47	0.47
Binge Alcohol Use <sup>2</sup>	0.64	0.63	0.29	0.29
Heavy Alcohol Use <sup>2</sup>	0.43	0.44	0.15	0.13

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.2947	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.6098		0.2650	
Cigars	0.4513		0.9842	
Pipe Tobacco	0.3110		0.2070	
<b>ALCOHOL</b>	0.8966		0.0026	
Binge Alcohol Use <sup>2</sup>	0.6242		0.9715	
Heavy Alcohol Use <sup>2</sup>	0.1045		0.5236	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.27C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	33	32
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	24	21	21	21
Cigars	40	38	25	24
Pipe Tobacco	17	15	10	9
<b>ALCOHOL</b>	63	55	74	73
Binge Alcohol Use <sup>2</sup>	56	48	54	50
Heavy Alcohol Use <sup>2</sup>	32	27	19	19

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.27D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.15	0.14
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.74	0.73	0.10	0.09
Cigars	1.08	1.16	0.11	0.11
Pipe Tobacco	0.53	0.54	0.04	0.04
<b>ALCOHOL</b>	1.22	1.24	0.30	0.29
Binge Alcohol Use <sup>2</sup>	1.26	1.22	0.23	0.22
Heavy Alcohol Use <sup>2</sup>	0.95	0.88	0.08	0.08

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.27P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.2085	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.6083		0.7184	
Cigars	0.6099		0.1564	
Pipe Tobacco	0.6327		0.9404	
<b>ALCOHOL</b>	0.3974		0.3351	
Binge Alcohol Use <sup>2</sup>	0.3637		0.0596	
Heavy Alcohol Use <sup>2</sup>	0.4454		0.7967	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.28C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	67	67
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	52	51	34	37
Cigars	91	84	57	53
Pipe Tobacco	28	31	12	21
<b>ALCOHOL</b>	197	199	188	207
Binge Alcohol Use <sup>2</sup>	174	180	138	155
Heavy Alcohol Use <sup>2</sup>	109	107	63	72

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.28D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.32	0.31
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.38	0.38	0.17	0.18
Cigars	0.55	0.57	0.28	0.25
Pipe Tobacco	0.21	0.24	0.06	0.10
<b>ALCOHOL</b>	0.60	0.58	0.63	0.68
Binge Alcohol Use <sup>2</sup>	0.70	0.69	0.56	0.61
Heavy Alcohol Use <sup>2</sup>	0.64	0.64	0.30	0.34

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.



**Table 7.28P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.5080	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.6425		0.0374	
Cigars	0.3431		0.1001	
Pipe Tobacco	0.7285		0.0105	
<b>ALCOHOL</b>	0.5796		0.4896	
Binge Alcohol Use <sup>2</sup>	0.5016		0.2567	
Heavy Alcohol Use <sup>2</sup>	0.6572		0.0700	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.29C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	354	348
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	120	128	223	224
Cigars	230	214	236	261
Pipe Tobacco	86	89	114	108
<b>ALCOHOL</b>	631	636	1,244	1,218
Binge Alcohol Use <sup>2</sup>	466	456	555	556
Heavy Alcohol Use <sup>2</sup>	259	255	264	242

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.29D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2004 and 2005**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A	N/A	0.24	0.24
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.27	0.28	0.15	0.16
Cigars	0.48	0.46	0.16	0.19
Pipe Tobacco	0.19	0.19	0.08	0.08
<b>ALCOHOL</b>	0.88	0.81	0.60	0.59
Binge Alcohol Use <sup>2</sup>	0.82	0.80	0.37	0.36
Heavy Alcohol Use <sup>2</sup>	0.54	0.53	0.19	0.17

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

**Table 7.29P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004**

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2004	2005	2004	2005
<b>TOBACCO PRODUCTS<sup>1</sup></b>	N/A		0.2911	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3144		0.4021	
Cigars	0.8731		0.6136	
Pipe Tobacco	0.2564		0.3721	
<b>ALCOHOL</b>	0.9001		0.0019	
Binge Alcohol Use <sup>2</sup>	0.6525		0.9723	
Heavy Alcohol Use <sup>2</sup>	0.1498		0.2546	

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.