

Table 7.30C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	300	320	426	430	454	488	548	508
Cigarettes	284	287	401	402	407	440	497	481
Smokeless Tobacco	98	116	122	122	121	122	177	172
Cigars	137	148	173	194	218	211	139	142
Pipe Tobacco	35	43	77	76	62	63	102	103

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.30D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	1.15	1.07	0.81	0.81	0.61	0.62	0.42	0.40
Cigarettes	1.18	1.14	0.82	0.79	0.57	0.57	0.40	0.38
Smokeless Tobacco	0.57	0.68	0.31	0.31	0.18	0.17	0.14	0.15
Cigars	0.77	0.86	0.43	0.48	0.32	0.29	0.12	0.12
Pipe Tobacco	0.21	0.27	0.20	0.19	0.09	0.09	0.09	0.09

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.30P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	0.2518		0.8797		0.3265		0.3127	
Cigarettes	0.7166		0.8673		0.3719		0.4158	
Smokeless Tobacco	0.0889		0.1961		0.5515		0.5913	
Cigars	0.6450		0.4166		0.1309		0.1628	
Pipe Tobacco	0.3460		0.2963		0.9844		0.2377	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.31C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	35	28	48	44	29	28	53	52
Cigarettes	32	27	44	40	25	25	49	46
Smokeless Tobacco	14	12	20	17	8	6	20	20
Cigars	21	20	27	26	17	14	26	27
Pipe Tobacco	8	10	9	10	6	7	13	8

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.31D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	1.82	2.32	1.57	1.61	1.49	1.42	0.24	0.24
Cigarettes	2.43	2.50	1.54	1.62	1.38	1.34	0.23	0.22
Smokeless Tobacco	1.95	1.92	0.89	0.86	0.47	0.35	0.09	0.09
Cigars	2.49	2.58	1.16	1.24	0.96	0.79	0.12	0.13
Pipe Tobacco	1.23	1.53	0.43	0.51	0.34	0.39	0.06	0.04

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.31P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	0.0261		0.9028		0.0979		0.2260	
Cigarettes	0.2132		0.9623		0.2328		0.1918	
Smokeless Tobacco	0.7481		0.9479		0.2163		0.9334	
Cigars	0.7801		0.3713		0.0388		0.6887	
Pipe Tobacco	0.8194		0.9573		0.5521		0.1729	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.32C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	119	117	120	132	76	75	96	85
Cigarettes	109	107	110	121	72	70	90	81
Smokeless Tobacco	41	43	33	35	19	17	22	24
Cigars	62	55	64	64	39	35	41	40
Pipe Tobacco	20	20	15	22	11	14	12	16

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.32D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	0.82	0.90	0.83	0.88	0.93	0.92	0.65	0.62
Cigarettes	0.91	1.03	0.85	0.91	0.94	0.89	0.63	0.60
Smokeless Tobacco	0.76	0.74	0.37	0.39	0.30	0.27	0.17	0.19
Cigars	0.97	0.91	0.66	0.65	0.59	0.54	0.31	0.31
Pipe Tobacco	0.40	0.39	0.18	0.25	0.18	0.23	0.09	0.13

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.32P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	0.0606		0.2930		0.5205		0.2603	
Cigarettes	0.0243		0.9840		0.7105		0.5318	
Smokeless Tobacco	0.8920		0.1895		0.3090		0.9358	
Cigars	0.0036		0.9796		0.2194		0.6761	
Pipe Tobacco	0.3253		0.0194		0.6246		0.2154	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.33C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	274	288	400	406	444	479	540	485
Cigarettes	259	255	373	378	395	428	491	459
Smokeless Tobacco	87	103	118	117	118	120	175	170
Cigars	121	135	164	179	217	208	129	133
Pipe Tobacco	28	36	76	72	60	62	101	102

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.33D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2004 and 2005

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	1.64	1.57	1.08	1.06	0.68	0.68	0.58	0.54
Cigarettes	1.69	1.64	1.09	1.02	0.63	0.63	0.54	0.52
Smokeless Tobacco	0.75	0.95	0.42	0.41	0.20	0.19	0.20	0.20
Cigars	1.02	1.21	0.56	0.61	0.36	0.32	0.15	0.16
Pipe Tobacco	0.25	0.34	0.28	0.25	0.10	0.10	0.12	0.12

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.

Table 7.33P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2005 Versus 2004

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2004	2005	2004	2005	2004	2005	2004	2005
TOBACCO PRODUCTS²	0.0917		0.9485		0.4770		0.1263	
Cigarettes	0.8701		0.8309		0.4962		0.2172	
Smokeless Tobacco	0.0789		0.2928		0.7044		0.5616	
Cigars	0.1579		0.5349		0.2762		0.0964	
Pipe Tobacco	0.1171		0.5965		0.9074		0.2320	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004 and 2005.