

**Table 7.53C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
Standard Errors of Numbers in Thousands, 2005**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	121	24	37	113
From More Than One Doctor	32	11	11	28
Wrote Fake Prescription	10	2	7	6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	18	3	6	17
From Friend or Relative for Free	229	35	78	212
Bought from Friend or Relative	71	18	32	60
Took from Friend or Relative without Asking	42	15	19	34
Bought from Drug Dealer or Other Stranger	56	11	21	49
Bought on the Internet	32	1	6	32
Some Other Way	69	14	19	65

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.53D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2005**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	0.98	1.42	0.89	1.78
From More Than One Doctor	0.29	0.73	0.29	0.48
Wrote Fake Prescription	0.09	0.11	0.19	0.11
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.21	0.15	0.30
From Friend or Relative for Free	1.23	1.70	1.27	2.16
Bought from Friend or Relative	0.63	1.14	0.78	1.03
Took from Friend or Relative without Asking	0.38	0.99	0.49	0.60
Bought from Drug Dealer or Other Stranger	0.49	0.75	0.52	0.83
Bought on the Internet	0.29	0.07	0.16	0.55
Some Other Way	0.62	0.95	0.48	1.11

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.54C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
Standard Errors of Numbers in Thousands, 2005**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	65	8	16	63
From More Than One Doctor	27	2	6	26
Wrote Fake Prescription	1	0	1	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	6	4	3	3
From Friend or Relative for Free	167	18	50	155
Bought from Friend or Relative	49	12	27	37
Took from Friend or Relative without Asking	30	7	14	25
Bought from Drug Dealer or Other Stranger	39	11	20	33
Bought on the Internet	23	*	5	22
Some Other Way	22	6	8	19

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.54D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2005**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	1.20	1.75	0.91	1.99
From More Than One Doctor	0.54	0.55	0.35	0.91
Wrote Fake Prescription	0.02	0.06	0.07	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.12	0.85	0.21	0.10
From Friend or Relative for Free	1.75	3.11	1.86	2.73
Bought from Friend or Relative	0.96	2.52	1.45	1.29
Took from Friend or Relative without Asking	0.60	1.63	0.82	0.86
Bought from Drug Dealer or Other Stranger	0.75	2.32	1.08	1.12
Bought on the Internet	0.46	*	0.29	0.77
Some Other Way	0.44	1.31	0.47	0.67

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.55C Source Where Prescription Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2005**

Source for Most Recent Nonmedical Use among Past Year Prescription Stimulant Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	17	7	7	14
From More Than One Doctor	15	4	4	14
Wrote Fake Prescription	6	*	5	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	*	4	*
From Friend or Relative for Free	78	19	37	63
Bought from Friend or Relative	27	8	22	13
Took from Friend or Relative without Asking	18	10	8	13
Bought from Drug Dealer or Other Stranger	20	5	10	17
Bought on the Internet	39	4	6	*
Some Other Way	12	3	7	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Past Year Prescription Stimulant Use refers only to nonmedical prescription stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.55D Source Where Prescription Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2005**

Source for Most Recent Nonmedical Use among Past Year Prescription Stimulant Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	0.88	1.85	0.83	1.92
From More Than One Doctor	0.79	1.07	0.50	1.90
Wrote Fake Prescription	0.30	*	0.63	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.26	*	0.50	*
From Friend or Relative for Free	2.49	3.59	2.61	5.36
Bought from Friend or Relative	1.34	2.09	2.18	1.74
Took from Friend or Relative without Asking	0.94	2.74	0.90	1.74
Bought from Drug Dealer or Other Stranger	1.03	1.53	1.16	2.24
Bought on the Internet	1.89	1.19	0.69	*
Some Other Way	0.62	0.79	0.88	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Past Year Prescription Stimulant Use refers only to nonmedical prescription stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.56C Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2005**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From Friend or Relative for Free	56	14	25	48
Bought from Friend or Relative	45	6	18	41
Took from Friend or Relative without Asking	16	3	3	*
Bought from Drug Dealer or Other Stranger	35	*	17	29
Bought on the Internet	11	*	5	*
Some Other Way	12	4	4	10

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.56D Source Where Methamphetamine Was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2005**

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From Friend or Relative for Free	3.37	5.60	3.93	5.70
Bought from Friend or Relative	3.04	3.95	3.28	5.31
Took from Friend or Relative without Asking	1.26	1.96	0.72	*
Bought from Drug Dealer or Other Stranger	2.51	*	3.21	4.20
Bought on the Internet	0.88	*	1.02	*
Some Other Way	0.95	2.56	0.92	1.62

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.



**Table 7.57C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:  
Standard Errors of Numbers in Thousands, 2005**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	19	*	5	*
From More Than One Doctor	9	*	*	*
Wrote Fake Prescription	1	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	*	4	*
From Friend or Relative for Free	65	*	12	*
Bought from Friend or Relative	10	*	6	*
Took from Friend or Relative without Asking	14	*	*	*
Bought from Drug Dealer or Other Stranger	14	*	6	*
Bought on the Internet	*	*	*	*
Some Other Way	6	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.

**Table 7.57D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2005**

Source for Most Recent Nonmedical Use among Past Year Sedative Users <sup>1</sup>	Total	AGE GROUP		
		12-17	18-25	26 or Older
From One Doctor	2.64	*	2.83	*
From More Than One Doctor	1.23	*	*	*
Wrote Fake Prescription	0.09	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.66	*	2.42	*
From Friend or Relative for Free	4.60	*	5.47	*
Bought from Friend or Relative	1.41	*	3.44	*
Took from Friend or Relative without Asking	1.98	*	*	*
Bought from Drug Dealer or Other Stranger	2.00	*	3.34	*
Bought on the Internet	*	*	*	*
Some Other Way	0.84	*	*	*

\*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

<sup>1</sup> Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2005.