

Table 7.68C Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
ILLCIT DRUGS²	147	132	13	13	145	131
Marijuana and Hashish	115	106	11	11	114	106
Cocaine	42	43	3	2	42	43
Crack	21	25	0	1	21	25
Heroin	8	8	1	2	8	8
Hallucinogens	27	19	3	2	27	18
LSD	7	5	0	2	7	5
PCP	4	3	1	*	4	3
Ecstasy	22	15	2	1	22	15
Inhalants	16	13	1	1	16	13
Nonmedical Use of Psychotherapeutics ³	85	83	7	7	85	83
Pain Relievers	71	68	7	7	71	68
OxyContin [®]	--	17	--	1	--	17
Tranquilizers	43	44	2	2	43	44
Stimulants	38	38	3	2	38	38
Methamphetamine	27	28	1	1	27	28
Sedatives	22	17	1	1	22	17
ILLCIT DRUGS OTHER THAN MARIJUANA²	97	94	8	8	97	93

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.68D Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
ILLCIT DRUGS²	0.22	0.20	0.54	0.51	0.23	0.21
Marijuana and Hashish	0.18	0.17	0.45	0.44	0.19	0.18
Cocaine	0.07	0.07	0.13	0.09	0.07	0.07
Crack	0.03	0.04	0.02	0.05	0.03	0.04
Heroin	0.01	0.01	0.03	0.07	0.01	0.01
Hallucinogens	0.04	0.03	0.11	0.09	0.04	0.03
LSD	0.01	0.01	0.02	0.07	0.01	0.01
PCP	0.01	0.00	0.06	*	0.01	0.01
Ecstasy	0.04	0.02	0.08	0.06	0.04	0.02
Inhalants	0.03	0.02	0.03	0.06	0.03	0.02
Nonmedical Use of Psychotherapeutics ³	0.14	0.13	0.31	0.29	0.14	0.14
Pain Relievers	0.11	0.11	0.28	0.28	0.12	0.11
OxyContin [®]	--	0.03	--	0.04	--	0.03
Tranquilizers	0.07	0.07	0.08	0.10	0.07	0.07
Stimulants	0.06	0.06	0.13	0.06	0.06	0.06
Methamphetamine	0.04	0.04	0.04	0.03	0.05	0.05
Sedatives	0.04	0.03	0.06	0.04	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.15	0.15	0.31	0.31	0.16	0.15

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.68P Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Drug	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
ILLCIT DRUGS²	0.0569		0.5799		0.0758	
Marijuana and Hashish	0.0396		0.3142		0.0650	
Cocaine	0.4031		0.7258		0.3828	
Crack	0.1178		0.3570		0.1260	
Heroin	0.7591		0.2844		0.8065	
Hallucinogens	0.0146		0.9689		0.0135	
LSD	0.4244		0.1610		0.2716	
PCP	0.2060		*		0.2315	
Ecstasy	0.0131		0.6446		0.0137	
Inhalants	0.7382		0.3985		0.6512	
Nonmedical Use of Psychotherapeutics ³	0.9071		0.8131		0.9106	
Pain Relievers	0.6792		0.4595		0.7164	
OxyContin [®]	--		--		--	
Tranquilizers	0.7962		0.7317		0.7789	
Stimulants	0.6101		0.3211		0.5856	
Methamphetamine	0.5587		0.7057		0.5567	
Sedatives	0.3962		0.8222		0.3965	
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.7493		0.6903		0.7479	

*Low precision; no estimate reported.

-- Not available.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.69C Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	147	132	13	13	145	131
AGE						
15-17	29	30	4	3	29	29
18-25	50	53	8	8	59	58
26-44	125	109	9	9	124	109
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	139	122	12	10	137	122
White	121	109	10	8	120	108
Black or African American	49	46	7	6	48	46
American Indian or Alaska Native	10	9	*	*	10	9
Native Hawaiian or Other Pacific Islander	8	*	*	*	8	*
Asian	22	12	*	*	22	12
Two or More Races	15	18	*	*	15	18
Hispanic or Latino	44	50	5	8	44	49
TRIMESTER²						
First	N/A	N/A	9	8	N/A	N/A
Second	N/A	N/A	6	8	N/A	N/A
Third	N/A	N/A	6	6	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.69D Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.22	0.20	0.54	0.51	0.23	0.21
AGE						
15-17	0.44	0.45	3.39	3.63	0.45	0.45
18-25	0.32	0.33	0.86	0.87	0.34	0.34
26-44	0.30	0.26	0.64	0.61	0.31	0.28
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.25	0.22	0.63	0.52	0.25	0.23
White	0.29	0.26	0.70	0.56	0.29	0.27
Black or African American	0.52	0.50	2.09	1.75	0.54	0.52
American Indian or Alaska Native	2.34	2.27	*	*	2.44	2.32
Native Hawaiian or Other Pacific Islander	3.89	*	*	*	4.01	*
Asian	0.72	0.38	*	*	0.76	0.40
Two or More Races	2.07	2.36	*	*	2.17	2.42
Hispanic or Latino	0.46	0.50	0.92	1.50	0.49	0.52
TRIMESTER²						
First	N/A	N/A	1.28	1.09	N/A	N/A
Second	N/A	N/A	0.61	0.97	N/A	N/A
Third	N/A	N/A	0.82	0.64	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.69P Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.0569		0.5799		0.0758	
AGE						
15-17	0.0049		0.9214		0.0028	
18-25	0.0528		0.6638		0.0822	
26-44	0.3431		0.9470		0.3686	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0257		0.5005		0.0369	
White	0.0705		0.6382		0.0877	
Black or African American	0.3637		0.6285		0.4671	
American Indian or Alaska Native	0.4791		*		0.3640	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0290		*		0.0361	
Two or More Races	0.2106		*		0.2480	
Hispanic or Latino	0.4814		0.9824		0.4915	
TRIMESTER²						
First	N/A		0.6514		N/A	
Second	N/A		0.9629		N/A	
Third	N/A		0.9631		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.70C Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	115	106	11	11	114	106
AGE						
15-17	26	25	3	3	25	25
18-25	46	47	7	7	53	50
26-44	94	87	7	8	93	87
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	111	99	10	8	110	99
White	97	90	9	8	97	90
Black or African American	41	36	4	3	40	36
American Indian or Alaska Native	8	8	*	*	8	8
Native Hawaiian or Other Pacific Islander	3	3	*	*	3	3
Asian	14	10	*	*	14	10
Two or More Races	12	12	*	*	12	12
Hispanic or Latino	31	36	4	7	31	35
TRIMESTER²						
First	N/A	N/A	7	7	N/A	N/A
Second	N/A	N/A	5	8	N/A	N/A
Third	N/A	N/A	6	4	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.70D Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.18	0.17	0.45	0.44	0.19	0.18
AGE						
15-17	0.40	0.38	2.91	3.50	0.40	0.39
18-25	0.30	0.29	0.76	0.73	0.31	0.31
26-44	0.23	0.22	0.52	0.55	0.24	0.23
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.20	0.19	0.52	0.42	0.21	0.20
White	0.23	0.22	0.63	0.50	0.24	0.23
Black or African American	0.46	0.39	1.38	0.99	0.47	0.41
American Indian or Alaska Native	2.02	1.98	*	*	2.15	2.02
Native Hawaiian or Other Pacific Islander	1.45	1.56	*	*	1.50	1.62
Asian	0.45	0.32	*	*	0.47	0.33
Two or More Races	1.82	1.68	*	*	1.92	1.73
Hispanic or Latino	0.34	0.37	0.81	1.43	0.35	0.38
TRIMESTER²						
First	N/A	N/A	1.05	0.93	N/A	N/A
Second	N/A	N/A	0.53	0.94	N/A	N/A
Third	N/A	N/A	0.81	0.39	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.70P Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.0396		0.3142		0.0650	
AGE						
15-17	0.0735		0.7902		0.0585	
18-25	0.0025		0.1963		0.0064	
26-44	0.4736		0.9926		0.5143	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0194		0.1941		0.0356	
White	0.0518		0.4554		0.0741	
Black or African American	0.1903		0.1667		0.3144	
American Indian or Alaska Native	0.7325		*		0.5705	
Native Hawaiian or Other Pacific Islander	0.6674		*		0.6422	
Asian	0.1688		*		0.2281	
Two or More Races	0.2838		*		0.3130	
Hispanic or Latino	0.5167		0.9676		0.5120	
TRIMESTER²						
First	N/A		0.4391		N/A	
Second	N/A		0.9430		N/A	
Third	N/A		0.4394		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.71C Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOBACCO PRODUCTS²	275	255	30	30	270	250
Cigarettes	269	251	29	29	265	246
Smokeless Tobacco	22	20	3	2	22	20
Cigars	68	68	7	6	66	67
Pipe Tobacco	19	23	4	1	19	23
ALCOHOL	394	364	24	32	391	360
Binge Alcohol Use ³	226	213	14	15	224	211
Heavy Alcohol Use ³	98	90	5	6	97	90

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.71D Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOBACCO PRODUCTS²	0.38	0.36	1.21	1.15	0.39	0.37
Cigarettes	0.37	0.35	1.18	1.14	0.38	0.36
Smokeless Tobacco	0.04	0.03	0.14	0.08	0.04	0.03
Cigars	0.11	0.11	0.28	0.22	0.11	0.11
Pipe Tobacco	0.03	0.04	0.16	0.05	0.03	0.04
ALCOHOL	0.38	0.38	0.94	1.19	0.39	0.39
Binge Alcohol Use ³	0.31	0.30	0.58	0.58	0.32	0.31
Heavy Alcohol Use ³	0.15	0.14	0.22	0.25	0.16	0.15

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.71P Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Substance	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOBACCO PRODUCTS²	0.0242		0.2652		0.0454	
Cigarettes	0.0155		0.4071		0.0245	
Smokeless Tobacco	0.9707		0.5328		0.9634	
Cigars	0.6315		0.3794		0.4558	
Pipe Tobacco	0.1233		0.1814		0.0829	
ALCOHOL	0.7920		0.1246		0.8400	
Binge Alcohol Use ³	0.8447		0.8142		0.7647	
Heavy Alcohol Use ³	0.9012		0.8075		0.9096	

*Low precision; no estimate reported.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.72C Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	269	251	29	29	265	246
AGE						
15-17	34	34	5	4	33	34
18-25	71	72	18	16	87	87
26-44	240	228	22	24	237	226
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	258	238	29	28	254	232
White	232	210	28	26	229	205
Black or African American	93	95	5	9	93	94
American Indian or Alaska Native	22	20	*	*	22	19
Native Hawaiian or Other Pacific Islander	13	*	*	*	13	*
Asian	45	36	*	*	45	36
Two or More Races	25	25	*	*	25	25
Hispanic or Latino	86	86	6	9	87	86
TRIMESTER²						
First	N/A	N/A	16	18	N/A	N/A
Second	N/A	N/A	17	11	N/A	N/A
Third	N/A	N/A	17	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.72D Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.37	0.35	1.18	1.14	0.38	0.36
AGE						
15-17	0.49	0.50	4.11	4.73	0.49	0.51
18-25	0.45	0.44	1.58	1.54	0.47	0.46
26-44	0.52	0.50	1.57	1.55	0.53	0.52
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.41	0.39	1.42	1.34	0.42	0.40
White	0.47	0.44	1.80	1.64	0.48	0.45
Black or African American	0.94	0.94	1.67	2.52	0.98	0.98
American Indian or Alaska Native	3.83	4.27	*	*	3.97	4.35
Native Hawaiian or Other Pacific Islander	5.41	*	*	*	5.55	*
Asian	1.41	1.08	*	*	1.50	1.13
Two or More Races	3.10	2.74	*	*	3.21	2.81
Hispanic or Latino	0.80	0.80	1.31	1.84	0.85	0.84
TRIMESTER²						
First	N/A	N/A	2.15	2.38	N/A	N/A
Second	N/A	N/A	1.75	1.41	N/A	N/A
Third	N/A	N/A	2.23	1.76	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.72P Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.0155		0.4071		0.0245	
AGE						
15-17	0.0021		0.5654		0.0019	
18-25	0.0266		0.5343		0.0363	
26-44	0.1404		0.8857		0.1801	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.0065		0.3136		0.0113	
White	0.0239		0.1584		0.0482	
Black or African American	0.7889		0.0579		0.7314	
American Indian or Alaska Native	0.1377		*		0.0526	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.0813		*		0.0796	
Two or More Races	0.7696		*		0.7545	
Hispanic or Latino	0.4325		0.8427		0.4404	
TRIMESTER²						
First	N/A		0.5857		N/A	
Second	N/A		0.2426		N/A	
Third	N/A		0.2038		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.73C Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	394	364	24	32	391	360
AGE						
15-17	42	43	4	3	42	42
18-25	81	76	11	10	140	122
26-44	343	331	21	30	340	328
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	364	346	22	28	361	341
White	333	312	19	24	328	309
Black or African American	125	124	4	10	124	122
American Indian or Alaska Native	22	19	*	*	21	19
Native Hawaiian or Other Pacific Islander	16	*	*	*	16	*
Asian	76	72	*	*	76	72
Two or More Races	34	29	*	*	34	29
Hispanic or Latino	145	120	10	15	143	118
TRIMESTER²						
First	N/A	N/A	17	22	N/A	N/A
Second	N/A	N/A	12	15	N/A	N/A
Third	N/A	N/A	11	16	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.73D Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.38	0.38	0.94	1.19	0.39	0.39
AGE						
15-17	0.58	0.58	3.53	3.76	0.59	0.58
18-25	0.52	0.47	1.05	1.02	0.54	0.48
26-44	0.54	0.55	1.43	1.85	0.55	0.56
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.42	0.41	1.07	1.31	0.43	0.42
White	0.46	0.45	1.26	1.49	0.46	0.45
Black or African American	1.01	0.99	1.39	2.78	1.05	1.01
American Indian or Alaska Native	3.64	4.38	*	*	3.81	4.48
Native Hawaiian or Other Pacific Islander	5.63	*	*	*	5.81	*
Asian	2.17	1.90	*	*	2.27	1.93
Two or More Races	3.37	2.66	*	*	3.50	2.71
Hispanic or Latino	1.05	0.96	1.88	2.79	1.12	0.99
TRIMESTER²						
First	N/A	N/A	2.36	2.67	N/A	N/A
Second	N/A	N/A	1.25	1.79	N/A	N/A
Third	N/A	N/A	1.39	1.66	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.73P Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.7920		0.1246		0.8400	
AGE						
15-17	0.2940		0.8950		0.2457	
18-25	0.9945		0.5727		0.8671	
26-44	0.6100		0.0496		0.6122	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.5203		0.0480		0.5627	
White	0.0806		0.1296		0.0825	
Black or African American	0.7293		0.0256		0.7137	
American Indian or Alaska Native	0.3529		*		0.2778	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2201		*		0.2605	
Two or More Races	0.0932		*		0.1566	
Hispanic or Latino	0.7090		0.6015		0.7296	
TRIMESTER²						
First	N/A		0.7635		N/A	
Second	N/A		0.0601		N/A	
Third	N/A		0.3663		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.74C Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	226	213	14	15	224	211
AGE						
15-17	32	32	2	2	32	31
18-25	73	71	6	7	99	91
26-44	190	184	12	12	188	183
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	214	200	14	12	213	199
White	183	174	10	9	182	173
Black or African American	80	76	2	9	79	75
American Indian or Alaska Native	18	16	*	*	17	16
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	34	41	*	*	32	41
Two or More Races	18	20	*	*	18	20
Hispanic or Latino	83	75	3	8	82	74
TRIMESTER²						
First	N/A	N/A	12	10	N/A	N/A
Second	N/A	N/A	5	7	N/A	N/A
Third	N/A	N/A	4	7	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.74D Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.31	0.30	0.58	0.58	0.32	0.31
AGE						
15-17	0.49	0.47	2.32	2.78	0.50	0.48
18-25	0.47	0.44	0.66	0.76	0.49	0.46
26-44	0.42	0.41	0.88	0.82	0.43	0.42
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.35	0.33	0.70	0.61	0.36	0.34
White	0.39	0.37	0.74	0.57	0.40	0.38
Black or African American	0.86	0.76	0.81	2.53	0.88	0.79
American Indian or Alaska Native	3.75	3.92	*	*	3.96	4.01
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	1.09	1.24	*	*	1.11	1.27
Two or More Races	2.47	2.56	*	*	2.60	2.63
Hispanic or Latino	0.75	0.71	0.66	1.55	0.80	0.75
TRIMESTER²						
First	N/A	N/A	1.75	1.37	N/A	N/A
Second	N/A	N/A	0.50	0.79	N/A	N/A
Third	N/A	N/A	0.49	0.76	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.74P Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.8447		0.8142		0.7647	
AGE						
15-17	0.9152		0.7118		0.9549	
18-25	0.5817		0.9142		0.6557	
26-44	0.8168		0.7828		0.9437	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8783		0.6160		0.7773	
White	0.6165		0.3762		0.5266	
Black or African American	0.4252		0.0308		0.4118	
American Indian or Alaska Native	0.6367		*		0.6172	
Native Hawaiian or Other Pacific Islander	*		*		*	
Asian	0.2636		*		0.1831	
Two or More Races	0.3319		*		0.4159	
Hispanic or Latino	0.7343		0.6209		0.7640	
TRIMESTER²						
First	N/A		0.1277		N/A	
Second	N/A		0.1989		N/A	
Third	N/A		0.2871		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.75C Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Numbers in Thousands, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	98	90	5	6	97	90
AGE						
15-17	13	14	0	1	13	14
18-25	42	41	2	2	47	45
26-44	80	77	5	6	80	76
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	94	87	5	5	93	87
White	83	80	5	5	83	79
Black or African American	31	29	1	1	31	29
American Indian or Alaska Native	8	6	*	*	7	6
Native Hawaiian or Other Pacific Islander	4	5	*	*	4	5
Asian	22	13	*	*	22	13
Two or More Races	8	8	*	*	8	8
Hispanic or Latino	24	25	1	*	24	25
TRIMESTER²						
First	N/A	N/A	4	5	N/A	N/A
Second	N/A	N/A	*	4	N/A	N/A
Third	N/A	N/A	1	1	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.75D Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: Standard Errors of Percentages, Annual Averages Based on 2002-2003 and 2004-2005

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.15	0.14	0.22	0.25	0.16	0.15
AGE						
15-17	0.22	0.23	0.09	1.51	0.22	0.23
18-25	0.27	0.25	0.26	0.26	0.28	0.27
26-44	0.20	0.19	0.35	0.38	0.20	0.20
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.17	0.16	0.28	0.24	0.18	0.17
White	0.20	0.20	0.34	0.31	0.21	0.21
Black or African American	0.37	0.33	0.26	0.22	0.39	0.35
American Indian or Alaska Native	2.05	1.46	*	*	1.95	1.49
Native Hawaiian or Other Pacific Islander	1.90	2.11	*	*	1.93	2.18
Asian	0.71	0.41	*	*	0.75	0.43
Two or More Races	1.16	1.16	*	*	1.22	1.20
Hispanic or Latino	0.25	0.25	0.19	*	0.27	0.27
TRIMESTER²						
First	N/A	N/A	0.54	0.63	N/A	N/A
Second	N/A	N/A	*	0.53	N/A	N/A
Third	N/A	N/A	0.08	0.07	N/A	N/A

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 7.75P Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status and Demographic Characteristics: *P* Values from Tests of Differences of Percentages, 2004-2005 Versus 2002-2003

Demographic Characteristic	Total ¹		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2002-2003	2004-2005	2002-2003	2004-2005	2002-2003	2004-2005
TOTAL	0.9012		0.8075		0.9096	
AGE						
15-17	0.7416		0.0607		0.6279	
18-25	0.9968		0.9630		1.0000	
26-44	0.7965		0.9656		0.8061	
HISPANIC ORIGIN AND RACE						
Not Hispanic or Latino	0.8214		0.7731		0.8440	
White	0.8053		0.9711		0.7946	
Black or African American	0.8234		0.6206		0.8814	
American Indian or Alaska Native	0.6318		*		0.6253	
Native Hawaiian or Other Pacific Islander	0.5719		*		0.6163	
Asian	0.2718		*		0.2688	
Two or More Races	0.3871		*		0.4361	
Hispanic or Latino	0.4863		*		0.5082	
TRIMESTER²						
First	N/A		0.9915		N/A	
Second	N/A		*		N/A	
Third	N/A		0.8429		N/A	

*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

¹ Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

² Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.