

**Table 8.22A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Numbers in Thousands, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	71,499	70,757	70,257	71,519
Cigarettes	61,136	60,434	59,896	60,532
Smokeless Tobacco	7,787	7,725	7,154	7,682
Cigars	12,751	12,837	13,727	13,640
Pipe Tobacco	1,816	1,619 <sup>b</sup>	1,835	2,190
<b>ALCOHOL</b>	119,820 <sup>b</sup>	118,965 <sup>b</sup>	120,934 <sup>b</sup>	126,028
Binge Alcohol Use <sup>2</sup>	53,787	53,770	54,725	55,090
Heavy Alcohol Use <sup>2</sup>	15,860	16,144	16,689	16,035
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	41,991	41,288	41,569	42,175
Cigarettes	32,636	32,263	32,278	32,312
Smokeless Tobacco	7,242	7,096	6,730	7,174
Cigars	10,669	10,372 <sup>a</sup>	11,375	11,355
Pipe Tobacco	1,487 <sup>a</sup>	1,400 <sup>a</sup>	1,579	1,877
<b>ALCOHOL</b>	65,210 <sup>b</sup>	65,927 <sup>b</sup>	66,317 <sup>a</sup>	68,497
Binge Alcohol Use <sup>2</sup>	35,456	35,565	36,195	36,025
Heavy Alcohol Use <sup>2</sup>	12,216	11,958	12,388	12,172
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	29,509	29,469	28,688	29,344
Cigarettes	28,500	28,171	27,618	28,220
Smokeless Tobacco	545	628	424	508
Cigars	2,082	2,465	2,352	2,285
Pipe Tobacco	330	219	256	313
<b>ALCOHOL</b>	54,610 <sup>b</sup>	53,038 <sup>b</sup>	54,616 <sup>b</sup>	57,531
Binge Alcohol Use <sup>2</sup>	18,331	18,205	18,530	19,065
Heavy Alcohol Use <sup>2</sup>	3,645	4,186	4,301	3,863

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.22B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Percentages, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	30.4 <sup>a</sup>	29.8	29.2	29.4
Cigarettes	26.0 <sup>a</sup>	25.4	24.9	24.9
Smokeless Tobacco	3.3	3.3	3.0	3.2
Cigars	5.4	5.4	5.7	5.6
Pipe Tobacco	0.8	0.7 <sup>b</sup>	0.8	0.9
<b>ALCOHOL</b>	51.0	50.1 <sup>b</sup>	50.3 <sup>b</sup>	51.8
Binge Alcohol Use <sup>2</sup>	22.9	22.6	22.8	22.7
Heavy Alcohol Use <sup>2</sup>	6.7	6.8	6.9	6.6
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	37.0	35.9	35.7	35.8
Cigarettes	28.7 <sup>a</sup>	28.1	27.7	27.4
Smokeless Tobacco	6.4	6.2	5.8	6.1
Cigars	9.4	9.0	9.8	9.6
Pipe Tobacco	1.3	1.2 <sup>a</sup>	1.4	1.6
<b>ALCOHOL</b>	57.4	57.3	56.9	58.1
Binge Alcohol Use <sup>2</sup>	31.2	30.9	31.1	30.5
Heavy Alcohol Use <sup>2</sup>	10.8	10.4	10.6	10.3
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	24.3	24.0	23.1	23.4
Cigarettes	23.4	23.0	22.3	22.5
Smokeless Tobacco	0.4	0.5	0.3	0.4
Cigars	1.7	2.0	1.9	1.8
Pipe Tobacco	0.3	0.2	0.2	0.3
<b>ALCOHOL</b>	44.9	43.2 <sup>b</sup>	44.0 <sup>a</sup>	45.9
Binge Alcohol Use <sup>2</sup>	15.1	14.8	14.9	15.2
Heavy Alcohol Use <sup>2</sup>	3.0	3.4	3.5 <sup>a</sup>	3.1

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.23A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Numbers in Thousands, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	3,767 <sup>b</sup>	3,610 <sup>a</sup>	3,635 <sup>b</sup>	3,324
Cigarettes	3,209 <sup>b</sup>	3,044 <sup>b</sup>	2,998 <sup>a</sup>	2,733
Smokeless Tobacco	487	511	571	523
Cigars	1,117	1,115	1,201 <sup>a</sup>	1,064
Pipe Tobacco	137	160	178	158
<b>ALCOHOL</b>	4,365	4,425 <sup>a</sup>	4,436 <sup>a</sup>	4,191
Binge Alcohol Use <sup>2</sup>	2,641	2,658	2,794 <sup>b</sup>	2,513
Heavy Alcohol Use <sup>2</sup>	630	652	671	611
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	2,019 <sup>a</sup>	1,988	1,972	1,843
Cigarettes	1,556 <sup>a</sup>	1,514	1,453	1,387
Smokeless Tobacco	436	479	521	476
Cigars	790	789	852 <sup>a</sup>	748
Pipe Tobacco	89	121	110	109
<b>ALCOHOL</b>	2,200	2,185	2,216	2,057
Binge Alcohol Use <sup>2</sup>	1,441	1,421	1,496 <sup>a</sup>	1,341
Heavy Alcohol Use <sup>2</sup>	395	374	407	383
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	1,748 <sup>b</sup>	1,622	1,663 <sup>a</sup>	1,481
Cigarettes	1,653 <sup>b</sup>	1,530 <sup>b</sup>	1,545 <sup>b</sup>	1,345
Smokeless Tobacco	51	32	50	47
Cigars	327	326	349	316
Pipe Tobacco	49	40	68	49
<b>ALCOHOL</b>	2,165	2,241	2,220	2,134
Binge Alcohol Use <sup>2</sup>	1,200	1,237	1,298	1,172
Heavy Alcohol Use <sup>2</sup>	235	278	264	229

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.23B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Percentages, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	15.2 <sup>b</sup>	14.4 <sup>b</sup>	14.4 <sup>b</sup>	13.1
Cigarettes	13.0 <sup>b</sup>	12.2 <sup>b</sup>	11.9 <sup>b</sup>	10.8
Smokeless Tobacco	2.0	2.0	2.3	2.1
Cigars	4.5	4.5	4.8 <sup>a</sup>	4.2
Pipe Tobacco	0.6	0.6	0.7	0.6
<b>ALCOHOL</b>	17.6 <sup>a</sup>	17.7 <sup>a</sup>	17.6 <sup>a</sup>	16.5
Binge Alcohol Use <sup>2</sup>	10.7 <sup>a</sup>	10.6	11.1 <sup>b</sup>	9.9
Heavy Alcohol Use <sup>2</sup>	2.5	2.6	2.7	2.4
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	16.0 <sup>b</sup>	15.6 <sup>a</sup>	15.3	14.2
Cigarettes	12.3 <sup>b</sup>	11.9 <sup>a</sup>	11.3	10.7
Smokeless Tobacco	3.4	3.7	4.0	3.7
Cigars	6.2	6.2	6.6 <sup>a</sup>	5.8
Pipe Tobacco	0.7	0.9	0.9	0.8
<b>ALCOHOL</b>	17.4 <sup>a</sup>	17.1	17.2 <sup>a</sup>	15.9
Binge Alcohol Use <sup>2</sup>	11.4 <sup>a</sup>	11.1	11.6 <sup>a</sup>	10.4
Heavy Alcohol Use <sup>2</sup>	3.1	2.9	3.2	3.0
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	14.4 <sup>b</sup>	13.3 <sup>a</sup>	13.5 <sup>b</sup>	11.9
Cigarettes	13.6 <sup>b</sup>	12.5 <sup>b</sup>	12.5 <sup>b</sup>	10.8
Smokeless Tobacco	0.4	0.3	0.4	0.4
Cigars	2.7	2.7	2.8	2.5
Pipe Tobacco	0.4	0.3	0.5	0.4
<b>ALCOHOL</b>	17.9	18.3	18.0	17.2
Binge Alcohol Use <sup>2</sup>	9.9	10.1	10.5 <sup>a</sup>	9.4
Heavy Alcohol Use <sup>2</sup>	1.9	2.3	2.1	1.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Numbers in Thousands, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	14,049	14,201	14,359	14,382
Cigarettes	12,650	12,764	12,731	12,667
Smokeless Tobacco	1,502	1,486 <sup>a</sup>	1,589	1,664
Cigars	3,417 <sup>b</sup>	3,623 <sup>a</sup>	4,103	3,895
Pipe Tobacco	332 <sup>b</sup>	292 <sup>b</sup>	396	471
<b>ALCOHOL</b>	18,759 <sup>b</sup>	19,466	19,474	19,783
Binge Alcohol Use <sup>2</sup>	12,699 <sup>b</sup>	13,188	13,271	13,597
Heavy Alcohol Use <sup>2</sup>	4,625 <sup>a</sup>	4,788	4,848	4,967
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	8,097 <sup>a</sup>	8,226	8,368	8,425
Cigarettes	6,897	7,040	7,041	7,007
Smokeless Tobacco	1,455	1,418 <sup>a</sup>	1,532	1,588
Cigars	2,605 <sup>b</sup>	2,758 <sup>a</sup>	3,179	2,997
Pipe Tobacco	262 <sup>b</sup>	228 <sup>b</sup>	335	372
<b>ALCOHOL</b>	10,129 <sup>b</sup>	10,652	10,501 <sup>a</sup>	10,830
Binge Alcohol Use <sup>2</sup>	7,787 <sup>b</sup>	8,163	8,101 <sup>a</sup>	8,442
Heavy Alcohol Use <sup>2</sup>	3,282 <sup>a</sup>	3,372	3,434	3,545
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	5,952	5,975	5,990	5,957
Cigarettes	5,753	5,724	5,690	5,660
Smokeless Tobacco	47	67	56	75
Cigars	813	865	924	898
Pipe Tobacco	69	65	61	100
<b>ALCOHOL</b>	8,630 <sup>a</sup>	8,814	8,974	8,953
Binge Alcohol Use <sup>2</sup>	4,911	5,025	5,170	5,155
Heavy Alcohol Use <sup>2</sup>	1,343	1,417	1,414	1,422

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	45.3	44.8	44.6	44.3
Cigarettes	40.8 <sup>b</sup>	40.2	39.5	39.0
Smokeless Tobacco	4.8	4.7	4.9	5.1
Cigars	11.0 <sup>a</sup>	11.4	12.7	12.0
Pipe Tobacco	1.1 <sup>b</sup>	0.9 <sup>b</sup>	1.2	1.5
<b>ALCOHOL</b>	60.5	61.4	60.5	60.9
Binge Alcohol Use <sup>2</sup>	40.9	41.6	41.2	41.9
Heavy Alcohol Use <sup>2</sup>	14.9	15.1	15.1	15.3
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	52.1	51.7	51.7	51.6
Cigarettes	44.4	44.2	43.5	42.9
Smokeless Tobacco	9.4	8.9	9.5	9.7
Cigars	16.8 <sup>a</sup>	17.3	19.7	18.3
Pipe Tobacco	1.7 <sup>a</sup>	1.4 <sup>b</sup>	2.1	2.3
<b>ALCOHOL</b>	65.2	66.9	64.9	66.3
Binge Alcohol Use <sup>2</sup>	50.2	51.3	50.1	51.7
Heavy Alcohol Use <sup>2</sup>	21.1	21.2	21.2	21.7
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	38.4	37.8	37.4	36.9
Cigarettes	37.1 <sup>a</sup>	36.2	35.5	35.0
Smokeless Tobacco	0.3	0.4	0.4	0.5
Cigars	5.2	5.5	5.8	5.6
Pipe Tobacco	0.4	0.4	0.4	0.6
<b>ALCOHOL</b>	55.7	55.8	56.0	55.4
Binge Alcohol Use <sup>2</sup>	31.7	31.8	32.3	31.9
Heavy Alcohol Use <sup>2</sup>	8.7	9.0	8.8	8.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	53,683	52,946	52,263	53,813
Cigarettes	45,278	44,626	44,166	45,133
Smokeless Tobacco	5,797	5,728	4,994	5,495
Cigars	8,217	8,099	8,423	8,681
Pipe Tobacco	1,347	1,166 <sup>a</sup>	1,261	1,561
<b>ALCOHOL</b>	96,696 <sup>b</sup>	95,073 <sup>b</sup>	97,023 <sup>b</sup>	102,054
Binge Alcohol Use <sup>2</sup>	38,447	37,924	38,661	38,980
Heavy Alcohol Use <sup>2</sup>	10,605	10,704	11,170	10,457
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	31,874	31,074	31,229	31,907
Cigarettes	24,184	23,709	23,784	23,917
Smokeless Tobacco	5,351	5,199	4,677	5,110
Cigars	7,274	6,825	7,344	7,610
Pipe Tobacco	1,136	1,051	1,133	1,396
<b>ALCOHOL</b>	52,880 <sup>b</sup>	53,090 <sup>b</sup>	53,601 <sup>a</sup>	55,610
Binge Alcohol Use <sup>2</sup>	26,227	25,981	26,598	26,242
Heavy Alcohol Use <sup>2</sup>	8,539	8,212	8,548	8,245
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	21,809	21,872	21,034	21,906
Cigarettes	21,094	20,918	20,382	21,215
Smokeless Tobacco	447	529	318	385
Cigars	942	1,274	1,079	1,071
Pipe Tobacco	212	115	128	165
<b>ALCOHOL</b>	43,816 <sup>b</sup>	41,983 <sup>b</sup>	43,423 <sup>b</sup>	46,444
Binge Alcohol Use <sup>2</sup>	12,220	11,943	12,063	12,738
Heavy Alcohol Use <sup>2</sup>	2,067	2,492	2,623	2,212

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Percentages, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	29.9	29.3	28.5	29.0
Cigarettes	25.2	24.7	24.1	24.3
Smokeless Tobacco	3.2	3.2	2.7	3.0
Cigars	4.6	4.5	4.6	4.7
Pipe Tobacco	0.8	0.6	0.7	0.8
<b>ALCOHOL</b>	53.9	52.5 <sup>b</sup>	53.0 <sup>b</sup>	55.1
Binge Alcohol Use <sup>2</sup>	21.4	21.0	21.1	21.0
Heavy Alcohol Use <sup>2</sup>	5.9	5.9	6.1	5.6
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	37.3	36.0	35.7	36.0
Cigarettes	28.3	27.5	27.2	27.0
Smokeless Tobacco	6.3	6.0	5.3	5.8
Cigars	8.5	7.9	8.4	8.6
Pipe Tobacco	1.3	1.2	1.3	1.6
<b>ALCOHOL</b>	61.9	61.5	61.3	62.7
Binge Alcohol Use <sup>2</sup>	30.7	30.1	30.4	29.6
Heavy Alcohol Use <sup>2</sup>	10.0	9.5	9.8	9.3
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	23.2	23.1	22.0	22.6
Cigarettes	22.5	22.1	21.3	21.9
Smokeless Tobacco	0.5	0.6	0.3	0.4
Cigars	1.0	1.3	1.1	1.1
Pipe Tobacco	0.2	0.1	0.1	0.2
<b>ALCOHOL</b>	46.6	44.3 <sup>b</sup>	45.4 <sup>b</sup>	48.0
Binge Alcohol Use <sup>2</sup>	13.0	12.6	12.6	13.2
Heavy Alcohol Use <sup>2</sup>	2.2	2.6	2.7 <sup>a</sup>	2.3

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.



**Table 8.26A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	67,732	67,147	66,622	68,195
Cigarettes	57,928	57,391	56,897	57,799
Smokeless Tobacco	7,299	7,214	6,583	7,158
Cigars	11,634	11,722	12,526	12,576
Pipe Tobacco	1,679	1,458 <sup>b</sup>	1,657	2,032
<b>ALCOHOL</b>	115,455 <sup>b</sup>	114,539 <sup>b</sup>	116,498 <sup>b</sup>	121,837
Binge Alcohol Use <sup>2</sup>	51,146	51,112	51,931	52,577
Heavy Alcohol Use <sup>2</sup>	15,231	15,492	16,018	15,424
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	39,972	39,300	39,597	40,332
Cigarettes	31,080	30,749	30,825	30,924
Smokeless Tobacco	6,806	6,618	6,209	6,698
Cigars	9,879	9,583 <sup>a</sup>	10,523	10,607
Pipe Tobacco	1,398	1,279 <sup>b</sup>	1,469	1,768
<b>ALCOHOL</b>	63,009 <sup>b</sup>	63,742 <sup>b</sup>	64,101 <sup>b</sup>	66,440
Binge Alcohol Use <sup>2</sup>	34,014	34,144	34,699	34,684
Heavy Alcohol Use <sup>2</sup>	11,821	11,584	11,981	11,790
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	27,761	27,847	27,025	27,863
Cigarettes	26,847	26,641	26,072	26,875
Smokeless Tobacco	493	596	374	460
Cigars	1,755	2,139	2,003	1,969
Pipe Tobacco	281	179	189	264
<b>ALCOHOL</b>	52,446 <sup>b</sup>	50,797 <sup>b</sup>	52,396 <sup>b</sup>	55,397
Binge Alcohol Use <sup>2</sup>	17,132	16,968	17,232	17,893
Heavy Alcohol Use <sup>2</sup>	3,409	3,908	4,037	3,634

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.26B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Percentages, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	32.2	31.6	30.9	31.3
Cigarettes	27.5	27.0	26.4	26.5
Smokeless Tobacco	3.5	3.4	3.1	3.3
Cigars	5.5	5.5	5.8	5.8
Pipe Tobacco	0.8	0.7 <sup>b</sup>	0.8	0.9
<b>ALCOHOL</b>	54.9	53.9 <sup>b</sup>	54.1 <sup>b</sup>	55.9
Binge Alcohol Use <sup>2</sup>	24.3	24.0	24.1	24.1
Heavy Alcohol Use <sup>2</sup>	7.2	7.3	7.4	7.1
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	39.6	38.4	38.2	38.4
Cigarettes	30.8	30.1	29.8	29.5
Smokeless Tobacco	6.7	6.5	6.0	6.4
Cigars	9.8	9.4	10.2	10.1
Pipe Tobacco	1.4	1.3 <sup>a</sup>	1.4	1.7
<b>ALCOHOL</b>	62.4	62.4	61.9	63.3
Binge Alcohol Use <sup>2</sup>	33.7	33.4	33.5	33.0
Heavy Alcohol Use <sup>2</sup>	11.7	11.3	11.6	11.2
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	25.4	25.2	24.2	24.7
Cigarettes	24.5	24.1	23.3	23.8
Smokeless Tobacco	0.5	0.5	0.3	0.4
Cigars	1.6	1.9	1.8	1.7
Pipe Tobacco	0.3	0.2	0.2	0.2
<b>ALCOHOL</b>	47.9	46.0 <sup>b</sup>	46.9 <sup>b</sup>	49.1
Binge Alcohol Use <sup>2</sup>	15.7	15.4	15.4	15.8
Heavy Alcohol Use <sup>2</sup>	3.1	3.5	3.6	3.2

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.