

Table 8.34A Nonmedical Use of Stimulants and Methamphetamine in the Past Year and Number of Days Used Stimulants and Methamphetamine Nonmedically in the Past Year among Past Year Users Aged 12 or Older: Numbers in Thousands, 2002-2005

Past Year Drug/Frequency of Use	2002	2003	2004	2005
STIMULANTS	3,181 ^a	2,751	2,918	2,771
Number of Days Used Nonmedically in Past Year among Past Year Users				
1-11	1,305 ^a	1,196	1,272	1,069
12-49	667	649	655	717
50-99	386	279	327	330
100-299	704	534	553	582
300 or More	118	93	112	73
METHAMPHETAMINE	1,541	1,315	1,440	1,297
Number of Days Used Nonmedically in Past Year among Past Year Users				
1-11	607 ^a	477	567	432
12-49	280	337	299	337
50-99	221	122	183	133
100-299	366	315	327	349
300 or More	65	64	64	46

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

Table 8.34B Nonmedical Use of Stimulants and Methamphetamine in the Past Year and Number of Days Used Stimulants and Methamphetamine Nonmedically in the Past Year among Past Year Users Aged 12 or Older: Percentages, 2002-2005

Past Year Drug/Frequency of Use	2002	2003	2004	2005
STIMULANTS	100.0	100.0	100.0	100.0
Number of Days Used Nonmedically in Past Year among Past Year Users				
1-11	41.0	43.5	43.6	38.6
12-49	21.0 ^a	23.6	22.4	25.9
50-99	12.1	10.1	11.2	11.9
100-299	22.1	19.4	18.9	21.0
300 or More	3.7	3.4	3.8	2.6
METHAMPHETAMINE	100.0	100.0	100.0	100.0
Number of Days Used Nonmedically in Past Year among Past Year Users				
1-11	39.4	36.3	39.4	33.3
12-49	18.2 ^a	25.7	20.8	26.0
50-99	14.4	9.3	12.7	10.3
100-299	23.8	23.9	22.7	26.9
300 or More	4.2	4.9	4.4	3.6

*Low precision; no estimate reported.

^a Difference between estimate and 2005 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2005 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.