

**Table 8.22C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	832	803	794	845
Cigarettes	795	789	768	777
Smokeless Tobacco	279	284	245	267
Cigars	350	331	323	354
Pipe Tobacco	166	131	147	146
<b>ALCOHOL</b>	989	919	957	975
Binge Alcohol Use <sup>2</sup>	732	682	698	701
Heavy Alcohol Use <sup>2</sup>	392	380	382	379
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	572	573	570	576
Cigarettes	527	545	537	515
Smokeless Tobacco	264	256	234	241
Cigars	327	305	290	324
Pipe Tobacco	139	127	139	141
<b>ALCOHOL</b>	656	599	626	621
Binge Alcohol Use <sup>2</sup>	540	517	513	529
Heavy Alcohol Use <sup>2</sup>	346	321	319	321
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	513	507	497	543
Cigarettes	501	488	491	538
Smokeless Tobacco	100	126	65	87
Cigars	114	123	122	112
Pipe Tobacco	90	35	43	42
<b>ALCOHOL</b>	644	600	664	637
Binge Alcohol Use <sup>2</sup>	415	382	390	402
Heavy Alcohol Use <sup>2</sup>	177	174	174	163

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.22D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: Standard Errors of Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.35	0.34	0.33	0.35
Cigarettes	0.34	0.33	0.32	0.32
Smokeless Tobacco	0.12	0.12	0.10	0.11
Cigars	0.15	0.14	0.13	0.15
Pipe Tobacco	0.07	0.06	0.06	0.06
<b>ALCOHOL</b>	0.42	0.39	0.40	0.40
Binge Alcohol Use <sup>2</sup>	0.31	0.29	0.29	0.29
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.16	0.16
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.50	0.50	0.49	0.49
Cigarettes	0.46	0.47	0.46	0.44
Smokeless Tobacco	0.23	0.22	0.20	0.20
Cigars	0.29	0.27	0.25	0.27
Pipe Tobacco	0.12	0.11	0.12	0.12
<b>ALCOHOL</b>	0.58	0.52	0.54	0.53
Binge Alcohol Use <sup>2</sup>	0.48	0.45	0.44	0.45
Heavy Alcohol Use <sup>2</sup>	0.30	0.28	0.27	0.27
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.41	0.40	0.43
Cigarettes	0.41	0.40	0.40	0.43
Smokeless Tobacco	0.08	0.10	0.05	0.07
Cigars	0.09	0.10	0.10	0.09
Pipe Tobacco	0.07	0.03	0.03	0.03
<b>ALCOHOL</b>	0.53	0.49	0.54	0.51
Binge Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.32
Heavy Alcohol Use <sup>2</sup>	0.15	0.14	0.14	0.13

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.22P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2005 Versus 2002, 2003, and 2004**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0436	0.4521	0.6856	
Cigarettes	0.0171	0.2427	0.9728	
Smokeless Tobacco	0.3429	0.5721	0.2195	
Cigars	0.3740	0.3038	0.6155	
Pipe Tobacco	0.1672	0.0073	0.1095	
<b>ALCOHOL</b>	0.1391	0.0016	0.0067	
Binge Alcohol Use <sup>2</sup>	0.5977	0.9463	0.8010	
Heavy Alcohol Use <sup>2</sup>	0.5053	0.3718	0.1202	
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0880	0.8384	0.9103	
Cigarettes	0.0374	0.3075	0.6257	
Smokeless Tobacco	0.3468	0.7714	0.2867	
Cigars	0.5510	0.1115	0.7124	
Pipe Tobacco	0.0981	0.0220	0.1633	
<b>ALCOHOL</b>	0.3815	0.3106	0.1259	
Binge Alcohol Use <sup>2</sup>	0.3126	0.5495	0.4053	
Heavy Alcohol Use <sup>2</sup>	0.2921	0.8434	0.4174	
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1560	0.3178	0.6229	
Cigarettes	0.1199	0.4550	0.6607	
Smokeless Tobacco	0.6907	0.3887	0.4675	
Cigars	0.3928	0.1672	0.5840	
Pipe Tobacco	0.7948	0.1057	0.3667	
<b>ALCOHOL</b>	0.1805	0.0001	0.0110	
Binge Alcohol Use <sup>2</sup>	0.7763	0.3982	0.5398	
Heavy Alcohol Use <sup>2</sup>	0.6669	0.0876	0.0450	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.23C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	81	80	80	78
Cigarettes	74	73	76	71
Smokeless Tobacco	27	30	32	29
Cigars	47	44	44	46
Pipe Tobacco	14	18	19	17
<b>ALCOHOL</b>	80	82	81	81
Binge Alcohol Use <sup>2</sup>	67	67	73	65
Heavy Alcohol Use <sup>2</sup>	31	33	36	32
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	59	56	55	59
Cigarettes	53	47	52	53
Smokeless Tobacco	26	29	30	28
Cigars	37	36	37	38
Pipe Tobacco	13	17	15	14
<b>ALCOHOL</b>	55	57	59	57
Binge Alcohol Use <sup>2</sup>	46	46	52	48
Heavy Alcohol Use <sup>2</sup>	24	23	27	27
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	51	52	53	50
Cigarettes	49	50	52	48
Smokeless Tobacco	9	6	11	8
Cigars	26	23	25	24
Pipe Tobacco	9	7	13	11
<b>ALCOHOL</b>	58	60	59	58
Binge Alcohol Use <sup>2</sup>	46	46	50	43
Heavy Alcohol Use <sup>2</sup>	19	23	23	20

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.23D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: Standard Errors of Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.33	0.32	0.32	0.31
Cigarettes	0.30	0.29	0.30	0.28
Smokeless Tobacco	0.11	0.12	0.13	0.12
Cigars	0.19	0.17	0.18	0.18
Pipe Tobacco	0.06	0.07	0.08	0.07
<b>ALCOHOL</b>	0.32	0.33	0.32	0.32
Binge Alcohol Use <sup>2</sup>	0.27	0.27	0.29	0.26
Heavy Alcohol Use <sup>2</sup>	0.12	0.13	0.14	0.13
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.44	0.43	0.46
Cigarettes	0.42	0.37	0.40	0.41
Smokeless Tobacco	0.21	0.23	0.23	0.22
Cigars	0.30	0.28	0.28	0.29
Pipe Tobacco	0.10	0.13	0.11	0.11
<b>ALCOHOL</b>	0.43	0.45	0.45	0.44
Binge Alcohol Use <sup>2</sup>	0.37	0.36	0.40	0.37
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.21	0.21
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.42	0.42	0.43	0.40
Cigarettes	0.40	0.41	0.42	0.39
Smokeless Tobacco	0.08	0.05	0.09	0.07
Cigars	0.21	0.19	0.20	0.19
Pipe Tobacco	0.07	0.06	0.10	0.09
<b>ALCOHOL</b>	0.48	0.49	0.48	0.47
Binge Alcohol Use <sup>2</sup>	0.38	0.38	0.41	0.35
Heavy Alcohol Use <sup>2</sup>	0.16	0.19	0.19	0.16

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.23P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Gender: *P* Values from Tests of Differences of Percentages, 2005 Versus 2002, 2003, and 2004**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0028	0.0031	
Cigarettes	0.0000	0.0006	0.0069	
Smokeless Tobacco	0.5548	0.9008	0.2407	
Cigars	0.2271	0.2930	0.0242	
Pipe Tobacco	0.4362	0.8574	0.4228	
<b>ALCOHOL</b>	0.0153	0.0105	0.0190	
Binge Alcohol Use <sup>2</sup>	0.0418	0.0511	0.0027	
Heavy Alcohol Use <sup>2</sup>	0.4475	0.2790	0.1856	
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0082	0.0359	0.0851	
Cigarettes	0.0065	0.0375	0.3191	
Smokeless Tobacco	0.4533	0.8166	0.2401	
Cigars	0.2556	0.3273	0.0394	
Pipe Tobacco	0.3313	0.5325	0.9199	
<b>ALCOHOL</b>	0.0139	0.0503	0.0361	
Binge Alcohol Use <sup>2</sup>	0.0451	0.1320	0.0212	
Heavy Alcohol Use <sup>2</sup>	0.5610	0.9280	0.4808	
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0000	0.0231	0.0092	
Cigarettes	0.0000	0.0032	0.0033	
Smokeless Tobacco	0.6950	0.1435	0.8330	
Cigars	0.5868	0.6478	0.3111	
Pipe Tobacco	0.9707	0.4911	0.2684	
<b>ALCOHOL</b>	0.3152	0.0980	0.2402	
Binge Alcohol Use <sup>2</sup>	0.3714	0.1900	0.0461	
Heavy Alcohol Use <sup>2</sup>	0.6548	0.0768	0.2305	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Numbers in Thousands, 2002-2005**

<b>Gender/Substance</b>	<b>2002</b>	<b>2003</b>	<b>2004</b>	<b>2005</b>
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	150	151	161	155
Cigarettes	150	151	159	152
Smokeless Tobacco	59	56	60	64
Cigars	83	84	97	92
Pipe Tobacco	24	25	30	37
<b>ALCOHOL</b>	164	160	164	166
Binge Alcohol Use <sup>2</sup>	161	155	166	171
Heavy Alcohol Use <sup>2</sup>	112	114	120	120
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	100	106	106	110
Cigarettes	99	109	105	107
Smokeless Tobacco	56	55	58	60
Cigars	73	72	83	79
Pipe Tobacco	22	21	27	33
<b>ALCOHOL</b>	111	100	110	101
Binge Alcohol Use <sup>2</sup>	110	105	113	111
Heavy Alcohol Use <sup>2</sup>	87	91	88	96
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	101	93	102	101
Cigarettes	102	92	101	101
Smokeless Tobacco	8	11	10	12
Cigars	38	43	44	41
Pipe Tobacco	12	12	12	16
<b>ALCOHOL</b>	102	114	104	112
Binge Alcohol Use <sup>2</sup>	99	100	96	104
Heavy Alcohol Use <sup>2</sup>	55	59	60	56

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: Standard Errors of Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.48	0.48	0.50	0.48
Cigarettes	0.48	0.47	0.49	0.47
Smokeless Tobacco	0.19	0.18	0.19	0.20
Cigars	0.27	0.26	0.30	0.28
Pipe Tobacco	0.08	0.08	0.09	0.11
<b>ALCOHOL</b>	0.53	0.50	0.51	0.51
Binge Alcohol Use <sup>2</sup>	0.52	0.49	0.52	0.53
Heavy Alcohol Use <sup>2</sup>	0.36	0.36	0.37	0.37
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.64	0.67	0.66	0.67
Cigarettes	0.64	0.69	0.65	0.66
Smokeless Tobacco	0.36	0.34	0.36	0.37
Cigars	0.47	0.45	0.51	0.48
Pipe Tobacco	0.14	0.13	0.17	0.20
<b>ALCOHOL</b>	0.71	0.63	0.68	0.62
Binge Alcohol Use <sup>2</sup>	0.71	0.66	0.70	0.68
Heavy Alcohol Use <sup>2</sup>	0.56	0.57	0.54	0.59
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.59	0.64	0.63
Cigarettes	0.66	0.58	0.63	0.63
Smokeless Tobacco	0.05	0.07	0.06	0.08
Cigars	0.25	0.27	0.27	0.26
Pipe Tobacco	0.08	0.08	0.08	0.10
<b>ALCOHOL</b>	0.66	0.72	0.65	0.69
Binge Alcohol Use <sup>2</sup>	0.64	0.63	0.60	0.65
Heavy Alcohol Use <sup>2</sup>	0.35	0.37	0.37	0.35

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.



**Table 8.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Gender: *P* Values from Tests of Differences of Percentages, 2005 Versus 2002, 2003, and 2004**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1362	0.4696	0.6334	
Cigarettes	0.0081	0.0633	0.4147	
Smokeless Tobacco	0.3064	0.0973	0.4928	
Cigars	0.0129	0.1408	0.0685	
Pipe Tobacco	0.0054	0.0001	0.1339	
<b>ALCOHOL</b>	0.5589	0.5250	0.5735	
Binge Alcohol Use <sup>2</sup>	0.2127	0.6875	0.3902	
Heavy Alcohol Use <sup>2</sup>	0.4602	0.7017	0.6611	
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.5430	0.9209	0.8589	
Cigarettes	0.0973	0.1631	0.4885	
Smokeless Tobacco	0.4974	0.1065	0.6311	
Cigars	0.0198	0.1212	0.0637	
Pipe Tobacco	0.0173	0.0005	0.4444	
<b>ALCOHOL</b>	0.2610	0.4913	0.1370	
Binge Alcohol Use <sup>2</sup>	0.1193	0.6661	0.1040	
Heavy Alcohol Use <sup>2</sup>	0.4894	0.5251	0.5586	
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.0913	0.2877	0.5705	
Cigarettes	0.0222	0.1739	0.5941	
Smokeless Tobacco	0.0727	0.7053	0.2438	
Cigars	0.3766	0.8128	0.5755	
Pipe Tobacco	0.1772	0.1008	0.0602	
<b>ALCOHOL</b>	0.7873	0.7452	0.5435	
Binge Alcohol Use <sup>2</sup>	0.8063	0.8891	0.6909	
Heavy Alcohol Use <sup>2</sup>	0.7777	0.7564	0.9622	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	797	748	752	798
Cigarettes	754	738	718	726
Smokeless Tobacco	270	274	236	256
Cigars	331	332	311	335
Pipe Tobacco	160	126	142	141
<b>ALCOHOL</b>	953	887	933	936
Binge Alcohol Use <sup>2</sup>	702	639	654	650
Heavy Alcohol Use <sup>2</sup>	355	349	356	347
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	553	546	552	554
Cigarettes	506	521	512	493
Smokeless Tobacco	252	247	226	232
Cigars	311	302	280	309
Pipe Tobacco	132	124	135	136
<b>ALCOHOL</b>	630	577	610	598
Binge Alcohol Use <sup>2</sup>	527	492	492	507
Heavy Alcohol Use <sup>2</sup>	323	306	307	302
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	488	477	469	515
Cigarettes	475	460	466	508
Smokeless Tobacco	99	125	63	85
Cigars	99	117	107	98
Pipe Tobacco	89	32	39	37
<b>ALCOHOL</b>	628	582	654	615
Binge Alcohol Use <sup>2</sup>	393	347	361	370
Heavy Alcohol Use <sup>2</sup>	158	155	160	146

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: Standard Errors of Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.44	0.41	0.41	0.43
Cigarettes	0.42	0.41	0.39	0.39
Smokeless Tobacco	0.15	0.15	0.13	0.14
Cigars	0.18	0.18	0.17	0.18
Pipe Tobacco	0.09	0.07	0.08	0.08
<b>ALCOHOL</b>	0.53	0.49	0.51	0.51
Binge Alcohol Use <sup>2</sup>	0.39	0.35	0.36	0.35
Heavy Alcohol Use <sup>2</sup>	0.20	0.19	0.19	0.19
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.65	0.63	0.63	0.62
Cigarettes	0.59	0.60	0.59	0.56
Smokeless Tobacco	0.29	0.29	0.26	0.26
Cigars	0.36	0.35	0.32	0.35
Pipe Tobacco	0.15	0.14	0.15	0.15
<b>ALCOHOL</b>	0.74	0.67	0.70	0.67
Binge Alcohol Use <sup>2</sup>	0.62	0.57	0.56	0.57
Heavy Alcohol Use <sup>2</sup>	0.38	0.35	0.35	0.34
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.52	0.50	0.49	0.53
Cigarettes	0.51	0.49	0.49	0.53
Smokeless Tobacco	0.11	0.13	0.07	0.09
Cigars	0.11	0.12	0.11	0.10
Pipe Tobacco	0.10	0.03	0.04	0.04
<b>ALCOHOL</b>	0.67	0.61	0.68	0.64
Binge Alcohol Use <sup>2</sup>	0.42	0.37	0.38	0.38
Heavy Alcohol Use <sup>2</sup>	0.17	0.16	0.17	0.15

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2005 Versus 2002, 2003, and 2004**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1458	0.7000	0.4141	
Cigarettes	0.1188	0.5776	0.6842	
Smokeless Tobacco	0.1899	0.3261	0.2108	
Cigars	0.6932	0.4215	0.7390	
Pipe Tobacco	0.4392	0.0557	0.1582	
<b>ALCOHOL</b>	0.1197	0.0004	0.0041	
Binge Alcohol Use <sup>2</sup>	0.4375	0.8883	0.8626	
Heavy Alcohol Use <sup>2</sup>	0.3194	0.3086	0.0895	
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1442	0.9892	0.7530	
Cigarettes	0.1029	0.5497	0.7857	
Smokeless Tobacco	0.2060	0.5012	0.2585	
Cigars	0.8895	0.1715	0.6956	
Pipe Tobacco	0.2593	0.0903	0.2005	
<b>ALCOHOL</b>	0.4022	0.2006	0.1400	
Binge Alcohol Use <sup>2</sup>	0.1929	0.5345	0.3098	
Heavy Alcohol Use <sup>2</sup>	0.1731	0.6631	0.3327	
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.4419	0.5293	0.3631	
Cigarettes	0.4713	0.8157	0.3822	
Smokeless Tobacco	0.5729	0.3124	0.5484	
Cigars	0.4762	0.1358	0.8927	
Pipe Tobacco	0.5888	0.3412	0.5118	
<b>ALCOHOL</b>	0.1395	0.0000	0.0051	
Binge Alcohol Use <sup>2</sup>	0.7801	0.2984	0.2988	
Heavy Alcohol Use <sup>2</sup>	0.7010	0.1219	0.0440	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.26C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Numbers in Thousands, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	829	784	793	839
Cigarettes	790	774	765	772
Smokeless Tobacco	276	283	243	264
Cigars	346	333	323	351
Pipe Tobacco	163	130	145	145
<b>ALCOHOL</b>	972	909	950	969
Binge Alcohol Use <sup>2</sup>	730	675	699	701
Heavy Alcohol Use <sup>2</sup>	389	376	383	378
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	565	561	568	576
Cigarettes	521	537	532	514
Smokeless Tobacco	260	255	233	239
Cigars	322	306	291	324
Pipe Tobacco	136	127	138	141
<b>ALCOHOL</b>	643	590	616	616
Binge Alcohol Use <sup>2</sup>	540	517	511	529
Heavy Alcohol Use <sup>2</sup>	346	320	318	322
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	509	498	494	541
Cigarettes	497	481	488	537
Smokeless Tobacco	99	126	64	87
Cigars	110	121	119	109
Pipe Tobacco	90	35	41	40
<b>ALCOHOL</b>	640	593	662	635
Binge Alcohol Use <sup>2</sup>	411	375	389	401
Heavy Alcohol Use <sup>2</sup>	175	170	173	162

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.26D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: Standard Errors of Percentages, 2002-2005**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.39	0.37	0.37	0.39
Cigarettes	0.38	0.36	0.36	0.35
Smokeless Tobacco	0.13	0.13	0.11	0.12
Cigars	0.16	0.16	0.15	0.16
Pipe Tobacco	0.08	0.06	0.07	0.07
<b>ALCOHOL</b>	0.46	0.43	0.44	0.44
Binge Alcohol Use <sup>2</sup>	0.35	0.32	0.32	0.32
Heavy Alcohol Use <sup>2</sup>	0.19	0.18	0.18	0.17
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.56	0.55	0.55	0.55
Cigarettes	0.52	0.53	0.51	0.49
Smokeless Tobacco	0.26	0.25	0.22	0.23
Cigars	0.32	0.30	0.28	0.31
Pipe Tobacco	0.13	0.12	0.13	0.13
<b>ALCOHOL</b>	0.64	0.58	0.59	0.59
Binge Alcohol Use <sup>2</sup>	0.53	0.51	0.49	0.50
Heavy Alcohol Use <sup>2</sup>	0.34	0.31	0.31	0.31
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.47	0.45	0.44	0.48
Cigarettes	0.45	0.44	0.44	0.48
Smokeless Tobacco	0.09	0.11	0.06	0.08
Cigars	0.10	0.11	0.11	0.10
Pipe Tobacco	0.08	0.03	0.04	0.04
<b>ALCOHOL</b>	0.59	0.54	0.59	0.56
Binge Alcohol Use <sup>2</sup>	0.38	0.34	0.35	0.36
Heavy Alcohol Use <sup>2</sup>	0.16	0.15	0.16	0.14

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.

**Table 8.26P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 or Older, by Gender: *P* Values from Tests of Differences of Percentages, 2005 Versus 2002, 2003, and 2004**

Gender/Substance	2002	2003	2004	2005
<b>TOTAL</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1057	0.6134	0.5022	
Cigarettes	0.0523	0.3717	0.8377	
Smokeless Tobacco	0.3033	0.5553	0.1681	
Cigars	0.2927	0.2463	0.8356	
Pipe Tobacco	0.1895	0.0064	0.0864	
<b>ALCOHOL</b>	0.1032	0.0008	0.0039	
Binge Alcohol Use <sup>2</sup>	0.7076	0.8230	0.9785	
Heavy Alcohol Use <sup>2</sup>	0.5298	0.4102	0.1477	
<b>MALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.1353	0.9722	0.7941	
Cigarettes	0.0624	0.3854	0.6796	
Smokeless Tobacco	0.2940	0.7819	0.2255	
Cigars	0.4725	0.0905	0.8999	
Pipe Tobacco	0.1160	0.0180	0.1593	
<b>ALCOHOL</b>	0.3100	0.2577	0.0889	
Binge Alcohol Use <sup>2</sup>	0.3752	0.6112	0.5229	
Heavy Alcohol Use <sup>2</sup>	0.2987	0.8172	0.4434	
<b>FEMALE</b>				
<b>TOBACCO PRODUCTS<sup>1</sup></b>	0.3044	0.4242	0.4566	
Cigarettes	0.2693	0.6306	0.4733	
Smokeless Tobacco	0.7170	0.3370	0.4469	
Cigars	0.3151	0.1885	0.7316	
Pipe Tobacco	0.7978	0.1334	0.2037	
<b>ALCOHOL</b>	0.1582	0.0001	0.0083	
Binge Alcohol Use <sup>2</sup>	0.7063	0.3191	0.3970	
Heavy Alcohol Use <sup>2</sup>	0.6289	0.1302	0.0621	

\*Low precision; no estimate reported.

<sup>1</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>2</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2002, 2003, 2004, and 2005.