

Table 1.74A (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	8,559	8,615	1,478	1,452	704	611
GEOGRAPHIC DIVISION						
Northeast	1,430	1,176	209	184	74	67
New England	419	300	48	55	22	14
Middle Atlantic	1,011	876	161	128	52	53
Midwest	1,789	1,776	360	361	181	141
East North Central	1,266	1,211	270	281	142	117
West North Central	523	566	89	80	38	25
South	2,898	3,158	594	639	307	285
South Atlantic	1,417	1,795	324	394	175	179
East South Central	541	539	152	103	74	51
West South Central	940	824	117	141	57	55
West	2,442	2,506	316	268	142	117
Mountain	776	773	60	100	20	43
Pacific	1,666	1,733	256	169	123	74
COUNTY TYPE						
Large Metro	4,759	4,865	768	748	447	298
Small Metro	2,563	2,360	483	445	176	245
250K – 1 Mil. Pop.	1,686	1,568	365	319	142	165
< 250K Pop.	877	792	117	126	35	80
Nonmetro	1,237	1,390	227	259	81	67
Urbanized	538	697	91	109	47	25
Less Urbanized	590	570	121	133	33	34
Completely Rural	109	123	15	17	*	9

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.74B (REVISED) Crack Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	3.5	3.5	0.6	0.6	0.3	0.2
GEOGRAPHIC DIVISION						
Northeast	3.1	2.6	0.5	0.4	0.2	0.1
New England	3.5	2.5	0.4	0.5	0.2	0.1
Middle Atlantic	3.0	2.6	0.5	0.4	0.2	0.2
Midwest	3.3	3.2	0.7	0.7	0.3	0.3
East North Central	3.3	3.2	0.7	0.7	0.4	0.3
West North Central	3.2	3.4	0.5	0.5	0.2	0.1
South	3.3	3.5	0.7	0.7	0.3	0.3
South Atlantic	3.0	3.8	0.7	0.8	0.4	0.4
East South Central	3.7	3.7	1.0	0.7	0.5	0.3
West South Central	3.4	3.0	0.4	0.5	0.2	0.2
West	4.3	4.4	0.6	0.5	0.3	0.2
Mountain	4.6	4.5	0.4	0.6	0.1	0.2
Pacific	4.2	4.3	0.6	0.4	0.3	0.2
COUNTY TYPE						
Large Metro	3.6	3.7	0.6	0.6	0.3	0.2
Small Metro	3.6	3.2	0.7	0.6	0.2	0.3
250K – 1 Mil. Pop.	3.6	3.2	0.8	0.6	0.3	0.3
< 250K Pop.	3.6	3.2	0.5	0.5	0.1	0.3
Nonmetro	2.9	3.4	0.5	0.6	0.2	0.2
Urbanized	3.3	4.1	0.6	0.7	0.3	0.1
Less Urbanized	2.8	2.9	0.6	0.7	0.2	0.2
Completely Rural	2.4	2.7	0.3	0.4	*	0.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.