

**Table 1.75A (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
**Numbers in Thousands, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
<b>TOTAL</b>	35,290	34,278	3,970	3,759	1,008	998
<b>GEOGRAPHIC DIVISION</b>						
Northeast	6,711	6,178	747	609	168	179
New England	2,181	1,830	250	172	48	40
Middle Atlantic	4,530	4,348	497	438	120	139
Midwest	7,158	7,113	875	756	217	176
East North Central	4,886	5,079	611	536	152	111
West North Central	2,272	2,034	264	220	65	65
South	11,057	10,868	1,303	1,315	321	351
South Atlantic	6,130	5,756	681	582	141	139
East South Central	1,453 <sup>a</sup>	1,813	222	230	55	61
West South Central	3,474	3,299	400	503	125	152
West	10,364	10,118	1,046	1,078	301	291
Mountain	3,300	3,228	320	380	100	107
Pacific	7,064	6,891	726	698	201	184
<b>COUNTY TYPE</b>						
Large Metro	20,331	19,021	2,277	2,040	584	576
Small Metro	10,059	10,595	1,175	1,226	310	320
250K – 1 Mil. Pop.	6,935	7,400	834	889	247	217
< 250K Pop.	3,125	3,195	341	337	63	103
Nonmetro	4,899	4,661	517	493	113	102
Urbanized	2,126	2,188	254	242	61	59
Less Urbanized	2,248	2,158	228	231	51	39
Completely Rural	525 <sup>a</sup>	315	35	20	1	4

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

<sup>a</sup> Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 1.75B (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
<b>TOTAL</b>	14.3	13.8	1.6	1.5	0.4	0.4
<b>GEOGRAPHIC DIVISION</b>						
Northeast	14.6	13.5	1.6	1.3	0.4	0.4
New England	18.1	15.2	2.1	1.4	0.4	0.3
Middle Atlantic	13.4	12.8	1.5	1.3	0.4	0.4
Midwest	13.1	13.0	1.6	1.4	0.4	0.3
East North Central	12.8	13.3	1.6	1.4	0.4	0.3
West North Central	13.8	12.3	1.6	1.3	0.4	0.4
South	12.4	12.1	1.5	1.5	0.4	0.4
South Atlantic	13.0	12.1	1.4	1.2	0.3	0.3
East South Central	9.9	12.3	1.5	1.6	0.4	0.4
West South Central	12.7	11.9	1.5	1.8	0.5	0.5
West	18.3	17.7	1.9	1.9	0.5	0.5
Mountain	19.6	18.7	1.9	2.2	0.6	0.6
Pacific	17.8	17.2	1.8	1.7	0.5	0.5
<b>COUNTY TYPE</b>						
Large Metro	15.3	14.3	1.7	1.5	0.4	0.4
Small Metro	14.2	14.3	1.7	1.7	0.4	0.4
250K – 1 Mil. Pop.	14.9	14.9	1.8	1.8	0.5	0.4
< 250K Pop.	12.9	13.0	1.4	1.4	0.3	0.4
Nonmetro	11.6	11.3	1.2	1.2	0.3	0.2
Urbanized	13.0	13.0	1.6	1.4	0.4	0.4
Less Urbanized	10.6	11.0	1.1	1.2	0.2	0.2
Completely Rural	11.4 <sup>b</sup>	6.8	0.8	0.4	0.0	0.1

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

<sup>a</sup> Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.