

Table 1.11A Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	111,774	114,275	35,775	35,692	20,357	19,857
12	472	398	269	216	123	107
13	674	675	412	419	192	163
14	994	885	676	608	299	278
15	1,447 ^a	1,273	1,037	939	504	483
16	1,614	1,673	1,227	1,272	645	657
17	1,807	1,717	1,351	1,272	718	720
18	2,299	2,236	1,724	1,658	950	987
19	2,291	2,337	1,553	1,606	934	978
20	2,403	2,332	1,538	1,495	950	898
21	2,649	2,463	1,544	1,539	864	944
22	2,459	2,261	1,355	1,242	772	757
23	2,443	2,482	1,219	1,250	716	692
24	2,448	2,446	1,233	1,130	701	636
25	2,319	2,239	1,086	944	596	572
26-29	9,971	9,484	4,067	3,772	2,321	2,092
30-34	10,287	10,493	3,250	3,201	1,860	1,772
35-39	11,740	11,811	2,973	2,931	1,679	1,543
40-44	13,626 ^a	12,437	3,046	2,773	1,860	1,480
45-49	13,569	13,778	2,615	2,689	1,477	1,615
50-54	11,391	12,794	1,901	2,297	1,252	1,242
55-59	7,601 ^b	9,505	851 ^b	1,472	418 ^a	751
60-64	3,783 ^a	4,658	461	588	288	251
65 or Older	3,487	3,899	389	378	239	238

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.11B Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	45.4	46.1	14.5	14.4	8.3	8.0
12	12.1 ^a	9.9	6.9 ^a	5.4	3.1	2.7
13	16.3	16.4	9.9	10.2	4.6	4.0
14	23.2	21.4	15.8	14.7	7.0	6.7
15	31.9 ^a	29.0	22.8	21.4	11.1	11.0
16	37.3	37.6	28.3	28.6	14.9	14.8
17	43.1	41.5	32.2	30.8	17.1	17.4
18	50.1	46.9	37.6	34.8	20.7	20.7
19	54.9	53.3	37.2	36.6	22.4	22.3
20	59.7	57.1	38.2	36.6	23.6	22.0
21	61.7	60.3	36.0	37.6	20.1	23.1
22	62.3	59.6	34.3	32.8	19.5	20.0
23	61.9	62.7	30.9	31.6	18.1	17.5
24	62.4	62.1	31.4	28.7	17.9	16.1
25	60.5	60.1	28.3	25.4	15.5	15.4
26-29	60.4	57.8	24.7	23.0	14.1	12.8
30-34	55.3	55.5	17.5	16.9	10.0	9.4
35-39	56.2	56.1	14.2	13.9	8.0	7.3
40-44	60.9	58.6	13.6	13.1	8.3	7.0
45-49	61.6	61.0	11.9	11.9	6.7	7.2
50-54	54.6	58.9	9.1	10.6	6.0	5.7
55-59	43.4 ^b	51.6	4.9 ^b	8.0	2.4	4.1
60-64	28.2 ^b	35.0	3.4	4.4	2.1	1.9
65 or Older	9.8	10.7	1.1	1.0	0.7	0.7

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.12A Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	97,825 ^a	100,518	25,378	25,085	14,813	14,448
12	55	51	40	41	12	23
13	190	167	136	120	59	52
14	460	410	363	321	159	154
15	937 ^b	787	723	630	348	334
16	1,259	1,224	974	970	514	506
17	1,493	1,444	1,112	1,065	596	617
18	1,974	1,883	1,433	1,348	783	814
19	1,983	2,050	1,309	1,376	780	829
20	2,136	2,075	1,295	1,287	812	786
21	2,377	2,201	1,283	1,267	705	773
22	2,209	2,014	1,096	1,037	634	631
23	2,222	2,256	1,001	992	614	578
24	2,192	2,174	933	921	538	499
25	2,064	1,966	820	756	456	467
26-29	8,745	8,528	2,837	2,851	1,674	1,600
30-34	9,040	9,220	2,181	2,102	1,307	1,199
35-39	10,541	10,494	1,965	1,874	1,203	1,145
40-44	12,420 ^a	11,340	1,952	1,704	1,185	949
45-49	12,577	12,849	1,653	1,775	1,068	1,110
50-54	10,593	11,910	1,329	1,465	865	822
55-59	6,938 ^a	8,507	479 ^a	818	225	395
60-64	3,096 ^a	3,936	291	255	202	85
65 or Older	2,324 ^a	3,031	174	108	72	78

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.12B Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	39.8	40.6	10.3	10.1	6.0	5.8
12	1.4	1.3	1.0	1.0	0.3	0.6
13	4.6	4.1	3.3	2.9	1.4	1.3
14	10.7	9.9	8.5	7.8	3.7	3.7
15	20.6 ^a	17.9	15.9	14.4	7.7	7.6
16	29.1	27.5	22.5	21.8	11.9	11.4
17	35.6	34.9	26.6	25.8	14.2	14.9
18	43.0 ^a	39.5	31.2	28.3	17.1	17.1
19	47.5	46.8	31.3	31.4	18.7	18.9
20	53.0	50.8	32.1	31.5	20.2	19.2
21	55.4	53.8	29.9	31.0	16.4	18.9
22	55.9	53.1	27.7	27.4	16.0	16.7
23	56.3	57.0	25.4	25.1	15.6	14.6
24	55.9	55.2	23.8	23.4	13.7	12.7
25	53.8	52.8	21.4	20.3	11.9	12.5
26-29	53.0	52.0	17.2	17.4	10.1	9.8
30-34	48.6	48.8	11.7	11.1	7.0	6.3
35-39	50.4	49.8	9.4	8.9	5.8	5.4
40-44	55.5	53.5	8.7	8.0	5.3	4.5
45-49	57.1	56.9	7.5	7.9	4.8	4.9
50-54	50.7	54.8	6.4	6.7	4.1	3.8
55-59	39.6 ^a	46.2	2.7 ^a	4.4	1.3	2.1
60-64	23.1 ^b	29.6	2.2	1.9	1.5	0.6
65 or Older	6.5	8.4	0.5	0.3	0.2	0.2

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.13A Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	35,298	35,882	6,069	5,738	2,421	2,075
12	4	6	2	3	1	2
13	13	14	8	14	2	5
14	31	26	27	23	16	6
15	95	78	75	63	19	12
16	152	169	114	108	31	32
17	274	250	184	173	39	47
18	474	384	341 ^a	242	121 ^b	61
19	513	490	330	280	109	85
20	584	569	307	299	106	83
21	668	593	289	286	100	84
22	696	657	240	306	54	65
23	694	784	240	266	75	69
24	784	753	271 ^a	201	88	65
25	718	688	240	203	76	60
26-29	3,191	2,947	846	718	316	246
30-34	3,207	2,910	703	543	292	252
35-39	4,006	4,011	578	431	264	166
40-44	6,155 ^a	5,272	552	403	306	188
45-49	5,724	6,131	291 ^a	547	148 ^a	311
50-54	4,097 ^a	5,291	236	397	137	114
55-59	2,091	2,589	130	159	78	68
60-64	781	854	55	2	35	*
65 or Older	346	418	9	71	7	53

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.13B Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	14.3	14.5	2.5	2.3	1.0	0.8
12	0.1	0.2	0.1	0.1	0.0	0.0
13	0.3	0.3	0.2	0.3	0.1	0.1
14	0.7	0.6	0.6	0.5	0.4	0.1
15	2.1	1.8	1.7	1.4	0.4	0.3
16	3.5	3.8	2.6	2.4	0.7	0.7
17	6.5	6.0	4.4	4.2	0.9	1.1
18	10.3 ^a	8.1	7.4 ^b	5.1	2.6 ^b	1.3
19	12.3	11.2	7.9	6.4	2.6	2.0
20	14.5	13.9	7.6	7.3	2.6	2.0
21	15.5	14.5	6.7	7.0	2.3	2.1
22	17.6	17.3	6.1 ^a	8.1	1.4	1.7
23	17.6	19.8	6.1	6.7	1.9	1.7
24	20.0	19.1	6.9 ^a	5.1	2.2	1.6
25	18.7	18.5	6.3	5.5	2.0	1.6
26-29	19.3	18.0	5.1	4.4	1.9	1.5
30-34	17.2	15.4	3.8	2.9	1.6	1.3
35-39	19.2	19.0	2.8	2.0	1.3	0.8
40-44	27.5	24.9	2.5	1.9	1.4	0.9
45-49	26.0	27.2	1.3 ^a	2.4	0.7 ^a	1.4
50-54	19.6 ^a	24.4	1.1	1.8	0.7	0.5
55-59	11.9	14.1	0.7	0.9	0.4	0.4
60-64	5.8	6.4	0.4	0.0	0.3	*
65 or Older	1.0	1.2	0.0	0.2	0.0	0.1

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.14A Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	8,554	8,581	1,479	1,451	702	610
12	*	*	*	*	*	*
13	0	5	*	3	*	0
14	9	9	9	7	3	2
15	25	18	14	16	1	3
16	37	31	21	21	4	6
17	56	43	28	17	4	4
18	131 ^b	57	74 ^b	22	11	7
19	95	85	32	30	7	9
20	136	102	32	40	8	7
21	133	128	30	24	9	3
22	150	149	16	36	2	9
23	164	144	39	30	16	7
24	197	156	32	37	14	6
25	170	160	38	28	5	10
26-29	921 ^a	669	148	115	92	59
30-34	950	885	123	89	46	37
35-39	1,308	1,076	187	172	107	77
40-44	1,644	1,407	244	186	140	103
45-49	1,051 ^b	1,554	132 ^a	297	73	174
50-54	747 ^a	1,232	165	170	84	42
55-59	320	470	84	76	45	33
60-64	195	77	23	*	23	*
65 or Older	116	125	7	32	7	13

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.14B Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	3.5	3.5	0.6	0.6	0.3	0.2
12	*	*	*	*	*	*
13	0.0	0.1	*	0.1	*	0.0
14	0.2	0.2	0.2	0.2	0.1	0.1
15	0.5	0.4	0.3	0.4	0.0	0.1
16	0.9	0.7	0.5	0.5	0.1	0.1
17	1.3	1.0	0.7	0.4	0.1	0.1
18	2.9 ^b	1.2	1.6 ^b	0.5	0.2	0.1
19	2.3	1.9	0.8	0.7	0.2	0.2
20	3.4	2.5	0.8	1.0	0.2	0.2
21	3.1	3.1	0.7	0.6	0.2	0.1
22	3.8	3.9	0.4 ^a	1.0	0.1	0.2
23	4.2	3.6	1.0	0.8	0.4	0.2
24	5.0	4.0	0.8	0.9	0.4	0.1
25	4.4	4.3	1.0	0.8	0.1	0.3
26-29	5.6 ^a	4.1	0.9	0.7	0.6	0.4
30-34	5.1	4.7	0.7	0.5	0.2	0.2
35-39	6.3	5.1	0.9	0.8	0.5	0.4
40-44	7.3	6.6	1.1	0.9	0.6	0.5
45-49	4.8 ^b	6.9	0.6 ^a	1.3	0.3	0.8
50-54	3.6 ^a	5.7	0.8	0.8	0.4	0.2
55-59	1.8	2.6	0.5	0.4	0.3	0.2
60-64	1.4	0.6	0.2	*	0.2	*
65 or Older	0.3	0.3	0.0	0.1	0.0	0.0

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.15A Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	35,281	34,215	3,956	3,762	1,006	996
12	26	20	13	14	6	1
13	36	45	23	31	9	18
14	91	85	60	59	20	15
15	165	196	123	143	31	37
16	270	239	195	163	58	45
17	397 ^a	311	236	219	57	60
18	592	499	375 ^a	278	123	87
19	626	641	351	362	91	81
20	732	716	343	341	106	73
21	847	844	292	311	70	64
22	887	777	223	244	37	47
23	903	958	215	172	54	41
24	1,047	1,001	190	193	48	40
25	979	852	158	147	35	40
26-29	3,895	3,942	532	463	134	84
30-34	3,875	3,812	307	359	55	147
35-39	3,677	3,788	144	124	55	46
40-44	4,601 ^b	3,648	74	52	7	24
45-49	4,272	4,152	55	51	11	34
50-54	4,190	4,472	37	30	*	10
55-59	2,278	2,417	11	*	*	*
60-64	583	590	*	*	*	*
65 or Older	311	211	*	9	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.15B Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	14.3	13.8	1.6	1.5	0.4	0.4
12	0.7	0.5	0.3	0.3	0.2	0.0
13	0.9	1.1	0.6	0.8	0.2	0.4
14	2.1	2.0	1.4	1.4	0.5	0.4
15	3.6	4.5	2.7	3.2	0.7	0.8
16	6.2	5.4	4.5	3.7	1.3	1.0
17	9.5 ^a	7.5	5.6	5.3	1.4	1.5
18	12.9 ^a	10.5	8.2 ^b	5.8	2.7	1.8
19	15.0	14.6	8.4	8.3	2.2	1.9
20	18.2	17.5	8.5	8.3	2.6	1.8
21	19.7	20.7	6.8	7.6	1.6	1.6
22	22.5	20.5	5.7	6.4	0.9	1.2
23	22.9	24.2	5.5	4.3	1.4	1.0
24	26.7	25.4	4.9	4.9	1.2	1.0
25	25.5	22.9	4.1	3.9	0.9	1.1
26-29	23.6	24.0	3.2	2.8	0.8	0.5
30-34	20.8	20.2	1.6	1.9	0.3	0.8
35-39	17.6	18.0	0.7	0.6	0.3	0.2
40-44	20.6 ^b	17.2	0.3	0.2	0.0	0.1
45-49	19.4	18.4	0.2	0.2	0.1	0.1
50-54	20.1	20.6	0.2	0.1	*	0.0
55-59	13.0	13.1	0.1	*	*	*
60-64	4.3	4.4	*	*	*	*
65 or Older	0.9	0.6	*	0.0	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.16A Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	22,879	22,477	2,218	2,080	761	616
12	301	245	141	106	39	41
13	411	418	201	198	56	50
14	473	431	206	203	69	55
15	562	486	241 ^a	186	83	60
16	432	424	183	166	52	52
17	383	418	138	135	42	36
18	464 ^a	369	168 ^a	113	36	23
19	437	422	133	116	26	23
20	486	400	73	77	28	21
21	479	430	55	62	10	14
22	561	477	42	60	8	24
23	518	514	41	38	4	14
24	625	523	43	28	6	3
25	519	556	22	31	11	8
26-29	2,355	2,283	146	132	54	43
30-34	2,721	2,644	134	99	78	26
35-39	2,484	2,675	18 ^a	123	11	5
40-44	2,854	2,502	78	71	67	28
45-49	2,450	2,550	25	40	16	30
50-54	1,892	2,124	57	69	29	33
55-59	791	1,010	32	25	32	25
60-64	428	209	41	*	4	*
65 or Older	254	368	*	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.16B Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	9.3	9.1	0.9	0.8	0.3	0.2
12	7.7 ^a	6.1	3.6	2.6	1.0	1.0
13	9.9	10.1	4.9	4.8	1.4	1.2
14	11.1	10.4	4.8	4.9	1.6	1.3
15	12.4	11.1	5.3	4.2	1.8	1.4
16	10.0	9.5	4.2	3.7	1.2	1.2
17	9.1	10.1	3.3	3.3	1.0	0.9
18	10.1 ^a	7.7	3.7 ^a	2.4	0.8	0.5
19	10.5	9.6	3.2	2.6	0.6	0.5
20	12.1 ^a	9.8	1.8	1.9	0.7	0.5
21	11.1	10.5	1.3	1.5	0.2	0.3
22	14.2	12.6	1.1	1.6	0.2	0.6
23	13.1	13.0	1.0	1.0	0.1	0.4
24	15.9 ^a	13.3	1.1	0.7	0.2	0.1
25	13.5	14.9	0.6	0.8	0.3	0.2
26-29	14.3	13.9	0.9	0.8	0.3	0.3
30-34	14.6	14.0	0.7	0.5	0.4	0.1
35-39	11.9	12.7	0.1 ^a	0.6	0.1	0.0
40-44	12.8	11.8	0.3	0.3	0.3	0.1
45-49	11.1	11.3	0.1	0.2	0.1	0.1
50-54	9.1	9.8	0.3	0.3	0.1	0.2
55-59	4.5	5.5	0.2	0.1	0.2	0.1
60-64	3.2	1.6	0.3	*	0.0	*
65 or Older	0.7	1.0	*	*	*	*

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.17A Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	50,965	50,415	16,482	16,280	7,095	6,895
12	232 ^a	171	144 ^a	97	77	53
13	266	262	172	160	89	65
14	426	357	295	244	122	108
15	624	610	428	463	153	183
16	760	761	534	535	201	188
17	863	788	609	535	201	232
18	1,212	1,141	795	762	310	305
19	1,161	1,156	674	700	288	285
20	1,220	1,247	705	666	319	260
21	1,321	1,344	650	711	259	277
22	1,330	1,214	643	556	238	239
23	1,255	1,267	564	563	236	197
24	1,288	1,274	586	507	268 ^a	197
25	1,264	1,146	505	449	197	190
26-29	5,152	4,822	1,853	1,669	802	644
30-34	5,021	4,769	1,622	1,534	704	594
35-39	4,898	5,145	1,452	1,469	651	564
40-44	6,190 ^b	5,111	1,591	1,281	746	625
45-49	5,837	5,790	1,118	1,077	455	520
50-54	4,705	5,157	640	883	290	468
55-59	2,904 ^b	4,011	506	727	249	331
60-64	1,563	1,681	185	387	72	193
65 or Older	1,473	1,191	212	307	167	178

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.17B Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	20.7	20.3	6.7	6.6	2.9	2.8
12	5.9 ^a	4.3	3.7 ^a	2.4	2.0	1.3
13	6.4	6.4	4.1	3.9	2.2	1.6
14	10.0	8.6	6.9	5.9	2.9	2.6
15	13.7	13.9	9.4	10.5	3.4	4.2
16	17.5	17.1	12.3	12.0	4.7	4.2
17	20.6	19.1	14.5	12.9	4.8	5.6
18	26.4	23.9	17.3	16.0	6.8	6.4
19	27.8	26.4	16.1	16.0	6.9	6.5
20	30.3	30.5	17.5	16.3	7.9	6.4
21	30.8	32.9	15.1	17.4	6.0	6.8
22	33.7	32.0	16.3	14.7	6.0	6.3
23	31.8	32.0	14.3	14.2	6.0	5.0
24	32.8	32.3	14.9	12.9	6.8 ^a	5.0
25	33.0	30.8	13.2	12.1	5.1	5.1
26-29	31.2	29.4	11.2	10.2	4.9	3.9
30-34	27.0	25.2	8.7	8.1	3.8	3.1
35-39	23.4	24.4	6.9	7.0	3.1	2.7
40-44	27.7 ^a	24.1	7.1	6.0	3.3	2.9
45-49	26.5	25.7	5.1	4.8	2.1	2.3
50-54	22.5	23.7	3.1	4.1	1.4	2.2
55-59	16.6 ^b	21.8	2.9	3.9	1.4	1.8
60-64	11.6	12.6	1.4	2.9	0.5	1.4
65 or Older	4.1	3.3	0.6	0.8	0.5	0.5

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.18A Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	33,472	33,060	12,649	12,466	5,220	5,174
12	191 ^a	139	124	87	66	45
13	207	195	141	130	70	52
14	361	311	253	200	103	82
15	524	498	361	370	125	135
16	636	653	438	448	173	154
17	728	656	501	446	153	200
18	1,048	926	661	610	239	237
19	996	995	552	590	216	239
20	1,036	1,041	568	519	245	203
21	1,066	1,146	510	582	204	217
22	1,118	1,032	511	442	188	188
23	1,036	1,055	439	436	185	153
24	1,023	1,003	436	423	196	143
25	1,019	930	394	368	147	137
26-29	3,884	3,713	1,483	1,279	568	522
30-34	3,538	3,479	1,107	1,173	489	464
35-39	3,158	3,505	1,044	1,120	454	406
40-44	3,846 ^b	3,086	1,220	959	513	432
45-49	3,261	3,227	754	808	291	393
50-54	2,286	2,380	486	643	237	350
55-59	1,345 ^a	1,857	396	427	182	226
60-64	534	762	80	201	25	93
65 or Older	631	471	192	205	150	102

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.18B Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	13.6	13.3	5.1	5.0	2.1	2.1
12	4.9 ^a	3.4	3.2	2.2	1.7	1.1
13	5.0	4.7	3.4	3.1	1.7	1.3
14	8.4	7.5	5.9	4.8	2.4	2.0
15	11.5	11.4	7.9	8.4	2.8	3.1
16	14.7	14.7	10.1	10.1	4.0	3.5
17	17.4	15.9	12.0	10.8	3.7 ^a	4.8
18	22.8 ^a	19.4	14.4	12.8	5.2	5.0
19	23.8	22.7	13.2	13.5	5.2	5.5
20	25.7	25.5	14.1	12.7	6.1	5.0
21	24.8 ^a	28.0	11.9	14.2	4.7	5.3
22	28.3	27.2	12.9	11.6	4.8	5.0
23	26.3	26.7	11.1	11.0	4.7	3.9
24	26.1	25.5	11.1	10.7	5.0	3.6
25	26.6	25.0	10.3	9.9	3.8	3.7
26-29	23.5	22.6	9.0	7.8	3.4	3.2
30-34	19.0	18.4	5.9	6.2	2.6	2.5
35-39	15.1	16.6	5.0	5.3	2.2	1.9
40-44	17.2 ^a	14.5	5.5	4.5	2.3	2.0
45-49	14.8	14.3	3.4	3.6	1.3	1.7
50-54	10.9	11.0	2.3	3.0	1.1	1.6
55-59	7.7	10.1	2.3	2.3	1.0	1.2
60-64	4.0	5.7	0.6	1.5	0.2	0.7
65 or Older	1.8	1.3	0.5	0.6	0.4	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.