

Table 1.76A (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Numbers in Thousands, 2006 and 2007

| Geographic Characteristic | TIME PERIOD | | | | | |
|----------------------------|--------------------|--------|------------------|-------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| TOTAL | 22,856 | 22,558 | 2,219 | 2,085 | 762 | 617 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 4,440 | 4,322 | 447 | 390 | 173 | 139 |
| New England | 1,372 ^a | 1,081 | 106 | 92 | 36 | 45 |
| Middle Atlantic | 3,068 | 3,241 | 341 | 298 | 137 | 95 |
| Midwest | 5,110 | 5,007 | 455 | 427 | 161 | 99 |
| East North Central | 3,531 | 3,474 | 291 | 314 | 93 | 66 |
| West North Central | 1,579 | 1,533 | 164 | 112 | 68 | 34 |
| South | 7,287 | 7,242 | 739 | 647 | 239 | 168 |
| South Atlantic | 4,210 | 3,885 | 415 | 342 | 140 | 89 |
| East South Central | 1,027 | 1,138 | 122 ^b | 58 | 33 | 13 |
| West South Central | 2,050 | 2,220 | 202 | 248 | 65 | 67 |
| West | 6,019 | 5,987 | 579 | 621 | 190 | 210 |
| Mountain | 1,878 | 1,866 | 180 | 200 | 39 | 67 |
| Pacific | 4,141 | 4,121 | 399 | 421 | 151 | 143 |
| COUNTY TYPE | | | | | | |
| Large Metro | 13,291 | 12,550 | 1,287 | 1,197 | 494 | 362 |
| Small Metro | 6,449 | 6,751 | 607 | 627 | 173 | 184 |
| 250K – 1 Mil. Pop. | 4,401 | 4,493 | 468 | 392 | 144 | 101 |
| < 250K Pop. | 2,049 | 2,259 | 140 ^a | 235 | 29 | 83 |
| Nonmetro | 3,115 | 3,256 | 325 | 261 | 95 | 71 |
| Urbanized | 1,317 | 1,536 | 122 | 114 | 29 | 31 |
| Less Urbanized | 1,498 | 1,512 | 173 ^a | 121 | 57 | 33 |
| Completely Rural | 300 | 208 | 30 | 26 | 9 | 7 |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.76B (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

| Geographic Characteristic | TIME PERIOD | | | | | |
|----------------------------|-------------------|------|------------------|------|------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| TOTAL | 9.3 | 9.1 | 0.9 | 0.8 | 0.3 | 0.2 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 9.7 | 9.4 | 1.0 | 0.8 | 0.4 | 0.3 |
| New England | 11.4 ^a | 9.0 | 0.9 | 0.8 | 0.3 | 0.4 |
| Middle Atlantic | 9.1 | 9.6 | 1.0 | 0.9 | 0.4 | 0.3 |
| Midwest | 9.3 | 9.1 | 0.8 | 0.8 | 0.3 | 0.2 |
| East North Central | 9.2 | 9.1 | 0.8 | 0.8 | 0.2 | 0.2 |
| West North Central | 9.6 | 9.3 | 1.0 | 0.7 | 0.4 | 0.2 |
| South | 8.2 | 8.1 | 0.8 | 0.7 | 0.3 | 0.2 |
| South Atlantic | 8.9 | 8.2 | 0.9 | 0.7 | 0.3 | 0.2 |
| East South Central | 7.0 | 7.7 | 0.8 ^b | 0.4 | 0.2 | 0.1 |
| West South Central | 7.5 | 8.0 | 0.7 | 0.9 | 0.2 | 0.2 |
| West | 10.7 | 10.5 | 1.0 | 1.1 | 0.3 | 0.4 |
| Mountain | 11.1 | 10.8 | 1.1 | 1.2 | 0.2 | 0.4 |
| Pacific | 10.5 | 10.3 | 1.0 | 1.1 | 0.4 | 0.4 |
| COUNTY TYPE | | | | | | |
| Large Metro | 10.0 | 9.5 | 1.0 | 0.9 | 0.4 | 0.3 |
| Small Metro | 9.1 | 9.1 | 0.9 | 0.8 | 0.2 | 0.2 |
| 250K – 1 Mil. Pop. | 9.5 | 9.1 | 1.0 | 0.8 | 0.3 | 0.2 |
| < 250K Pop. | 8.5 | 9.2 | 0.6 ^a | 1.0 | 0.1 | 0.3 |
| Nonmetro | 7.4 | 7.9 | 0.8 | 0.6 | 0.2 | 0.2 |
| Urbanized | 8.0 | 9.1 | 0.7 | 0.7 | 0.2 | 0.2 |
| Less Urbanized | 7.1 | 7.7 | 0.8 | 0.6 | 0.3 | 0.2 |
| Completely Rural | 6.5 | 4.5 | 0.7 | 0.6 | 0.2 | 0.1 |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.77A (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Numbers in Thousands, 2006 and 2007

| Geographic Characteristic | TIME PERIOD | | | | | |
|----------------------------|------------------|-------|------------------|------|-----------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| TOTAL | 2,559 | 2,428 | 1,109 | 995 | 340 | 294 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 438 | 377 | 180 | 150 | 56 | 42 |
| New England | 111 | 113 | 46 | 44 | 13 | 17 |
| Middle Atlantic | 327 ^a | 264 | 134 | 106 | 43 | 25 |
| Midwest | 546 | 520 | 229 | 226 | 72 | 54 |
| East North Central | 386 | 371 | 158 | 164 | 44 | 38 |
| West North Central | 160 | 148 | 71 | 62 | 28 | 16 |
| South | 925 | 883 | 414 | 350 | 114 | 97 |
| South Atlantic | 484 | 453 | 211 | 182 | 55 | 52 |
| East South Central | 164 | 137 | 70 | 45 | 15 | 9 |
| West South Central | 277 | 293 | 133 | 123 | 44 | 36 |
| West | 650 | 648 | 286 | 268 | 98 | 101 |
| Mountain | 207 | 210 | 85 | 90 | 24 | 31 |
| Pacific | 443 | 438 | 201 | 178 | 74 | 70 |
| COUNTY TYPE | | | | | | |
| Large Metro | 1,369 | 1,290 | 621 ^a | 511 | 196 | 148 |
| Small Metro | 733 | 716 | 295 | 323 | 79 | 99 |
| 250K – 1 Mil. Pop. | 499 | 477 | 203 | 198 | 59 | 59 |
| < 250K Pop. | 234 | 239 | 92 | 125 | 20 | 40 |
| Nonmetro | 456 | 422 | 194 | 160 | 64 | 48 |
| Urbanized | 169 | 171 | 77 | 64 | 22 | 24 |
| Less Urbanized | 234 | 205 | 97 | 75 | 34 ^a | 17 |
| Completely Rural | 53 | 46 | 20 | 21 | 9 | 6 |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.77B (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2006 and 2007

| Geographic Characteristic | TIME PERIOD | | | | | |
|----------------------------|-------------|------|------------------|------|------------------|------|
| | Lifetime | | Past Year | | Past Month | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| TOTAL | 10.1 | 9.6 | 4.4 | 3.9 | 1.3 | 1.2 |
| GEOGRAPHIC DIVISION | | | | | | |
| Northeast | 9.7 | 8.5 | 4.0 | 3.4 | 1.2 | 0.9 |
| New England | 9.5 | 9.7 | 3.9 | 3.8 | 1.1 | 1.5 |
| Middle Atlantic | 9.8 | 8.0 | 4.0 | 3.2 | 1.3 | 0.8 |
| Midwest | 9.6 | 9.3 | 4.0 | 4.0 | 1.3 | 1.0 |
| East North Central | 9.7 | 9.4 | 4.0 | 4.2 | 1.1 | 1.0 |
| West North Central | 9.4 | 8.9 | 4.2 | 3.7 | 1.6 | 0.9 |
| South | 10.1 | 9.7 | 4.5 | 3.8 | 1.3 | 1.1 |
| South Atlantic | 10.4 | 9.8 | 4.5 | 3.9 | 1.2 | 1.1 |
| East South Central | 11.0 | 9.3 | 4.7 | 3.1 | 1.0 | 0.6 |
| West South Central | 9.2 | 9.7 | 4.4 | 4.1 | 1.5 | 1.2 |
| West | 10.7 | 10.7 | 4.7 | 4.4 | 1.6 | 1.7 |
| Mountain | 11.5 | 11.6 | 4.7 | 5.0 | 1.3 | 1.7 |
| Pacific | 10.4 | 10.4 | 4.7 | 4.2 | 1.7 | 1.7 |
| COUNTY TYPE | | | | | | |
| Large Metro | 10.0 | 9.4 | 4.5 ^a | 3.7 | 1.4 | 1.1 |
| Small Metro | 9.7 | 9.7 | 3.9 | 4.4 | 1.0 | 1.3 |
| 250K – 1 Mil. Pop. | 9.8 | 9.6 | 4.0 | 4.0 | 1.2 | 1.2 |
| < 250K Pop. | 9.4 | 10.0 | 3.7 | 5.2 | 0.8 | 1.7 |
| Nonmetro | 11.0 | 10.1 | 4.7 | 3.8 | 1.6 | 1.1 |
| Urbanized | 10.1 | 10.0 | 4.6 | 3.8 | 1.3 | 1.4 |
| Less Urbanized | 11.8 | 10.1 | 4.9 | 3.7 | 1.7 ^a | 0.9 |
| Completely Rural | 11.4 | 10.7 | 4.2 | 5.0 | 1.8 | 1.5 |

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.