

Table 1.78A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	50,983	50,541	16,482	16,334	7,098	6,907
GEOGRAPHIC DIVISION						
Northeast	8,607	8,038	2,656	2,583	1,172	987
New England	2,617 ^a	2,168	845 ^a	665	382	287
Middle Atlantic	5,990	5,871	1,812	1,919	790	700
Midwest	10,414	10,869	3,425	3,395	1,406	1,468
East North Central	7,200	7,637	2,463	2,499	1,013	1,051
West North Central	3,214	3,233	962	896	394	417
South	17,557	18,417	5,978	6,155	2,650	2,676
South Atlantic	8,974	9,350	2,892	2,880	1,177	1,174
East South Central	2,773 ^a	3,317	1,018 ^a	1,271	474	633
West South Central	5,809	5,751	2,068	2,005	998	870
West	14,406 ^a	13,216	4,423	4,200	1,869	1,777
Mountain	4,421	4,306	1,417	1,374	585	563
Pacific	9,985 ^a	8,910	3,006	2,826	1,284	1,214
COUNTY TYPE						
Large Metro	28,582	27,288	9,059	8,750	3,946	3,653
Small Metro	14,859	15,302	4,955	4,900	2,082	2,100
250K – 1 Mil. Pop.	10,100	10,290	3,348	3,190	1,359	1,297
< 250K Pop.	4,759	5,012	1,608	1,710	723	803
Nonmetro	7,542	7,951	2,468	2,684	1,070	1,154
Urbanized	3,201	3,522	1,044	1,187	428	440
Less Urbanized	3,616	3,723	1,125	1,346	505	635
Completely Rural	726	707	299 ^a	152	137	78

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.78B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	20.7	20.4	6.7	6.6	2.9	2.8
GEOGRAPHIC DIVISION						
Northeast	18.8	17.5	5.8	5.6	2.6	2.2
New England	21.8 ^a	18.0	7.0 ^a	5.5	3.2	2.4
Middle Atlantic	17.7	17.3	5.4	5.7	2.3	2.1
Midwest	19.0	19.8	6.3	6.2	2.6	2.7
East North Central	18.8	19.9	6.4	6.5	2.6	2.7
West North Central	19.6	19.6	5.9	5.4	2.4	2.5
South	19.7	20.5	6.7	6.8	3.0	3.0
South Atlantic	19.1	19.7	6.1	6.1	2.5	2.5
East South Central	19.0 ^a	22.5	7.0	8.6	3.2	4.3
West South Central	21.3	20.8	7.6	7.3	3.7	3.1
West	25.5 ^a	23.1	7.8	7.3	3.3	3.1
Mountain	26.2	24.9	8.4	8.0	3.5	3.3
Pacific	25.2 ^a	22.3	7.6	7.1	3.2	3.0
COUNTY TYPE						
Large Metro	21.4	20.6	6.8	6.6	3.0	2.8
Small Metro	21.0	20.7	7.0	6.6	2.9	2.8
250K – 1 Mil. Pop.	21.8	20.8	7.2	6.4	2.9	2.6
< 250K Pop.	19.6	20.4	6.6	7.0	3.0	3.3
Nonmetro	17.9	19.4	5.9	6.5	2.5	2.8
Urbanized	19.5	21.0	6.4	7.1	2.6	2.6
Less Urbanized	17.1	18.9	5.3 ^a	6.8	2.4 ^a	3.2
Completely Rural	15.8	15.4	6.5 ^a	3.3	3.0	1.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.79A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	3,175 ^a	2,954	2,184	2,037	846	830
GEOGRAPHIC DIVISION						
Northeast	471	406	319	268	114	90
New England	118	118	82	80	30	23
Middle Atlantic	352 ^a	287	237	188	84	68
Midwest	696	672	477	469	187	190
East North Central	496	479	344	339	133	131
West North Central	199	194	134	130	54	59
South	1,176	1,149	829	812	341	351
South Atlantic	534	531	376	354	159	135
East South Central	215	202	154	140	66	57
West South Central	427	417	299	318	116	158
West	831	726	559	487	204	198
Mountain	258	233	161	173	68	68
Pacific	573	494	399	314	136	130
COUNTY TYPE						
Large Metro	1,634	1,538	1,165	1,067	462	409
Small Metro	982	898	661	631	251	278
250K – 1 Mil. Pop.	638	584	444	399	162	166
< 250K Pop.	345	314	217	232	89	112
Nonmetro	558	518	358	339	132	143
Urbanized	218	219	142	142	54	57
Less Urbanized	277	248	178	160	66	64
Completely Rural	63	50	39	37	13	22

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.79B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	12.5	11.7	8.6	8.1	3.3	3.3
GEOGRAPHIC DIVISION						
Northeast	10.4	9.1	7.1	6.0	2.5	2.0
New England	10.2	10.2	7.0	6.9	2.6	2.0
Middle Atlantic	10.5	8.7	7.1	5.7	2.5	2.1
Midwest	12.3	12.0	8.4	8.4	3.3	3.4
East North Central	12.5	12.1	8.6	8.6	3.3	3.3
West North Central	11.8	11.6	7.9	7.8	3.2	3.5
South	12.9	12.6	9.1	8.9	3.7	3.8
South Atlantic	11.5	11.4	8.1	7.6	3.4	2.9
East South Central	14.5	13.6	10.3	9.5	4.5	3.9
West South Central	14.2	13.9	9.9	10.6	3.9	5.3
West	13.7	12.0	9.2	8.1	3.4	3.3
Mountain	14.4	12.9	9.0	9.6	3.8	3.8
Pacific	13.4	11.7	9.4	7.4	3.2	3.1
COUNTY TYPE						
Large Metro	11.9	11.2	8.5	7.8	3.4	3.0
Small Metro	13.0	12.2	8.7	8.5	3.3	3.8
250K – 1 Mil. Pop.	12.6	11.7	8.7	8.0	3.2	3.3
< 250K Pop.	13.9	13.1	8.8	9.7	3.6	4.7
Nonmetro	13.5	12.4	8.7	8.1	3.2	3.4
Urbanized	13.1	12.8	8.5	8.3	3.2	3.3
Less Urbanized	13.9	12.3	8.9	7.9	3.3	3.2
Completely Rural	13.5	11.8	8.2	8.6	2.8	5.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.80A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	10,067	9,807	5,130	4,920	2,122	1,947
GEOGRAPHIC DIVISION						
Northeast	1,664	1,586	850	807	329	294
New England	534	507	304	266	116	94
Middle Atlantic	1,129	1,079	546	541	213	200
Midwest	2,242	2,168	1,160	1,055	454	438
East North Central	1,542	1,546	816	791	325	319
West North Central	701	623	344 ^a	265	129	119
South	3,609	3,503	1,882	1,792	830	745
South Atlantic	1,732	1,733	954	912	428	377
East South Central	652	630	344	330	161	149
West South Central	1,225	1,140	584	550	242	220
West	2,552	2,550	1,237	1,266	509	470
Mountain	883	808	394	397	155	158
Pacific	1,670	1,742	843	870	354	312
COUNTY TYPE						
Large Metro	5,172	4,967	2,615	2,485	1,023	998
Small Metro	3,277	3,242	1,718	1,597	765 ^b	593
250K – 1 Mil. Pop.	2,117	2,166	1,127	1,084	524 ^b	389
< 250K Pop.	1,160	1,076	591	512	242	204
Nonmetro	1,619	1,598	796	838	334	356
Urbanized	682	735	330	381	126	160
Less Urbanized	793	757	391	394	172	168
Completely Rural	144	106	76	63	35	29

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.80B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	30.7	30.0	15.7	15.0	6.5	5.9
GEOGRAPHIC DIVISION						
Northeast	28.4	26.9	14.5	13.7	5.6	5.0
New England	35.0	33.2	20.0	17.4	7.6	6.2
Middle Atlantic	26.0	24.7	12.6	12.4	4.9	4.6
Midwest	30.4	29.8	15.7	14.5	6.1	6.0
East North Central	30.2	30.8	16.0	15.7	6.4	6.3
West North Central	30.7	27.5	15.1 ^a	11.7	5.6	5.3
South	30.9	30.0	16.1	15.3	7.1	6.4
South Atlantic	29.5	29.3	16.2	15.4	7.3	6.4
East South Central	34.4	33.7	18.2	17.6	8.5	8.0
West South Central	31.3	29.1	14.9	14.1	6.2	5.6
West	32.7	32.5	15.8	16.1	6.5	6.0
Mountain	37.5	34.3	16.7	16.8	6.6	6.7
Pacific	30.6	31.7	15.5	15.8	6.5	5.7
COUNTY TYPE						
Large Metro	29.5	28.7	14.9	14.4	5.8	5.8
Small Metro	32.4	31.4	17.0	15.5	7.6 ^b	5.7
250K – 1 Mil. Pop.	31.7	32.0	16.9	16.0	7.8 ^b	5.7
< 250K Pop.	33.6	30.3	17.1	14.4	7.0	5.7
Nonmetro	32.0	31.3	15.7	16.4	6.6	7.0
Urbanized	31.9	32.2	15.4	16.7	5.9	7.0
Less Urbanized	32.8	30.9	16.2	16.1	7.1	6.9
Completely Rural	28.4	28.4	14.9	17.0	6.9	7.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.81A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	37,741	37,781	9,168	9,377	4,130	4,130
GEOGRAPHIC DIVISION						
Northeast	6,472	6,046	1,487	1,508	729	602
New England	1,964 ^a	1,542	458	319	236	170
Middle Atlantic	4,508	4,505	1,029	1,189	493	433
Midwest	7,475	8,029	1,788	1,871	766	840
East North Central	5,162	5,612	1,304	1,370	555	602
West North Central	2,313	2,416	484	501	211	238
South	12,771	13,766	3,267	3,551	1,478	1,579
South Atlantic	6,708	7,086	1,562	1,614	591	661
East South Central	1,906 ^b	2,485	520 ^a	801	247 ^a	427
West South Central	4,158	4,194	1,185	1,137	640	492
West	11,023 ^a	9,940	2,627	2,447	1,156	1,109
Mountain	3,280	3,265	862	804	361	337
Pacific	7,743 ^a	6,675	1,764	1,642	795	771
COUNTY TYPE						
Large Metro	21,775	20,782	5,279	5,198	2,461	2,246
Small Metro	10,600	11,163	2,576	2,672	1,065	1,230
250K – 1 Mil. Pop.	7,345	7,540	1,776	1,706	673	743
< 250K Pop.	3,254	3,623	799	966	392	487
Nonmetro	5,366	5,836	1,314	1,507	604	655
Urbanized	2,301	2,568	571	664	248	224
Less Urbanized	2,547	2,717	557 ^a	791	267	403
Completely Rural	519	551	185 ^a	52	89	27

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.81B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	20.1	19.9	4.9	4.9	2.2	2.2
GEOGRAPHIC DIVISION						
Northeast	18.2	17.0	4.2	4.2	2.1	1.7
New England	21.0 ^a	16.5	4.9	3.4	2.5	1.8
Middle Atlantic	17.2	17.2	3.9	4.5	1.9	1.7
Midwest	17.9	19.2	4.3	4.5	1.8	2.0
East North Central	17.7	19.1	4.5	4.7	1.9	2.1
West North Central	18.6	19.2	3.9	4.0	1.7	1.9
South	18.7	19.9	4.8	5.1	2.2	2.3
South Atlantic	18.4	19.1	4.3	4.4	1.6	1.8
East South Central	17.0 ^b	21.8	4.6 ^a	7.0	2.2 ^a	3.7
West South Central	20.4	20.2	5.8	5.5	3.1	2.4
West	25.9 ^a	22.9	6.2	5.6	2.7	2.6
Mountain	25.8	24.9	6.8	6.1	2.8	2.6
Pacific	25.9 ^a	22.1	5.9	5.4	2.7	2.6
COUNTY TYPE						
Large Metro	21.3	20.4	5.2	5.1	2.4	2.2
Small Metro	20.0	19.8	4.9	4.7	2.0	2.2
250K – 1 Mil. Pop.	21.2	19.9	5.1	4.5	1.9	2.0
< 250K Pop.	17.8	19.5	4.4	5.2	2.1	2.6
Nonmetro	16.3 ^a	18.3	4.0	4.7	1.8	2.1
Urbanized	18.3	20.0	4.5	5.2	2.0	1.7
Less Urbanized	15.2 ^a	17.9	3.3 ^b	5.2	1.6 ^a	2.7
Completely Rural	14.3	14.5	5.1 ^a	1.4	2.5	0.7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.82A (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	47,809	47,588	14,298	14,297	6,252	6,077
GEOGRAPHIC DIVISION						
Northeast	8,136	7,633	2,338	2,315	1,058	897
New England	2,498 ^a	2,049	763 ^a	585	352	264
Middle Atlantic	5,638	5,583	1,575	1,731	706	633
Midwest	9,718	10,197	2,948	2,926	1,220	1,278
East North Central	6,704	7,158	2,120	2,160	880	920
West North Central	3,014	3,039	828	766	340	358
South	16,380	17,268	5,149	5,343	2,309	2,325
South Atlantic	8,440	8,819	2,516	2,526	1,019	1,038
East South Central	2,558 ^a	3,115	864 ^a	1,130	408	575
West South Central	5,382	5,334	1,769	1,687	882	711
West	13,575	12,490	3,864	3,713	1,665	1,578
Mountain	4,163	4,073	1,257	1,201	517	495
Pacific	9,412 ^a	8,416	2,607	2,512	1,149	1,083
COUNTY TYPE						
Large Metro	26,947	25,749	7,894	7,683	3,484	3,244
Small Metro	13,877	14,405	4,294	4,269	1,830	1,823
250K – 1 Mil. Pop.	9,462	9,706	2,904	2,791	1,197	1,132
< 250K Pop.	4,415	4,698	1,390	1,478	634	691
Nonmetro	6,985	7,434	2,110	2,345	938	1,011
Urbanized	2,983	3,303	902	1,045	374	383
Less Urbanized	3,339	3,474	948	1,185	440	571
Completely Rural	663	656	261 ^a	115	124	56

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.82B (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	21.7	21.4	6.5	6.4	2.8	2.7
GEOGRAPHIC DIVISION						
Northeast	19.7	18.4	5.7	5.6	2.6	2.2
New England	23.0 ^a	18.9	7.0 ^a	5.4	3.2	2.4
Middle Atlantic	18.5	18.3	5.2	5.7	2.3	2.1
Midwest	19.8	20.7	6.0	5.9	2.5	2.6
East North Central	19.6	20.8	6.2	6.3	2.6	2.7
West North Central	20.4	20.5	5.6	5.2	2.3	2.4
South	20.5	21.4	6.4	6.6	2.9	2.9
South Atlantic	19.9	20.5	5.9	5.9	2.4	2.4
East South Central	19.5 ^a	23.5	6.6 ^a	8.5	3.1	4.3
West South Central	22.1	21.7	7.3	6.8	3.6	2.9
West	26.9 ^a	24.4	7.7	7.3	3.3	3.1
Mountain	27.6	26.3	8.3	7.8	3.4	3.2
Pacific	26.6 ^a	23.6	7.4	7.0	3.2	3.0
COUNTY TYPE						
Large Metro	22.5	21.6	6.6	6.5	2.9	2.7
Small Metro	22.0	21.6	6.8	6.4	2.9	2.7
250K – 1 Mil. Pop.	22.9	21.8	7.0	6.3	2.9	2.5
< 250K Pop.	20.3	21.2	6.4	6.7	2.9	3.1
Nonmetro	18.4 ^a	20.1	5.5	6.4	2.5	2.7
Urbanized	20.2	21.9	6.1	6.9	2.5	2.5
Less Urbanized	17.4	19.7	4.9 ^b	6.7	2.3 ^a	3.2
Completely Rural	16.1	15.7	6.3 ^a	2.8	3.0	1.3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.