

Table 1.83A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	33,478	33,132	12,650	12,517	5,226	5,188
GEOGRAPHIC DIVISION						
Northeast	5,633	5,230	2,056	1,872	918	744
New England	1,740	1,487	692 ^a	535	314	209
Middle Atlantic	3,892	3,743	1,365	1,337	604	535
Midwest	7,024	7,474	2,771	2,685	1,066	1,123
East North Central	5,091	5,419	2,057	2,057	792	855
West North Central	1,932	2,055	714	628	274	267
South	11,562	11,815	4,532	4,709	1,972	2,001
South Atlantic	5,854	5,778	2,219	2,146	912	900
East South Central	1,989	2,287	822	1,045	368	524
West South Central	3,719	3,751	1,491	1,519	691	577
West	9,260	8,614	3,290	3,251	1,270	1,321
Mountain	2,966	2,788	1,054	1,082	452	449
Pacific	6,294	5,826	2,236	2,169	818	872
COUNTY TYPE						
Large Metro	18,475	17,808	6,881	6,540	2,907	2,659
Small Metro	10,176	10,080	3,809	3,847	1,505	1,597
250K – 1 Mil. Pop.	6,929	6,611	2,621	2,425	1,018	960
< 250K Pop.	3,246	3,469	1,188	1,422	488	637
Nonmetro	4,828	5,244	1,960	2,130	814	932
Urbanized	2,028	2,334	840	934	320	358
Less Urbanized	2,342	2,467	893	1,065	381	511
Completely Rural	457	443	227	130	113	63

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.83B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	13.6	13.4	5.1	5.1	2.1	2.1
GEOGRAPHIC DIVISION						
Northeast	12.3	11.4	4.5	4.1	2.0	1.6
New England	14.5	12.4	5.8 ^a	4.4	2.6	1.7
Middle Atlantic	11.5	11.1	4.0	3.9	1.8	1.6
Midwest	12.8	13.6	5.1	4.9	1.9	2.0
East North Central	13.3	14.1	5.4	5.4	2.1	2.2
West North Central	11.8	12.5	4.3	3.8	1.7	1.6
South	13.0	13.1	5.1	5.2	2.2	2.2
South Atlantic	12.4	12.1	4.7	4.5	1.9	1.9
East South Central	13.6	15.5	5.6	7.1	2.5	3.6
West South Central	13.6	13.6	5.5	5.5	2.5	2.1
West	16.4	15.1	5.8	5.7	2.2	2.3
Mountain	17.6	16.1	6.2	6.3	2.7	2.6
Pacific	15.9	14.6	5.6	5.4	2.1	2.2
COUNTY TYPE						
Large Metro	13.9	13.4	5.2	4.9	2.2	2.0
Small Metro	14.4	13.6	5.4	5.2	2.1	2.2
250K – 1 Mil. Pop.	14.9 ^a	13.3	5.6 ^a	4.9	2.2	1.9
< 250K Pop.	13.4	14.1	4.9	5.8	2.0	2.6
Nonmetro	11.5 ^a	12.8	4.6	5.2	1.9	2.3
Urbanized	12.4	13.9	5.1	5.6	2.0	2.1
Less Urbanized	11.1	12.5	4.2 ^a	5.4	1.8 ^a	2.6
Completely Rural	9.9	9.6	4.9	2.8	2.4	1.4

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.84A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	2,650 ^a	2,457	1,821	1,684	692	671
GEOGRAPHIC DIVISION						
Northeast	381	335	264	229	90	75
New England	89	90	60	61	19	12
Middle Atlantic	292	245	203	169	71	62
Midwest	582	570	392	388	151	152
East North Central	420	407	286	285	107	105
West North Central	162	162	106	103	44	46
South	980	939	697	674	282	295
South Atlantic	437	432	316	303	134	124
East South Central	190	167	138	114	49	48
West South Central	353	340	242	256	99	122
West	707	613	468	393	168	150
Mountain	227	187	145	139	59	53
Pacific	480	426	324	254	110	97
COUNTY TYPE						
Large Metro	1,357	1,271	974	884	385	328
Small Metro	807	752	532	525	193	227
250K – 1 Mil. Pop.	509	477	349	328	119	133
< 250K Pop.	298	275	182	197	75	94
Nonmetro	486	434	315	276	114	116
Urbanized	196	180	126	110	48	46
Less Urbanized	234	206	152	131	53	49
Completely Rural	57	48	37	34	12	21

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.84B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	10.4	9.7	7.2	6.7	2.7	2.7
GEOGRAPHIC DIVISION						
Northeast	8.4	7.5	5.8	5.1	2.0	1.7
New England	7.6	7.8	5.2	5.2	1.7	1.1
Middle Atlantic	8.7	7.4	6.1	5.1	2.1	1.9
Midwest	10.3	10.1	6.9	6.9	2.7	2.7
East North Central	10.5	10.3	7.2	7.2	2.7	2.7
West North Central	9.6	9.7	6.3	6.2	2.6	2.8
South	10.7	10.3	7.6	7.4	3.1	3.2
South Atlantic	9.4	9.3	6.8	6.5	2.9	2.7
East South Central	12.8	11.3	9.3	7.7	3.3	3.3
West South Central	11.7	11.3	8.0	8.5	3.3	4.1
West	11.7	10.2	7.7	6.5	2.8	2.5
Mountain	12.7	10.4	8.1	7.7	3.3	2.9
Pacific	11.3	10.1	7.6	6.0	2.6	2.3
COUNTY TYPE						
Large Metro	9.9	9.3	7.1	6.5	2.8	2.4
Small Metro	10.7	10.2	7.0	7.1	2.6	3.1
250K – 1 Mil. Pop.	10.0	9.6	6.9	6.6	2.3	2.7
< 250K Pop.	12.0	11.5	7.3	8.2	3.0	3.9
Nonmetro	11.8	10.4	7.6	6.6	2.8	2.8
Urbanized	11.7	10.5	7.6	6.4	2.9	2.7
Less Urbanized	11.8	10.2	7.6	6.5	2.7	2.4
Completely Rural	12.2	11.2	7.8	8.1	2.6	4.8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.85A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	8,353	8,143	4,077	3,981	1,627	1,518
GEOGRAPHIC DIVISION						
Northeast	1,377	1,307	665	640	246	216
New England	460	406	254	214	87	64
Middle Atlantic	918	901	411	426	159	152
Midwest	1,891	1,841	975	871	365	333
East North Central	1,333	1,348	701	664	267	246
West North Central	558	493	275 ^a	207	98	87
South	2,953	2,848	1,467	1,425	634	590
South Atlantic	1,452	1,396	759	720	320	310
East South Central	543	527	278	270	137	122
West South Central	958	924	430	434	178	158
West	2,131	2,147	969	1,046	381	379
Mountain	731	679	313	312	129	120
Pacific	1,400	1,469	656	734	252	259
COUNTY TYPE						
Large Metro	4,259	4,125	2,016	2,004	744	765
Small Metro	2,747	2,697	1,406	1,291	606 ^b	467
250K – 1 Mil. Pop.	1,782	1,824	925	906	432 ^b	308
< 250K Pop.	964	874	481 ^a	385	174	159
Nonmetro	1,347	1,320	655	686	276	286
Urbanized	562	591	267	320	106	129
Less Urbanized	660	637	323	314	143	134
Completely Rural	125	93	66	52	28	23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.85B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	25.5	24.9	12.5	12.2	5.0	4.6
GEOGRAPHIC DIVISION						
Northeast	23.5	22.1	11.3	10.8	4.2	3.7
New England	30.1	26.6	16.6	14.0	5.7	4.2
Middle Atlantic	21.2	20.6	9.5	9.7	3.7	3.5
Midwest	25.6	25.3	13.2	12.0	4.9	4.6
East North Central	26.1	26.8	13.7	13.2	5.2	4.9
West North Central	24.5	21.8	12.0 ^a	9.2	4.3	3.8
South	25.3	24.4	12.6	12.2	5.4	5.1
South Atlantic	24.7	23.6	12.9	12.2	5.4	5.3
East South Central	28.7	28.2	14.7	14.4	7.2	6.5
West South Central	24.5	23.6	11.0	11.1	4.5	4.0
West	27.3	27.3	12.4	13.3	4.9	4.8
Mountain	31.0	28.8	13.3	13.2	5.5	5.1
Pacific	25.7	26.7	12.0	13.3	4.6	4.7
COUNTY TYPE						
Large Metro	24.3	23.8	11.5	11.6	4.2	4.4
Small Metro	27.1	26.1	13.9	12.5	6.0 ^b	4.5
250K – 1 Mil. Pop.	26.7	26.9	13.9	13.4	6.5 ^b	4.6
< 250K Pop.	27.9	24.6	13.9 ^a	10.8	5.1	4.5
Nonmetro	26.6	25.9	12.9	13.4	5.5	5.6
Urbanized	26.3	25.9	12.5	14.0	4.9	5.6
Less Urbanized	27.3	26.0	13.4	12.8	5.9	5.5
Completely Rural	24.6	24.9	12.9	14.0	5.5	6.3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.86A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	22,475	22,533	6,752	6,851	2,908	2,998
GEOGRAPHIC DIVISION						
Northeast	3,875	3,588	1,128	1,003	582	453
New England	1,192	991	378	261	207	132
Middle Atlantic	2,683	2,597	750	742	375	321
Midwest	4,551	5,063	1,403	1,425	550	638
East North Central	3,339	3,664	1,070	1,107	418	504
West North Central	1,212	1,399	333	318	132	134
South	7,628	8,029	2,368	2,611	1,055	1,115
South Atlantic	3,965	3,950	1,144	1,122	458	466
East South Central	1,255 ^a	1,593	406 ^a	661	183 ^a	353
West South Central	2,408	2,486	818	828	414	296
West	6,422	5,853	1,853	1,813	721	792
Mountain	2,008	1,922	596	631	265	276
Pacific	4,414	3,931	1,257	1,181	456	516
COUNTY TYPE						
Large Metro	12,859	12,412	3,890	3,652	1,778	1,566
Small Metro	6,622	6,631	1,872	2,032	706	902
250K – 1 Mil. Pop.	4,638	4,311	1,347	1,192	467	519
< 250K Pop.	1,984	2,320	525 ^a	840	239	383
Nonmetro	2,995	3,490	990	1,168	424	531
Urbanized	1,270	1,563	447	504	166	184
Less Urbanized	1,449	1,624	418 ^a	621	185 ^a	328
Completely Rural	275	303	124	43	73	19

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.86B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	12.0	11.9	3.6	3.6	1.5	1.6
GEOGRAPHIC DIVISION						
Northeast	10.9	10.1	3.2	2.8	1.6	1.3
New England	12.8	10.6	4.0	2.8	2.2	1.4
Middle Atlantic	10.3	9.9	2.9	2.8	1.4	1.2
Midwest	10.9	12.1	3.4	3.4	1.3	1.5
East North Central	11.4	12.5	3.7	3.8	1.4	1.7
West North Central	9.7	11.1	2.7	2.5	1.1	1.1
South	11.2	11.6	3.5	3.8	1.5	1.6
South Atlantic	10.9	10.7	3.1	3.0	1.3	1.3
East South Central	11.2	14.0	3.6 ^a	5.8	1.6 ^a	3.1
West South Central	11.8	12.0	4.0	4.0	2.0	1.4
West	15.1	13.5	4.3	4.2	1.7	1.8
Mountain	15.8	14.7	4.7	4.8	2.1	2.1
Pacific	14.8	13.0	4.2	3.9	1.5	1.7
COUNTY TYPE						
Large Metro	12.6	12.2	3.8	3.6	1.7	1.5
Small Metro	12.5	11.8	3.5	3.6	1.3	1.6
250K – 1 Mil. Pop.	13.4 ^a	11.4	3.9	3.2	1.3	1.4
< 250K Pop.	10.8	12.5	2.9 ^a	4.5	1.3	2.1
Nonmetro	9.1 ^a	11.0	3.0	3.7	1.3	1.7
Urbanized	10.1	12.2	3.6	3.9	1.3	1.4
Less Urbanized	8.7 ^a	10.7	2.5 ^b	4.1	1.1 ^a	2.2
Completely Rural	7.6	8.0	3.4	1.1	2.0	0.5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.87A (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	30,828	30,676	10,829	10,833	4,534	4,517
GEOGRAPHIC DIVISION						
Northeast	5,252	4,895	1,793	1,642	828	669
New England	1,651	1,397	631 ^a	474	295	197
Middle Atlantic	3,600	3,498	1,162	1,168	534	473
Midwest	6,442	6,904	2,379	2,297	915	971
East North Central	4,672	5,012	1,771	1,772	685	750
West North Central	1,770	1,892	608	525	230	221
South	10,581	10,876	3,835	4,036	1,690	1,706
South Atlantic	5,417	5,346	1,903	1,843	778	776
East South Central	1,798	2,120	684 ^a	931	320	475
West South Central	3,366	3,411	1,249	1,262	592	454
West	8,553	8,000	2,822	2,858	1,102	1,171
Mountain	2,739	2,600	909	943	394	396
Pacific	5,814	5,400	1,913	1,915	708	775
COUNTY TYPE						
Large Metro	17,118	16,537	5,906	5,656	2,522	2,331
Small Metro	9,369	9,328	3,278	3,323	1,312	1,369
250K – 1 Mil. Pop.	6,421	6,134	2,272	2,098	899	827
< 250K Pop.	2,948	3,194	1,006	1,225	413	542
Nonmetro	4,341	4,810	1,645	1,854	701	817
Urbanized	1,833	2,154	714	824	272	312
Less Urbanized	2,109	2,261	741	934	328	462
Completely Rural	400	395	190	95	100	42

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.87B (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	14.0	13.8	4.9	4.9	2.1	2.0
GEOGRAPHIC DIVISION						
Northeast	12.7	11.8	4.3	4.0	2.0	1.6
New England	15.2	12.9	5.8 ^a	4.4	2.7	1.8
Middle Atlantic	11.8	11.5	3.8	3.8	1.8	1.5
Midwest	13.1	14.0	4.9	4.7	1.9	2.0
East North Central	13.6	14.6	5.2	5.2	2.0	2.2
West North Central	12.0	12.8	4.1	3.5	1.6	1.5
South	13.3	13.5	4.8	5.0	2.1	2.1
South Atlantic	12.8	12.5	4.5	4.3	1.8	1.8
East South Central	13.7	16.0	5.2 ^a	7.0	2.4	3.6
West South Central	13.8	13.9	5.1	5.1	2.4	1.8
West	17.0	15.6	5.6	5.6	2.2	2.3
Mountain	18.2	16.8	6.0	6.1	2.6	2.6
Pacific	16.5	15.1	5.4	5.4	2.0	2.2
COUNTY TYPE						
Large Metro	14.3	13.9	4.9	4.8	2.1	2.0
Small Metro	14.9	14.0	5.2	5.0	2.1	2.1
250K – 1 Mil. Pop.	15.5 ^a	13.8	5.5 ^a	4.7	2.2	1.9
< 250K Pop.	13.6	14.4	4.6	5.5	1.9	2.4
Nonmetro	11.4 ^a	13.0	4.3	5.0	1.8	2.2
Urbanized	12.4	14.3	4.8	5.5	1.8	2.1
Less Urbanized	11.0	12.8	3.9 ^a	5.3	1.7 ^a	2.6
Completely Rural	9.7	9.5	4.6	2.3	2.4	1.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.