

**Table 1.75C (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
**Standard Errors of Numbers in Thousands, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
<b>TOTAL</b>	644	640	149	140	70	75
<b>GEOGRAPHIC DIVISION</b>						
Northeast	268	243	64	55	25	37
New England	147	103	36	19	12	11
Middle Atlantic	228	219	52	51	22	35
Midwest	226	235	58	46	26	21
East North Central	173	201	48	39	22	17
West North Central	147	126	31	22	14	11
South	351	384	85	90	38	45
South Atlantic	283	290	63	54	24	21
East South Central	115	138	36	34	15	18
West South Central	170	198	46	61	25	35
West	404	386	89	80	46	44
Mountain	167	180	35	46	28	28
Pacific	365	342	81	65	36	33
<b>COUNTY TYPE</b>						
Large Metro	590	581	124	107	53	58
Small Metro	325	397	73	85	42	44
250K – 1 Mil. Pop.	270	350	63	73	40	37
< 250K Pop.	191	218	39	41	12	23
Nonmetro	229	219	44	39	17	17
Urbanized	145	157	33	25	13	12
Less Urbanized	162	156	27	29	12	12
Completely Rural	99	49	12	6	1	2

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 1.75D (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:  
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
<b>TOTAL</b>	0.26	0.26	0.06	0.06	0.03	0.03
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.59	0.53	0.14	0.12	0.06	0.08
New England	1.22	0.86	0.30	0.16	0.10	0.09
Middle Atlantic	0.67	0.65	0.15	0.15	0.07	0.10
Midwest	0.41	0.43	0.11	0.08	0.05	0.04
East North Central	0.45	0.52	0.13	0.10	0.06	0.04
West North Central	0.90	0.76	0.19	0.14	0.08	0.07
South	0.39	0.43	0.10	0.10	0.04	0.05
South Atlantic	0.60	0.61	0.13	0.11	0.05	0.04
East South Central	0.79	0.93	0.25	0.23	0.11	0.12
West South Central	0.62	0.72	0.17	0.22	0.09	0.13
West	0.72	0.68	0.16	0.14	0.08	0.08
Mountain	0.99	1.04	0.21	0.26	0.16	0.16
Pacific	0.92	0.86	0.20	0.16	0.09	0.08
<b>COUNTY TYPE</b>						
Large Metro	0.40	0.38	0.09	0.08	0.04	0.04
Small Metro	0.43	0.45	0.10	0.11	0.06	0.06
250K – 1 Mil. Pop.	0.54	0.57	0.14	0.14	0.09	0.07
< 250K Pop.	0.71	0.78	0.16	0.17	0.05	0.10
Nonmetro	0.47	0.48	0.10	0.10	0.04	0.04
Urbanized	0.74	0.74	0.19	0.15	0.08	0.07
Less Urbanized	0.65	0.72	0.12	0.15	0.06	0.06
Completely Rural	1.52	0.94	0.27	0.13	0.01	0.04

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 1.75P (REVISED) Hallucinogen Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:**  
***P* Values from Tests of Differences of Percentages, 2007 Versus 2006**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
<b>TOTAL</b>	0.1598		0.2386		0.8674	
<b>GEOGRAPHIC DIVISION</b>						
Northeast	0.1361		0.0926		0.7978	
New England	0.0579		0.0612		0.6384	
Middle Atlantic	0.5533		0.3952		0.6355	
Midwest	0.8609		0.0850		0.2113	
East North Central	0.4819		0.2050		0.1369	
West North Central	0.2032		0.2262		0.9810	
South	0.5485		0.9883		0.6624	
South Atlantic	0.2523		0.1888		0.9040	
East South Central	0.0558		0.8967		0.8297	
West South Central	0.4181		0.1895		0.5510	
West	0.4889		0.8750		0.8256	
Mountain	0.5586		0.3513		0.9048	
Pacific	0.6246		0.7396		0.7082	
<b>COUNTY TYPE</b>						
Large Metro	0.0890		0.1687		0.9364	
Small Metro	0.9291		0.9462		0.9367	
250K – 1 Mil. Pop.	0.9865		0.9840		0.4224	
< 250K Pop.	0.9091		0.8652		0.1320	
Nonmetro	0.6835		0.8446		0.7270	
Urbanized	0.9602		0.6298		0.8351	
Less Urbanized	0.7127		0.5972		0.6252	
Completely Rural	0.0097		0.2444		0.2030	

\*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.