

Table 1.11C Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	897	971	566	596	426	445
12	30	27	23	20	14	15
13	33	38	26	30	19	18
14	40	42	34	35	22	23
15	55	49	43	43	32	32
16	55	57	47	49	34	37
17	58	59	51	51	36	38
18	83	84	72	72	50	52
19	86	86	70	72	56	53
20	91	90	74	74	56	54
21	94	92	69	75	50	57
22	87	81	62	60	47	46
23	84	81	59	60	41	42
24	83	86	59	59	42	45
25	77	77	55	49	40	40
26-29	312	309	203	185	156	135
30-34	305	319	172	176	127	136
35-39	352	368	180	183	135	129
40-44	392	376	193	170	157	125
45-49	399	407	186	169	129	128
50-54	476	550	192	241	160	184
55-59	386	502	117	179	84	132
60-64	270	342	95	130	67	80
65 or Older	271	296	99	87	71	71

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.11D Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.36	0.39	0.23	0.24	0.17	0.18
12	0.71	0.63	0.58	0.48	0.35	0.36
13	0.75	0.82	0.61	0.68	0.44	0.44
14	0.84	0.90	0.74	0.78	0.50	0.53
15	0.97	0.92	0.85	0.85	0.66	0.69
16	0.98	1.01	0.91	0.92	0.73	0.76
17	1.07	1.08	1.00	1.00	0.79	0.82
18	1.17	1.20	1.13	1.13	0.94	0.95
19	1.22	1.30	1.17	1.24	1.07	1.04
20	1.35	1.29	1.33	1.30	1.16	1.13
21	1.22	1.26	1.17	1.28	1.01	1.11
22	1.29	1.30	1.24	1.25	1.01	1.06
23	1.21	1.23	1.14	1.20	0.89	0.97
24	1.29	1.27	1.21	1.21	0.93	1.00
25	1.31	1.25	1.21	1.15	0.97	0.96
26-29	1.14	1.21	1.02	0.97	0.85	0.75
30-34	1.13	1.18	0.86	0.87	0.65	0.69
35-39	1.17	1.17	0.81	0.81	0.62	0.58
40-44	1.15	1.17	0.80	0.72	0.66	0.56
45-49	1.09	1.08	0.79	0.70	0.57	0.54
50-54	1.55	1.67	0.87	1.06	0.75	0.83
55-59	1.71	1.92	0.65	0.93	0.47	0.70
60-64	1.69	1.96	0.70	0.95	0.50	0.59
65 or Older	0.76	0.81	0.28	0.24	0.20	0.19

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.11P Illicit Drug Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.2061		0.6550		0.2782
12	0.0216		0.0479		0.3450
13	0.9164		0.7948		0.2695
14	0.1528		0.3111		0.7339
15	0.0357		0.2248		0.9322
16	0.8236		0.8490		0.9019
17	0.2879		0.2846		0.8165
18	0.0671		0.0948		0.9973
19	0.3821		0.7361		0.9644
20	0.1655		0.4016		0.3203
21	0.4083		0.3417		0.0525
22	0.1486		0.3995		0.7685
23	0.6209		0.6513		0.6239
24	0.8472		0.1127		0.2155
25	0.8522		0.0810		0.8989
26-29	0.1111		0.2324		0.2388
30-34	0.8765		0.6867		0.5306
35-39	0.9562		0.7926		0.4274
40-44	0.1582		0.6116		0.1188
45-49	0.7163		0.9651		0.5717
50-54	0.0588		0.2731		0.7960
55-59	0.0014		0.0077		0.0613
60-64	0.0086		0.3779		0.7458
65 or Older	0.3964		0.8943		0.9632

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.12C Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	908	963	477	492	356	360
12	11	9	10	9	4	7
13	18	18	15	15	10	9
14	27	29	24	25	17	17
15	42	39	37	34	27	25
16	49	48	43	42	30	31
17	52	54	45	45	32	35
18	77	75	67	65	46	49
19	78	78	63	65	51	50
20	87	86	68	69	51	49
21	90	84	64	66	45	50
22	80	74	56	55	43	42
23	81	78	52	53	38	39
24	76	83	50	55	35	40
25	74	72	48	45	35	36
26-29	287	290	167	160	129	116
30-34	284	290	145	145	107	114
35-39	339	340	148	144	114	110
40-44	375	352	151	136	128	102
45-49	383	391	144	138	109	109
50-54	462	532	162	198	132	154
55-59	373	487	90	129	63	90
60-64	233	318	68	91	55	39
65 or Older	231	261	75	47	36	41

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.12D Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.37	0.39	0.19	0.20	0.14	0.15
12	0.28	0.23	0.27	0.21	0.10	0.17
13	0.42	0.43	0.36	0.35	0.24	0.21
14	0.61	0.65	0.55	0.58	0.38	0.40
15	0.82	0.78	0.74	0.72	0.57	0.55
16	0.93	0.94	0.87	0.83	0.66	0.66
17	0.99	1.05	0.93	0.93	0.71	0.77
18	1.17	1.17	1.11	1.10	0.88	0.92
19	1.24	1.28	1.14	1.15	1.00	1.01
20	1.42	1.31	1.29	1.24	1.09	1.05
21	1.30	1.31	1.14	1.22	0.93	1.04
22	1.32	1.32	1.16	1.21	0.95	0.99
23	1.25	1.28	1.05	1.12	0.85	0.91
24	1.30	1.35	1.09	1.17	0.83	0.92
25	1.33	1.30	1.10	1.07	0.86	0.90
26-29	1.17	1.20	0.90	0.87	0.73	0.66
30-34	1.14	1.16	0.75	0.74	0.56	0.58
35-39	1.21	1.16	0.68	0.66	0.53	0.51
40-44	1.18	1.13	0.65	0.61	0.56	0.47
45-49	1.11	1.10	0.64	0.58	0.50	0.47
50-54	1.56	1.68	0.75	0.90	0.62	0.70
55-59	1.69	1.92	0.51	0.68	0.36	0.48
60-64	1.53	1.89	0.50	0.67	0.41	0.29
65 or Older	0.65	0.72	0.21	0.13	0.10	0.11

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.12P Marijuana Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.1262		0.4708		0.3387
12	0.6974		1.0000		0.1830
13	0.3811		0.4890		0.6122
14	0.3608		0.3830		0.9816
15	0.0152		0.1172		0.9506
16	0.2481		0.5729		0.5689
17	0.6278		0.5298		0.5036
18	0.0354		0.0716		0.9900
19	0.6885		0.9756		0.8569
20	0.2391		0.7193		0.5475
21	0.4136		0.5080		0.0861
22	0.1303		0.8284		0.6455
23	0.6672		0.8408		0.4592
24	0.6913		0.7967		0.4018
25	0.5714		0.4982		0.6144
26-29	0.5495		0.8828		0.6900
30-34	0.8930		0.5900		0.4166
35-39	0.7255		0.6110		0.6720
40-44	0.2041		0.4356		0.2569
45-49	0.9168		0.6832		0.9156
50-54	0.0717		0.7441		0.7018
55-59	0.0112		0.0456		0.1514
60-64	0.0074		0.7660		0.0864
65 or Older	0.0578		0.4409		0.9354

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.13C Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	654	659	230	242	153	149
12	2	3	1	2	1	2
13	4	5	3	5	2	3
14	8	6	7	6	5	3
15	14	13	12	13	6	4
16	16	19	14	14	6	7
17	22	21	19	18	8	9
18	36	37	30	30	18	11
19	38	36	30	28	15	14
20	42	40	29	29	17	15
21	44	42	29	28	14	17
22	42	42	23	26	10	11
23	42	47	23	27	12	11
24	46	42	28	22	14	12
25	44	44	25	25	14	14
26-29	185	166	94	83	68	46
30-34	175	169	77	76	49	50
35-39	216	212	85	70	55	42
40-44	271	245	81	59	62	42
45-49	259	263	57	84	40	66
50-54	291	373	70	111	57	54
55-59	197	255	56	60	45	38
60-64	128	163	29	2	20	*
65 or Older	109	110	8	39	7	34

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.13D Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.27	0.09	0.10	0.06	0.06
12	0.06	0.08	0.04	0.05	0.03	0.04
13	0.10	0.12	0.07	0.12	0.04	0.08
14	0.18	0.16	0.17	0.15	0.13	0.07
15	0.30	0.30	0.27	0.28	0.12	0.10
16	0.36	0.41	0.33	0.31	0.14	0.17
17	0.51	0.49	0.43	0.41	0.20	0.21
18	0.73	0.74	0.63	0.60	0.38	0.22
19	0.83	0.78	0.67	0.61	0.35	0.32
20	0.99	0.89	0.69	0.67	0.42	0.36
21	0.93	0.94	0.66	0.66	0.33	0.40
22	1.00	0.97	0.57	0.66	0.25	0.29
23	0.96	1.06	0.57	0.65	0.30	0.28
24	1.03	0.96	0.67	0.56	0.36	0.30
25	1.00	1.07	0.64	0.66	0.37	0.38
26-29	0.99	0.90	0.55	0.49	0.41	0.28
30-34	0.86	0.83	0.41	0.40	0.26	0.27
35-39	0.97	0.89	0.40	0.33	0.26	0.20
40-44	1.02	0.98	0.36	0.28	0.28	0.20
45-49	0.99	0.95	0.26	0.37	0.18	0.29
50-54	1.24	1.49	0.34	0.51	0.27	0.25
55-59	1.05	1.26	0.32	0.33	0.26	0.20
60-64	0.92	1.18	0.21	0.01	0.15	*
65 or Older	0.31	0.30	0.02	0.11	0.02	0.09

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.13P Cocaine Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.7302		0.2511		0.0799
12	0.6644		0.8483		0.8162
13	0.9208		0.3592		0.5761
14	0.6501		0.7204		0.1127
15	0.4643		0.5846		0.3738
16	0.6263		0.6453		0.9954
17	0.4810		0.7354		0.4989
18	0.0308		0.0083		0.0027
19	0.3401		0.0914		0.1735
20	0.6684		0.7646		0.2850
21	0.4323		0.7820		0.5928
22	0.8424		0.0197		0.3395
23	0.1178		0.4538		0.6988
24	0.5379		0.0379		0.1990
25	0.8676		0.3865		0.4907
26-29	0.2863		0.2971		0.3998
30-34	0.1366		0.1174		0.5369
35-39	0.9261		0.1708		0.1524
40-44	0.0574		0.2010		0.1554
45-49	0.3886		0.0142		0.0382
50-54	0.0163		0.2506		0.7201
55-59	0.1933		0.7912		0.8239
60-64	0.6553		0.0635		*
65 or Older	0.6783		0.1218		0.1989

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.14C Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	329	339	135	123	101	80
12	*	*	*	*	*	*
13	0	2	*	2	*	0
14	5	4	5	3	2	2
15	7	6	6	6	1	2
16	7	7	5	6	2	3
17	10	8	7	7	2	3
18	19	10	14	6	5	4
19	16	15	7	8	3	5
20	19	15	9	10	4	3
21	18	20	7	6	4	1
22	18	19	5	9	2	4
23	19	18	9	8	7	4
24	21	18	8	10	5	3
25	20	18	9	8	2	5
26-29	100	68	50	29	46	21
30-34	88	93	30	27	21	15
35-39	124	103	45	40	34	29
40-44	134	126	55	39	43	29
45-49	108	135	42	60	29	43
50-54	124	174	64	66	47	29
55-59	82	111	46	37	34	27
60-64	67	43	16	*	16	*
65 or Older	68	62	7	23	7	13

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.14D Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.13	0.14	0.05	0.05	0.04	0.03
12	*	*	*	*	*	*
13	0.00	0.05	*	0.04	*	0.01
14	0.11	0.10	0.11	0.08	0.06	0.05
15	0.16	0.14	0.13	0.14	0.01	0.05
16	0.17	0.17	0.12	0.13	0.05	0.07
17	0.25	0.19	0.17	0.16	0.06	0.07
18	0.40	0.21	0.31	0.12	0.11	0.08
19	0.38	0.33	0.18	0.17	0.07	0.11
20	0.46	0.36	0.23	0.25	0.10	0.08
21	0.41	0.48	0.17	0.15	0.10	0.03
22	0.46	0.49	0.14	0.24	0.04	0.10
23	0.47	0.46	0.24	0.20	0.17	0.10
24	0.51	0.45	0.19	0.26	0.12	0.07
25	0.51	0.48	0.23	0.22	0.05	0.14
26-29	0.58	0.41	0.30	0.18	0.28	0.13
30-34	0.46	0.48	0.16	0.14	0.11	0.08
35-39	0.57	0.48	0.21	0.19	0.16	0.14
40-44	0.58	0.57	0.25	0.18	0.19	0.14
45-49	0.49	0.58	0.19	0.26	0.13	0.19
50-54	0.58	0.79	0.31	0.30	0.23	0.14
55-59	0.46	0.59	0.26	0.20	0.19	0.15
60-64	0.49	0.32	0.12	*	0.12	*
65 or Older	0.19	0.17	0.02	0.06	0.02	0.04

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.14P Crack Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.9378		0.8263		0.4485
12	*		*		*
13	0.0519		*		*
14	0.9080		0.8340		0.7806
15	0.5365		0.7483		0.3164
16	0.4959		0.9330		0.6255
17	0.3286		0.2512		0.9187
18	0.0003		0.0007		0.4967
19	0.5179		0.7603		0.6525
20	0.1286		0.5819		0.7481
21	0.9636		0.6387		0.1597
22	0.8265		0.0494		0.1152
23	0.4148		0.4631		0.2567
24	0.1194		0.6856		0.1156
25	0.8300		0.4606		0.3631
26-29	0.0396		0.5750		0.5133
30-34	0.5273		0.3970		0.7069
35-39	0.1123		0.7903		0.4865
40-44	0.3782		0.4834		0.5510
45-49	0.0057		0.0293		0.0570
50-54	0.0245		0.9860		0.4312
55-59	0.3119		0.8370		0.7302
60-64	0.1387		*		*
65 or Older	0.9397		0.3065		0.6989

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.15C Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	644	638	149	140	70	74
12	7	5	5	4	3	1
13	7	8	6	6	3	5
14	13	11	11	9	6	5
15	17	22	15	19	7	8
16	22	22	18	18	10	9
17	28	23	20	19	10	10
18	38	41	29	29	19	14
19	42	42	31	31	16	13
20	48	46	34	31	23	14
21	51	51	29	30	14	13
22	50	43	24	24	9	9
23	48	54	21	20	12	9
24	54	53	21	23	11	10
25	51	47	21	23	10	11
26-29	194	188	73	64	40	28
30-34	188	207	53	61	18	43
35-39	206	193	51	37	24	25
40-44	231	196	31	18	6	12
45-49	225	207	22	19	8	17
50-54	288	330	22	18	*	10
55-59	213	233	10	*	*	*
60-64	96	123	*	*	*	*
65 or Older	101	71	*	9	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.15D Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.26	0.26	0.06	0.06	0.03	0.03
12	0.17	0.13	0.13	0.11	0.09	0.03
13	0.16	0.20	0.14	0.16	0.08	0.12
14	0.31	0.27	0.25	0.22	0.13	0.12
15	0.36	0.50	0.32	0.43	0.16	0.19
16	0.49	0.48	0.40	0.39	0.22	0.19
17	0.64	0.52	0.47	0.46	0.24	0.25
18	0.77	0.81	0.61	0.58	0.42	0.30
19	0.85	0.89	0.66	0.67	0.37	0.30
20	1.08	0.99	0.81	0.70	0.55	0.34
21	1.09	1.05	0.66	0.70	0.33	0.32
22	1.11	1.00	0.59	0.61	0.22	0.24
23	1.05	1.17	0.52	0.48	0.30	0.23
24	1.16	1.16	0.51	0.56	0.27	0.25
25	1.18	1.10	0.54	0.60	0.26	0.29
26-29	1.04	0.96	0.44	0.38	0.24	0.17
30-34	0.94	0.97	0.28	0.32	0.10	0.23
35-39	0.92	0.81	0.25	0.18	0.11	0.12
40-44	0.92	0.82	0.14	0.08	0.03	0.06
45-49	0.92	0.82	0.10	0.09	0.04	0.08
50-54	1.25	1.35	0.11	0.08	*	0.05
55-59	1.13	1.17	0.06	*	*	*
60-64	0.70	0.90	*	*	*	*
65 or Older	0.28	0.20	*	0.02	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.15P Hallucinogen Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.1428		0.2720		0.8641
12	0.4050		0.9828		0.1810
13	0.3670		0.3691		0.1713
14	0.8620		0.9425		0.5701
15	0.1925		0.3231		0.5631
16	0.2072		0.1387		0.2595
17	0.0146		0.5985		0.7905
18	0.0287		0.0054		0.1004
19	0.7595		0.8916		0.5028
20	0.6549		0.8675		0.1932
21	0.5464		0.4118		0.8851
22	0.1981		0.3519		0.3416
23	0.3916		0.1197		0.3893
24	0.4463		0.9650		0.5702
25	0.1029		0.8333		0.6372
26-29	0.7624		0.4918		0.3102
30-34	0.6422		0.5580		0.0522
35-39	0.7495		0.7394		0.7785
40-44	0.0067		0.5982		0.1739
45-49	0.4186		0.8602		0.2419
50-54	0.7837		0.7751		*
55-59	0.9441		*		*
60-64	0.9292		*		*
65 or Older	0.3994		*		*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.16C Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	471	503	107	106	73	67
12	23	22	17	15	7	11
13	27	30	19	21	9	11
14	26	28	18	19	11	10
15	32	30	21	19	13	10
16	30	29	19	17	11	10
17	25	27	16	16	10	8
18	34	31	20	17	8	7
19	34	34	20	18	7	7
20	35	33	13	14	8	7
21	38	33	11	13	5	7
22	37	35	9	12	4	9
23	35	37	10	9	2	6
24	40	35	10	8	3	2
25	36	39	7	11	7	5
26-29	148	143	44	36	28	23
30-34	152	161	36	26	28	12
35-39	149	164	9	43	8	5
40-44	177	164	36	26	35	17
45-49	161	155	16	18	14	16
50-54	205	232	35	42	21	31
55-59	130	167	26	25	26	25
60-64	107	70	22	*	4	*
65 or Older	69	92	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.16D Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.19	0.20	0.04	0.04	0.03	0.03
12	0.58	0.52	0.43	0.37	0.19	0.27
13	0.62	0.69	0.45	0.49	0.22	0.26
14	0.59	0.64	0.42	0.46	0.26	0.25
15	0.65	0.65	0.44	0.42	0.28	0.22
16	0.65	0.63	0.42	0.39	0.25	0.22
17	0.58	0.63	0.37	0.37	0.24	0.20
18	0.69	0.62	0.43	0.37	0.19	0.14
19	0.75	0.72	0.46	0.41	0.18	0.17
20	0.81	0.76	0.33	0.34	0.21	0.18
21	0.82	0.76	0.27	0.32	0.11	0.18
22	0.85	0.85	0.22	0.32	0.10	0.23
23	0.81	0.86	0.25	0.22	0.06	0.15
24	0.93	0.83	0.26	0.19	0.07	0.04
25	0.88	0.95	0.18	0.28	0.19	0.13
26-29	0.81	0.79	0.26	0.21	0.17	0.14
30-34	0.76	0.78	0.19	0.14	0.15	0.06
35-39	0.68	0.74	0.05	0.20	0.04	0.02
40-44	0.73	0.72	0.16	0.12	0.16	0.08
45-49	0.69	0.65	0.07	0.08	0.06	0.07
50-54	0.93	1.02	0.17	0.19	0.10	0.14
55-59	0.71	0.87	0.15	0.14	0.15	0.14
60-64	0.77	0.52	0.16	*	0.03	*
65 or Older	0.19	0.25	*	*	*	*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.16P Inhalant Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.4154		0.3174		0.1264
12	0.0416		0.0930		0.9131
13	0.8088		0.9407		0.6754
14	0.4719		0.8999		0.4645
15	0.1702		0.0755		0.1876
16	0.6345		0.3892		0.9580
17	0.2822		0.9517		0.6844
18	0.0116		0.0254		0.1963
19	0.3917		0.3466		0.6974
20	0.0433		0.8857		0.5188
21	0.5752		0.5602		0.6529
22	0.1909		0.2126		0.0833
23	0.9079		0.8483		0.1330
24	0.0313		0.2494		0.3592
25	0.2763		0.4317		0.7550
26-29	0.7541		0.8215		0.7740
30-34	0.5644		0.4220		0.0905
35-39	0.4263		0.0167		0.5031
40-44	0.3649		0.9515		0.3499
45-49	0.8525		0.5554		0.5302
50-54	0.6019		0.8537		0.9328
55-59	0.3895		0.8151		0.8151
60-64	0.0639		*		*
65 or Older	0.3420		*		*

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.17C Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	700	696	360	380	243	251
12	22	18	18	14	12	10
13	22	24	18	18	13	11
14	29	26	24	22	15	15
15	35	37	26	33	16	20
16	37	39	31	34	19	19
17	41	39	35	31	19	20
18	58	58	47	49	27	28
19	57	57	42	43	27	27
20	59	64	42	46	28	29
21	63	67	44	47	26	29
22	59	60	41	37	24	25
23	61	59	40	39	23	22
24	59	57	41	35	28	21
25	58	51	36	32	23	20
26-29	237	214	135	116	91	70
30-34	208	209	114	120	74	69
35-39	223	238	129	132	84	81
40-44	278	239	143	113	94	82
45-49	262	248	111	95	68	66
50-54	300	342	105	147	73	111
55-59	236	295	96	136	72	96
60-64	179	194	71	99	37	75
65 or Older	175	162	66	78	61	64

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.17D Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.28	0.28	0.15	0.15	0.10	0.10
12	0.54	0.44	0.45	0.33	0.31	0.25
13	0.52	0.56	0.43	0.43	0.30	0.26
14	0.65	0.61	0.54	0.51	0.34	0.35
15	0.71	0.77	0.56	0.69	0.34	0.43
16	0.78	0.80	0.65	0.70	0.42	0.41
17	0.89	0.82	0.77	0.67	0.44	0.47
18	1.03	0.99	0.91	0.90	0.57	0.56
19	1.14	1.13	0.92	0.91	0.61	0.60
20	1.20	1.22	0.94	0.97	0.67	0.66
21	1.16	1.25	0.92	1.00	0.59	0.68
22	1.25	1.28	0.96	0.88	0.60	0.62
23	1.20	1.23	0.89	0.90	0.56	0.54
24	1.21	1.16	0.94	0.82	0.67	0.51
25	1.22	1.11	0.88	0.80	0.60	0.52
26-29	1.17	1.02	0.77	0.67	0.54	0.42
30-34	0.95	0.95	0.59	0.61	0.39	0.36
35-39	0.96	0.96	0.60	0.60	0.40	0.38
40-44	1.04	0.95	0.61	0.51	0.41	0.38
45-49	0.98	0.91	0.48	0.41	0.31	0.29
50-54	1.27	1.38	0.50	0.66	0.35	0.50
55-59	1.22	1.45	0.54	0.72	0.41	0.52
60-64	1.24	1.34	0.52	0.73	0.27	0.55
65 or Older	0.49	0.45	0.18	0.22	0.17	0.18

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.17P Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.3493		0.5185		0.4643
12	0.0141		0.0231		0.0964
13	0.9470		0.6788		0.1414
14	0.1521		0.1899		0.6047
15	0.8892		0.2115		0.1518
16	0.6888		0.7517		0.4794
17	0.1916		0.1202		0.2132
18	0.0920		0.3169		0.6478
19	0.3699		0.8905		0.6539
20	0.8880		0.3755		0.0899
21	0.2147		0.1006		0.3993
22	0.3804		0.2099		0.7305
23	0.8882		0.9524		0.1933
24	0.7736		0.1046		0.0336
25	0.1836		0.3585		0.9446
26-29	0.2299		0.2828		0.1677
30-34	0.2035		0.4868		0.2341
35-39	0.4621		0.9741		0.4380
40-44	0.0130		0.1812		0.4816
45-49	0.5187		0.6385		0.5668
50-54	0.5294		0.2108		0.2007
55-59	0.0056		0.2637		0.6057
60-64	0.5856		0.0630		0.1407
65 or Older	0.2010		0.3781		0.9250

*Low precision; no estimate reported.

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.18C Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	539	562	306	316	216	215
12	20	16	18	13	12	10
13	19	20	16	16	11	9
14	27	24	22	19	13	12
15	32	34	24	29	14	17
16	34	37	27	31	18	17
17	37	34	31	27	16	19
18	54	51	42	42	23	23
19	53	53	37	40	22	24
20	53	60	37	41	24	27
21	55	60	40	40	23	24
22	55	56	37	34	23	23
23	55	54	33	33	21	19
24	51	51	32	32	22	17
25	52	48	32	29	20	17
26-29	200	189	121	99	79	65
30-34	173	184	93	104	60	62
35-39	178	198	101	120	74	70
40-44	226	189	126	99	83	67
45-49	199	180	92	84	55	57
50-54	211	218	93	129	69	100
55-59	159	195	84	98	62	76
60-64	99	123	36	72	16	56
65 or Older	105	88	64	61	60	44

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.18D Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.22	0.23	0.12	0.13	0.09	0.09
12	0.50	0.40	0.44	0.32	0.30	0.23
13	0.46	0.48	0.39	0.38	0.26	0.22
14	0.60	0.56	0.51	0.46	0.31	0.28
15	0.66	0.71	0.52	0.63	0.31	0.37
16	0.70	0.76	0.58	0.65	0.39	0.38
17	0.80	0.75	0.69	0.62	0.38	0.44
18	0.99	0.93	0.83	0.81	0.48	0.48
19	1.05	1.07	0.83	0.86	0.51	0.54
20	1.10	1.17	0.84	0.86	0.59	0.61
21	1.10	1.21	0.87	0.91	0.53	0.57
22	1.18	1.25	0.85	0.81	0.56	0.57
23	1.15	1.16	0.77	0.78	0.51	0.48
24	1.11	1.12	0.77	0.77	0.55	0.43
25	1.16	1.07	0.78	0.72	0.52	0.45
26-29	1.03	0.95	0.70	0.57	0.47	0.38
30-34	0.86	0.85	0.49	0.53	0.32	0.33
35-39	0.81	0.85	0.48	0.56	0.35	0.33
40-44	0.91	0.80	0.55	0.45	0.36	0.31
45-49	0.83	0.71	0.41	0.37	0.25	0.25
50-54	0.96	0.96	0.44	0.58	0.33	0.46
55-59	0.88	1.03	0.47	0.52	0.35	0.41
60-64	0.71	0.90	0.27	0.54	0.12	0.42
65 or Older	0.30	0.24	0.18	0.17	0.17	0.12

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.18P Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.4043		0.5138		0.7810
12	0.0208		0.0617		0.1278
13	0.6978		0.6230		0.2325
14	0.2833		0.1196		0.3115
15	0.8581		0.5571		0.4983
16	0.9909		0.9627		0.3447
17	0.1653		0.2168		0.0462
18	0.0137		0.1767		0.7130
19	0.4471		0.8355		0.6960
20	0.8810		0.2419		0.1808
21	0.0481		0.0562		0.4511
22	0.5446		0.2763		0.7991
23	0.8003		0.9273		0.2477
24	0.6853		0.7349		0.0500
25	0.3211		0.7212		0.8137
26-29	0.5154		0.1784		0.6723
30-34	0.6281		0.7226		0.7055
35-39	0.1857		0.6560		0.6151
40-44	0.0286		0.1922		0.5894
45-49	0.6448		0.7741		0.2374
50-54	0.9960		0.3782		0.3847
55-59	0.0777		0.9311		0.7252
60-64	0.1313		0.1266		0.2410
65 or Older	0.2191		0.9116		0.5030

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.