

Table 1.76C (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	471	503	107	106	73	67
GEOGRAPHIC DIVISION						
Northeast	191	205	56	59	40	44
New England	96	94	21	17	13	15
Middle Atlantic	166	183	52	56	38	42
Midwest	181	184	43	32	33	14
East North Central	139	145	27	28	17	11
West North Central	121	115	33	16	28	9
South	292	284	54	52	39	27
South Atlantic	236	235	44	36	34	19
East South Central	83	90	20	10	13	4
West South Central	134	135	25	37	14	17
West	276	305	57	61	31	40
Mountain	132	124	30	39	8	27
Pacific	242	279	48	48	30	29
COUNTY TYPE						
Large Metro	436	434	90	92	63	56
Small Metro	244	265	54	53	36	34
250K – 1 Mil. Pop.	204	220	51	38	35	19
< 250K Pop.	146	161	18	38	6	29
Nonmetro	154	175	27	21	13	11
Urbanized	102	124	16	14	7	6
Less Urbanized	109	122	19	16	9	9
Completely Rural	55	35	9	5	4	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 1.76D (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.19	0.20	0.04	0.04	0.03	0.03
GEOGRAPHIC DIVISION						
Northeast	0.42	0.45	0.12	0.13	0.09	0.10
New England	0.80	0.78	0.17	0.14	0.11	0.12
Middle Atlantic	0.49	0.54	0.15	0.17	0.11	0.12
Midwest	0.33	0.34	0.08	0.06	0.06	0.03
East North Central	0.36	0.38	0.07	0.07	0.05	0.03
West North Central	0.74	0.70	0.20	0.09	0.17	0.06
South	0.33	0.32	0.06	0.06	0.04	0.03
South Atlantic	0.50	0.49	0.09	0.07	0.07	0.04
East South Central	0.57	0.61	0.14	0.07	0.09	0.03
West South Central	0.49	0.49	0.09	0.13	0.05	0.06
West	0.49	0.53	0.10	0.11	0.06	0.07
Mountain	0.78	0.72	0.18	0.22	0.05	0.16
Pacific	0.61	0.70	0.12	0.12	0.08	0.07
COUNTY TYPE						
Large Metro	0.30	0.31	0.07	0.07	0.05	0.04
Small Metro	0.33	0.32	0.08	0.07	0.05	0.05
250K – 1 Mil. Pop.	0.42	0.39	0.11	0.07	0.07	0.04
< 250K Pop.	0.53	0.60	0.07	0.15	0.03	0.12
Nonmetro	0.33	0.40	0.06	0.05	0.03	0.03
Urbanized	0.56	0.65	0.09	0.09	0.04	0.04
Less Urbanized	0.45	0.56	0.09	0.08	0.04	0.04
Completely Rural	1.01	0.76	0.19	0.11	0.09	0.06

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.76P (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
***P* Values from Tests of Differences of Percentages, 2007 Versus 2006**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.5050		0.3301		0.1278	
GEOGRAPHIC DIVISION						
Northeast	0.6679		0.4907		0.5784	
New England	0.0330		0.6593		0.6654	
Middle Atlantic	0.4894		0.5713		0.4511	
Midwest	0.6663		0.5942		0.0827	
East North Central	0.7576		0.5354		0.1627	
West North Central	0.7683		0.1604		0.2408	
South	0.7670		0.1804		0.1288	
South Atlantic	0.2772		0.1686		0.1816	
East South Central	0.3893		0.0042		0.1353	
West South Central	0.4401		0.3083		0.9601	
West	0.7796		0.6999		0.7221	
Mountain	0.7559		0.7489		0.3485	
Pacific	0.8732		0.8054		0.8281	
COUNTY TYPE						
Large Metro	0.2287		0.5209		0.1134	
Small Metro	0.9641		0.8921		0.9558	
250K – 1 Mil. Pop.	0.4693		0.0779		0.2086	
< 250K Pop.	0.3597		0.0284		0.0695	
Nonmetro	0.3249		0.0934		0.1969	
Urbanized	0.2104		0.6111		0.9169	
Less Urbanized	0.4015		0.0688		0.1034	
Completely Rural	0.1228		0.7089		0.7079	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.77C (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	66	68	45	43	26	26
GEOGRAPHIC DIVISION						
Northeast	25	24	17	15	10	8
New England	11	13	10	7	4	5
Middle Atlantic	22	21	14	13	9	7
Midwest	24	28	17	19	10	9
East North Central	20	24	13	16	7	8
West North Central	13	14	11	9	6	4
South	41	42	27	27	14	16
South Atlantic	31	31	20	20	10	11
East South Central	16	16	11	8	5	3
West South Central	22	22	15	16	9	11
West	41	38	26	23	16	16
Mountain	19	15	12	10	6	6
Pacific	36	35	23	21	15	14
COUNTY TYPE						
Large Metro	55	60	37	34	21	20
Small Metro	39	39	23	25	12	15
250K – 1 Mil. Pop.	31	29	19	17	11	10
< 250K Pop.	23	26	13	19	5	12
Nonmetro	27	25	17	15	10	8
Urbanized	15	14	10	9	6	6
Less Urbanized	21	20	12	11	7	4
Completely Rural	9	7	5	5	4	3

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.77D (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.26	0.27	0.18	0.17	0.10	0.10
GEOGRAPHIC DIVISION						
Northeast	0.55	0.55	0.37	0.33	0.22	0.18
New England	0.96	1.14	0.82	0.61	0.37	0.39
Middle Atlantic	0.67	0.62	0.41	0.40	0.27	0.20
Midwest	0.42	0.49	0.31	0.33	0.18	0.16
East North Central	0.50	0.61	0.34	0.42	0.19	0.21
West North Central	0.80	0.85	0.65	0.54	0.38	0.26
South	0.45	0.46	0.30	0.29	0.16	0.18
South Atlantic	0.66	0.67	0.42	0.43	0.21	0.24
East South Central	1.11	1.05	0.71	0.52	0.32	0.21
West South Central	0.73	0.74	0.49	0.55	0.30	0.37
West	0.67	0.64	0.43	0.39	0.26	0.26
Mountain	1.08	0.83	0.67	0.58	0.34	0.33
Pacific	0.84	0.83	0.54	0.50	0.35	0.34
COUNTY TYPE						
Large Metro	0.37	0.39	0.26	0.23	0.15	0.14
Small Metro	0.48	0.48	0.30	0.32	0.15	0.20
250K – 1 Mil. Pop.	0.57	0.54	0.36	0.33	0.21	0.20
< 250K Pop.	0.83	0.93	0.52	0.69	0.20	0.46
Nonmetro	0.56	0.57	0.39	0.36	0.25	0.18
Urbanized	0.81	0.76	0.59	0.50	0.36	0.34
Less Urbanized	0.89	0.91	0.55	0.55	0.35	0.20
Completely Rural	1.73	1.66	1.10	1.08	0.84	0.64

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.77P (REVISED) Inhalant Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.2196		0.0829		0.2328	
GEOGRAPHIC DIVISION						
Northeast	0.0994		0.2119		0.3123	
New England	0.8859		0.9060		0.4839	
Middle Atlantic	0.0515		0.1686		0.1297	
Midwest	0.5749		1.0000		0.1943	
East North Central	0.7217		0.7149		0.6018	
West North Central	0.6267		0.5908		0.1269	
South	0.4940		0.1064		0.4344	
South Atlantic	0.4830		0.3040		0.8084	
East South Central	0.2736		0.0758		0.3229	
West South Central	0.5915		0.6756		0.5900	
West	1.0000		0.6232		0.8703	
Mountain	0.9529		0.7715		0.4718	
Pacific	0.9712		0.4857		0.9004	
COUNTY TYPE						
Large Metro	0.2940		0.0193		0.0818	
Small Metro	1.0000		0.2730		0.2635	
250K – 1 Mil. Pop.	0.7273		0.9556		0.9587	
< 250K Pop.	0.6800		0.0856		0.1016	
Nonmetro	0.2563		0.1180		0.1842	
Urbanized	0.9015		0.2714		0.8495	
Less Urbanized	0.2095		0.1422		0.0353	
Completely Rural	0.7748		0.6150		0.7828	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.