

Table 1.78C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	701	698	360	381	244	251
GEOGRAPHIC DIVISION						
Northeast	276	277	147	152	102	84
New England	143	144	76	62	57	37
Middle Atlantic	236	237	125	139	84	76
Midwest	269	285	139	143	89	97
East North Central	213	234	116	124	76	83
West North Central	154	163	71	69	43	48
South	406	439	221	251	147	161
South Atlantic	307	317	163	177	104	111
East South Central	148	168	85	112	52	80
West South Central	215	237	121	131	89	83
West	403	400	204	204	136	153
Mountain	162	179	103	100	75	71
Pacific	369	359	174	177	112	136
COUNTY TYPE						
Large Metro	685	657	309	310	211	209
Small Metro	392	481	187	216	110	138
250K – 1 Mil. Pop.	321	415	155	171	88	105
< 250K Pop.	244	274	113	134	68	92
Nonmetro	266	308	133	154	81	89
Urbanized	168	225	85	109	44	55
Less Urbanized	207	213	90	114	53	73
Completely Rural	95	110	59	23	44	15

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.78D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.28	0.28	0.15	0.15	0.10	0.10
GEOGRAPHIC DIVISION						
Northeast	0.60	0.60	0.32	0.33	0.22	0.18
New England	1.19	1.20	0.64	0.52	0.48	0.31
Middle Atlantic	0.70	0.70	0.37	0.41	0.25	0.22
Midwest	0.49	0.52	0.25	0.26	0.16	0.18
East North Central	0.56	0.61	0.30	0.32	0.20	0.22
West North Central	0.93	0.99	0.43	0.42	0.26	0.29
South	0.46	0.49	0.25	0.28	0.16	0.18
South Atlantic	0.65	0.67	0.35	0.37	0.22	0.23
East South Central	1.01	1.14	0.58	0.76	0.36	0.54
West South Central	0.79	0.86	0.44	0.48	0.33	0.30
West	0.71	0.70	0.36	0.36	0.24	0.27
Mountain	0.96	1.04	0.61	0.58	0.44	0.41
Pacific	0.93	0.90	0.44	0.44	0.28	0.34
COUNTY TYPE						
Large Metro	0.42	0.42	0.22	0.22	0.15	0.15
Small Metro	0.48	0.50	0.26	0.27	0.15	0.17
250K – 1 Mil. Pop.	0.59	0.62	0.32	0.31	0.18	0.19
< 250K Pop.	0.86	0.88	0.44	0.52	0.28	0.37
Nonmetro	0.51	0.60	0.30	0.35	0.19	0.21
Urbanized	0.80	0.97	0.47	0.57	0.26	0.30
Less Urbanized	0.74	0.85	0.37	0.50	0.25	0.35
Completely Rural	1.53	1.73	1.14	0.45	0.94	0.31

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.78P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.4086		0.5906		0.4809	
GEOGRAPHIC DIVISION						
Northeast	0.1396		0.7194		0.1466	
New England	0.0347		0.0382		0.1395	
Middle Atlantic	0.7052		0.5727		0.4138	
Midwest	0.2758		0.8536		0.6519	
East North Central	0.1774		0.8364		0.7336	
West North Central	0.9791		0.4793		0.7416	
South	0.2591		0.7172		0.9897	
South Atlantic	0.5354		0.8375		0.9102	
East South Central	0.0176		0.0531		0.0767	
West South Central	0.7081		0.6293		0.2443	
West	0.0148		0.3346		0.5948	
Mountain	0.3882		0.5932		0.7576	
Pacific	0.0211		0.4132		0.6611	
COUNTY TYPE						
Large Metro	0.1459		0.5148		0.3451	
Small Metro	0.5606		0.2491		0.6171	
250K – 1 Mil. Pop.	0.2282		0.0625		0.2405	
< 250K Pop.	0.5074		0.6090		0.5228	
Nonmetro	0.0642		0.1419		0.3368	
Urbanized	0.2548		0.3395		0.9762	
Less Urbanized	0.0961		0.0125		0.0435	
Completely Rural	0.8586		0.0103		0.2038	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.79C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	75	74	62	63	40	40
GEOGRAPHIC DIVISION						
Northeast	28	23	23	21	14	12
New England	13	13	10	11	7	5
Middle Atlantic	25	20	20	17	12	11
Midwest	28	31	24	25	15	15
East North Central	23	24	19	21	12	12
West North Central	16	18	14	13	10	9
South	44	47	37	40	24	26
South Atlantic	28	35	25	29	16	18
East South Central	17	17	15	15	10	9
West South Central	28	27	24	24	15	18
West	44	42	37	34	23	22
Mountain	22	18	14	16	11	11
Pacific	38	37	34	30	20	20
COUNTY TYPE						
Large Metro	61	64	51	53	32	32
Small Metro	45	44	36	38	22	23
250K – 1 Mil. Pop.	36	36	30	30	17	17
< 250K Pop.	28	26	22	22	13	15
Nonmetro	29	29	22	23	14	14
Urbanized	19	20	14	15	9	9
Less Urbanized	22	21	16	16	9	9
Completely Rural	9	8	7	7	4	5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.79D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.30	0.29	0.24	0.25	0.16	0.16
GEOGRAPHIC DIVISION						
Northeast	0.62	0.52	0.50	0.47	0.31	0.26
New England	1.12	1.09	0.90	0.99	0.58	0.43
Middle Atlantic	0.73	0.59	0.60	0.53	0.36	0.32
Midwest	0.49	0.55	0.42	0.44	0.27	0.28
East North Central	0.57	0.62	0.48	0.53	0.30	0.31
West North Central	0.92	1.08	0.81	0.75	0.57	0.57
South	0.48	0.51	0.41	0.44	0.27	0.29
South Atlantic	0.61	0.75	0.53	0.63	0.35	0.38
East South Central	1.17	1.17	1.00	1.00	0.70	0.62
West South Central	0.92	0.90	0.79	0.81	0.49	0.60
West	0.73	0.69	0.62	0.56	0.38	0.37
Mountain	1.21	1.00	0.79	0.90	0.61	0.58
Pacific	0.88	0.89	0.80	0.70	0.47	0.46
COUNTY TYPE						
Large Metro	0.42	0.42	0.35	0.35	0.23	0.22
Small Metro	0.52	0.51	0.44	0.44	0.28	0.29
250K – 1 Mil. Pop.	0.63	0.61	0.55	0.52	0.34	0.32
< 250K Pop.	0.94	0.93	0.76	0.81	0.50	0.61
Nonmetro	0.62	0.62	0.47	0.51	0.32	0.33
Urbanized	0.98	0.90	0.75	0.75	0.52	0.50
Less Urbanized	0.87	0.88	0.69	0.71	0.44	0.45
Completely Rural	1.47	1.95	1.25	1.61	0.83	1.25

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.79P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0543		0.1300		0.8486	
GEOGRAPHIC DIVISION						
Northeast	0.1060		0.1262		0.2288	
New England	0.9647		0.9318		0.3720	
Middle Atlantic	0.0567		0.0840		0.3676	
Midwest	0.6948		0.9211		0.8146	
East North Central	0.7041		0.9572		0.9861	
West North Central	0.8784		0.9185		0.6837	
South	0.6985		0.7913		0.7699	
South Atlantic	0.9560		0.5537		0.3248	
East South Central	0.6058		0.5497		0.5496	
West South Central	0.8218		0.5719		0.0766	
West	0.0810		0.1535		0.8702	
Mountain	0.3385		0.6407		0.9604	
Pacific	0.1346		0.0582		0.8687	
COUNTY TYPE						
Large Metro	0.2190		0.1466		0.2191	
Small Metro	0.2712		0.7554		0.2867	
250K – 1 Mil. Pop.	0.3538		0.3515		0.7917	
< 250K Pop.	0.5535		0.3935		0.1474	
Nonmetro	0.2176		0.4118		0.6213	
Urbanized	0.8312		0.8197		0.8956	
Less Urbanized	0.2026		0.3077		0.8365	
Completely Rural	0.4993		0.8571		0.1009	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.80C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	154	145	118	110	77	69
GEOGRAPHIC DIVISION						
Northeast	61	60	46	43	30	24
New England	29	26	28	23	17	14
Middle Atlantic	53	53	37	35	24	20
Midwest	59	59	45	42	28	26
East North Central	50	45	38	34	24	21
West North Central	32	36	23	23	15	16
South	90	89	74	68	48	43
South Atlantic	68	64	53	48	37	31
East South Central	32	36	29	25	19	16
West South Central	52	51	41	41	24	27
West	91	85	66	66	42	40
Mountain	35	37	24	29	16	19
Pacific	83	77	61	59	39	36
COUNTY TYPE						
Large Metro	154	158	101	105	59	60
Small Metro	113	115	76	72	47	40
250K – 1 Mil. Pop.	91	92	62	59	37	31
< 250K Pop.	67	72	46	43	31	26
Nonmetro	75	77	50	48	28	27
Urbanized	53	52	34	32	17	18
Less Urbanized	56	55	36	36	21	20
Completely Rural	23	22	17	15	11	8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.80D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.47	0.44	0.36	0.34	0.23	0.21
GEOGRAPHIC DIVISION						
Northeast	1.04	1.02	0.78	0.72	0.50	0.41
New England	1.88	1.73	1.81	1.51	1.14	0.93
Middle Atlantic	1.23	1.22	0.84	0.80	0.55	0.45
Midwest	0.80	0.81	0.61	0.57	0.38	0.36
East North Central	0.97	0.89	0.74	0.68	0.47	0.42
West North Central	1.42	1.60	1.02	1.02	0.66	0.72
South	0.77	0.76	0.63	0.58	0.41	0.37
South Atlantic	1.15	1.08	0.91	0.81	0.64	0.52
East South Central	1.71	1.94	1.52	1.35	1.03	0.84
West South Central	1.33	1.31	1.05	1.04	0.61	0.68
West	1.17	1.09	0.84	0.84	0.54	0.51
Mountain	1.47	1.56	1.03	1.25	0.68	0.82
Pacific	1.51	1.40	1.11	1.08	0.71	0.65
COUNTY TYPE						
Large Metro	0.68	0.63	0.51	0.50	0.32	0.31
Small Metro	0.76	0.81	0.61	0.57	0.42	0.36
250K – 1 Mil. Pop.	0.93	0.95	0.76	0.69	0.50	0.42
< 250K Pop.	1.35	1.49	1.12	0.99	0.82	0.68
Nonmetro	0.95	1.08	0.77	0.77	0.48	0.50
Urbanized	1.36	1.72	1.16	1.20	0.69	0.76
Less Urbanized	1.43	1.51	1.16	1.12	0.75	0.74
Completely Rural	3.02	3.92	2.55	3.21	1.97	1.91

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.80P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.2228		0.2048		0.0858	
GEOGRAPHIC DIVISION						
Northeast	0.2746		0.4248		0.3366	
New England	0.4656		0.2619		0.3552	
Middle Atlantic	0.3936		0.8466		0.6187	
Midwest	0.5793		0.1368		0.7884	
East North Central	0.6574		0.7997		0.9541	
West North Central	0.1556		0.0221		0.7295	
South	0.3926		0.3790		0.1900	
South Atlantic	0.9172		0.5108		0.2688	
East South Central	0.7789		0.7983		0.6774	
West South Central	0.2551		0.5695		0.5466	
West	0.8939		0.8117		0.4706	
Mountain	0.1616		0.9539		0.9286	
Pacific	0.5750		0.8123		0.4053	
COUNTY TYPE						
Large Metro	0.3921		0.4469		0.8845	
Small Metro	0.3853		0.0749		0.0012	
250K – 1 Mil. Pop.	0.8084		0.3859		0.0009	
< 250K Pop.	0.0994		0.0846		0.2491	
Nonmetro	0.6744		0.5257		0.5787	
Urbanized	0.8736		0.4764		0.2813	
Less Urbanized	0.3965		0.9746		0.7987	
Completely Rural	0.9916		0.5876		0.7715	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.81C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	663	662	323	346	222	237
GEOGRAPHIC DIVISION						
Northeast	264	267	136	141	98	81
New England	134	134	72	50	56	35
Middle Atlantic	229	232	115	132	80	74
Midwest	253	272	121	131	79	93
East North Central	201	224	100	115	68	80
West North Central	143	159	63	62	38	47
South	387	420	203	229	138	150
South Atlantic	295	304	149	157	95	99
East South Central	139	160	74	109	48	78
West South Central	204	222	109	118	83	77
West	386	381	187	182	126	142
Mountain	153	174	96	86	70	65
Pacific	354	338	159	158	103	126
COUNTY TYPE						
Large Metro	637	603	276	281	194	193
Small Metro	361	435	166	183	96	126
250K – 1 Mil. Pop.	294	382	138	142	77	96
< 250K Pop.	218	237	98	118	59	82
Nonmetro	234	276	114	133	73	80
Urbanized	144	192	67	92	36	47
Less Urbanized	179	190	75	97	48	67
Completely Rural	83	104	54	14	42	10

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.81D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.35	0.35	0.17	0.18	0.12	0.12
GEOGRAPHIC DIVISION						
Northeast	0.74	0.75	0.38	0.40	0.28	0.23
New England	1.43	1.43	0.77	0.53	0.60	0.38
Middle Atlantic	0.87	0.89	0.44	0.51	0.31	0.28
Midwest	0.61	0.65	0.29	0.31	0.19	0.22
East North Central	0.69	0.76	0.34	0.39	0.23	0.27
West North Central	1.14	1.27	0.50	0.49	0.31	0.38
South	0.57	0.61	0.30	0.33	0.20	0.22
South Atlantic	0.81	0.82	0.41	0.42	0.26	0.27
East South Central	1.23	1.41	0.66	0.96	0.43	0.68
West South Central	1.00	1.07	0.53	0.57	0.41	0.37
West	0.91	0.88	0.44	0.42	0.30	0.33
Mountain	1.21	1.33	0.75	0.66	0.55	0.49
Pacific	1.19	1.12	0.53	0.52	0.34	0.42
COUNTY TYPE						
Large Metro	0.52	0.52	0.26	0.27	0.19	0.19
Small Metro	0.62	0.62	0.31	0.31	0.18	0.21
250K – 1 Mil. Pop.	0.75	0.78	0.39	0.35	0.22	0.24
< 250K Pop.	1.08	1.05	0.52	0.62	0.32	0.43
Nonmetro	0.62	0.74	0.34	0.40	0.22	0.24
Urbanized	1.00	1.18	0.52	0.67	0.29	0.35
Less Urbanized	0.89	1.05	0.43	0.59	0.28	0.42
Completely Rural	1.84	2.19	1.40	0.38	1.16	0.28

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.81P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.7035		0.8059		0.8933	
GEOGRAPHIC DIVISION						
Northeast	0.2502		0.9207		0.2912	
New England	0.0357		0.0787		0.2646	
Middle Atlantic	0.9772		0.3615		0.5622	
Midwest	0.1854		0.6781		0.5697	
East North Central	0.1643		0.6836		0.6739	
West North Central	0.7127		0.8831		0.6819	
South	0.1531		0.4032		0.6868	
South Atlantic	0.5033		0.8776		0.6562	
East South Central	0.0068		0.0190		0.0325	
West South Central	0.9264		0.6754		0.1476	
West	0.0205		0.3970		0.7444	
Mountain	0.6404		0.5266		0.7449	
Pacific	0.0177		0.5323		0.8529	
COUNTY TYPE						
Large Metro	0.2247		0.8636		0.4466	
Small Metro	0.7880		0.7646		0.5449	
250K – 1 Mil. Pop.	0.2487		0.2355		0.9467	
< 250K Pop.	0.2234		0.2880		0.3746	
Nonmetro	0.0319		0.1555		0.4867	
Urbanized	0.2543		0.4445		0.6331	
Less Urbanized	0.0476		0.0078		0.0296	
Completely Rural	0.9551		0.0102		0.1474	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.82C (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	696	692	354	377	236	248
GEOGRAPHIC DIVISION						
Northeast	274	276	146	152	102	84
New England	142	142	75	61	57	37
Middle Atlantic	234	238	124	139	84	75
Midwest	265	284	137	141	88	96
East North Central	211	233	114	121	76	83
West North Central	150	165	69	69	42	48
South	408	434	219	248	144	159
South Atlantic	308	314	162	174	102	108
East South Central	147	167	84	111	52	80
West South Central	216	231	119	128	86	81
West	401	398	201	201	133	150
Mountain	161	177	101	96	71	69
Pacific	366	358	172	175	110	133
COUNTY TYPE						
Large Metro	676	645	301	307	205	206
Small Metro	384	470	185	207	108	135
250K – 1 Mil. Pop.	315	405	154	164	87	102
< 250K Pop.	236	267	110	128	66	90
Nonmetro	260	302	128	150	80	87
Urbanized	163	218	81	106	42	54
Less Urbanized	201	209	87	110	52	72
Completely Rural	92	107	58	21	44	13

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.82D (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.32	0.31	0.16	0.17	0.11	0.11
GEOGRAPHIC DIVISION						
Northeast	0.66	0.67	0.35	0.37	0.25	0.20
New England	1.31	1.30	0.69	0.56	0.53	0.34
Middle Atlantic	0.77	0.78	0.41	0.46	0.27	0.25
Midwest	0.54	0.58	0.28	0.29	0.18	0.20
East North Central	0.62	0.68	0.33	0.35	0.22	0.24
West North Central	1.02	1.11	0.47	0.47	0.28	0.33
South	0.51	0.54	0.27	0.31	0.18	0.20
South Atlantic	0.73	0.73	0.38	0.41	0.24	0.25
East South Central	1.12	1.26	0.64	0.84	0.40	0.61
West South Central	0.89	0.94	0.49	0.52	0.36	0.33
West	0.80	0.78	0.40	0.39	0.26	0.29
Mountain	1.07	1.14	0.67	0.62	0.47	0.45
Pacific	1.04	1.00	0.49	0.49	0.31	0.37
COUNTY TYPE						
Large Metro	0.46	0.46	0.24	0.25	0.17	0.17
Small Metro	0.54	0.55	0.29	0.29	0.17	0.19
250K – 1 Mil. Pop.	0.67	0.68	0.35	0.33	0.20	0.21
< 250K Pop.	0.95	0.97	0.49	0.56	0.30	0.40
Nonmetro	0.57	0.67	0.32	0.38	0.21	0.23
Urbanized	0.90	1.06	0.51	0.63	0.28	0.34
Less Urbanized	0.81	0.94	0.41	0.55	0.26	0.38
Completely Rural	1.71	1.94	1.28	0.47	1.05	0.31

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.82P (REVISED) Nonmedical Use of Prescription-Type Psychotherapeutics in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.5108		0.7958		0.4960	
GEOGRAPHIC DIVISION						
Northeast	0.1795		0.8944		0.1982	
New England	0.0326		0.0346		0.1709	
Middle Atlantic	0.8367		0.4162		0.4920	
Midwest	0.2607		0.8715		0.6714	
East North Central	0.1637		0.8212		0.7260	
West North Central	0.9735		0.4862		0.7978	
South	0.2460		0.6697		0.9528	
South Atlantic	0.5383		0.9193		0.9649	
East South Central	0.0147		0.0379		0.0601	
West South Central	0.7154		0.5478		0.1147	
West	0.0229		0.4697		0.6047	
Mountain	0.4403		0.5285		0.7537	
Pacific	0.0297		0.6240		0.6752	
COUNTY TYPE						
Large Metro	0.1779		0.6698		0.4352	
Small Metro	0.5784		0.2916		0.4946	
250K – 1 Mil. Pop.	0.2303		0.0988		0.2204	
< 250K Pop.	0.4785		0.6916		0.6766	
Nonmetro	0.0424		0.1090		0.3765	
Urbanized	0.2389		0.3116		0.9961	
Less Urbanized	0.0623		0.0073		0.0374	
Completely Rural	0.9049		0.0109		0.1322	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

NOTE: Estimates in this table include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.