

Table 1.83C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	540	563	306	316	217	214
GEOGRAPHIC DIVISION						
Northeast	227	218	127	115	94	73
New England	121	106	73	51	54	32
Middle Atlantic	192	190	103	102	76	65
Midwest	205	233	122	127	76	86
East North Central	168	188	103	112	68	76
West North Central	109	136	61	58	36	39
South	312	331	188	203	134	131
South Atlantic	225	242	139	141	95	90
East South Central	124	140	74	102	45	75
West South Central	171	171	100	103	75	57
West	321	319	172	176	120	128
Mountain	148	138	92	92	70	68
Pacific	284	290	145	150	96	109
COUNTY TYPE						
Large Metro	493	482	263	250	184	169
Small Metro	301	344	151	181	93	119
250K – 1 Mil. Pop.	246	286	128	132	76	82
< 250K Pop.	189	205	88	124	56	86
Nonmetro	196	229	116	124	73	77
Urbanized	125	164	75	85	38	46
Less Urbanized	146	163	76	94	46	65
Completely Rural	72	74	51	20	43	12

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.83D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.22	0.23	0.12	0.13	0.09	0.09
GEOGRAPHIC DIVISION						
Northeast	0.49	0.48	0.28	0.25	0.20	0.16
New England	1.00	0.88	0.61	0.42	0.45	0.27
Middle Atlantic	0.57	0.56	0.30	0.30	0.23	0.19
Midwest	0.37	0.42	0.22	0.23	0.14	0.16
East North Central	0.44	0.49	0.27	0.29	0.18	0.20
West North Central	0.66	0.82	0.37	0.35	0.22	0.24
South	0.35	0.37	0.21	0.23	0.15	0.15
South Atlantic	0.48	0.51	0.30	0.30	0.20	0.19
East South Central	0.85	0.95	0.51	0.69	0.31	0.51
West South Central	0.62	0.62	0.36	0.37	0.27	0.21
West	0.57	0.56	0.31	0.31	0.21	0.22
Mountain	0.88	0.80	0.55	0.53	0.42	0.39
Pacific	0.72	0.73	0.37	0.37	0.24	0.27
COUNTY TYPE						
Large Metro	0.33	0.33	0.19	0.18	0.14	0.13
Small Metro	0.39	0.39	0.21	0.23	0.13	0.15
250K – 1 Mil. Pop.	0.49	0.48	0.26	0.25	0.16	0.16
< 250K Pop.	0.71	0.72	0.36	0.48	0.23	0.34
Nonmetro	0.39	0.49	0.26	0.29	0.17	0.18
Urbanized	0.62	0.77	0.43	0.46	0.22	0.26
Less Urbanized	0.53	0.68	0.32	0.43	0.21	0.31
Completely Rural	1.40	1.47	1.02	0.41	0.92	0.26

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.83P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.4542		0.5953		0.8009	
GEOGRAPHIC DIVISION						
Northeast	0.2108		0.2605		0.1321	
New England	0.1231		0.0411		0.0842	
Middle Atlantic	0.5884		0.8404		0.4769	
Midwest	0.1646		0.5919		0.6281	
East North Central	0.1944		0.9872		0.5190	
West North Central	0.5399		0.3009		0.8818	
South	0.7752		0.6245		0.9670	
South Atlantic	0.6665		0.5931		0.8622	
East South Central	0.1206		0.0628		0.0722	
West South Central	0.9678		0.9394		0.1834	
West	0.0926		0.7372		0.8564	
Mountain	0.2476		0.9790		0.9008	
Pacific	0.1937		0.6775		0.7578	
COUNTY TYPE						
Large Metro	0.3489		0.3692		0.3479	
Small Metro	0.1394		0.5033		0.9100	
250K – 1 Mil. Pop.	0.0178		0.0348		0.2640	
< 250K Pop.	0.4674		0.1185		0.1551	
Nonmetro	0.0352		0.1750		0.1771	
Urbanized	0.1127		0.4882		0.5955	
Less Urbanized	0.0894		0.0222		0.0333	
Completely Rural	0.8830		0.0601		0.2653	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.84C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	68	69	56	59	36	36
GEOGRAPHIC DIVISION						
Northeast	26	22	21	20	12	11
New England	11	11	9	10	6	3
Middle Atlantic	23	19	19	17	11	10
Midwest	24	28	20	22	14	14
East North Central	20	23	17	19	11	11
West North Central	14	15	11	11	9	8
South	41	43	34	37	22	24
South Atlantic	27	31	23	27	15	17
East South Central	17	17	14	14	9	9
West South Central	25	25	21	22	13	16
West	40	39	33	31	20	19
Mountain	21	16	13	15	10	9
Pacific	34	35	30	27	17	16
COUNTY TYPE						
Large Metro	54	57	46	47	30	28
Small Metro	40	40	31	33	19	21
250K – 1 Mil. Pop.	31	33	25	26	14	15
< 250K Pop.	26	25	19	21	13	15
Nonmetro	27	26	20	21	12	13
Urbanized	18	17	13	13	8	8
Less Urbanized	20	18	15	14	8	8
Completely Rural	8	8	7	7	4	5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.84D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.27	0.22	0.23	0.14	0.14
GEOGRAPHIC DIVISION						
Northeast	0.57	0.49	0.47	0.44	0.27	0.24
New England	0.93	0.99	0.79	0.86	0.47	0.26
Middle Atlantic	0.70	0.56	0.57	0.51	0.33	0.31
Midwest	0.43	0.49	0.35	0.39	0.25	0.25
East North Central	0.50	0.58	0.42	0.48	0.28	0.29
West North Central	0.82	0.89	0.66	0.68	0.53	0.51
South	0.44	0.48	0.38	0.41	0.24	0.27
South Atlantic	0.58	0.67	0.50	0.59	0.33	0.36
East South Central	1.12	1.12	0.96	0.94	0.62	0.60
West South Central	0.82	0.82	0.71	0.72	0.45	0.52
West	0.66	0.65	0.55	0.51	0.33	0.31
Mountain	1.16	0.91	0.74	0.83	0.57	0.51
Pacific	0.79	0.83	0.71	0.64	0.41	0.38
COUNTY TYPE						
Large Metro	0.37	0.38	0.32	0.32	0.21	0.20
Small Metro	0.48	0.48	0.39	0.41	0.24	0.27
250K – 1 Mil. Pop.	0.56	0.57	0.47	0.47	0.27	0.28
< 250K Pop.	0.91	0.91	0.68	0.79	0.48	0.60
Nonmetro	0.58	0.56	0.43	0.46	0.29	0.29
Urbanized	0.94	0.84	0.69	0.67	0.48	0.44
Less Urbanized	0.82	0.82	0.64	0.66	0.40	0.38
Completely Rural	1.44	1.87	1.22	1.52	0.82	1.23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.84P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0671		0.1240		0.7476	
GEOGRAPHIC DIVISION						
Northeast	0.2329		0.2852		0.3799	
New England	0.9224		0.9704		0.2769	
Middle Atlantic	0.1584		0.2203		0.6142	
Midwest	0.8626		0.9956		0.9108	
East North Central	0.7827		0.9552		0.9768	
West North Central	0.9282		0.9224		0.8240	
South	0.4998		0.6681		0.6857	
South Atlantic	0.9082		0.7105		0.6680	
East South Central	0.3291		0.2460		0.9815	
West South Central	0.7345		0.6301		0.2655	
West	0.0973		0.1068		0.5065	
Mountain	0.1145		0.7426		0.6350	
Pacific	0.2918		0.0942		0.6154	
COUNTY TYPE						
Large Metro	0.2335		0.1471		0.1460	
Small Metro	0.4716		0.8944		0.1554	
250K – 1 Mil. Pop.	0.5718		0.6444		0.4008	
< 250K Pop.	0.6668		0.3939		0.2098	
Nonmetro	0.0858		0.0939		0.9657	
Urbanized	0.3410		0.2237		0.7501	
Less Urbanized	0.1851		0.2054		0.6158	
Completely Rural	0.6826		0.9151		0.1158	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.85C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	145	136	103	97	66	60
GEOGRAPHIC DIVISION						
Northeast	59	54	42	37	24	21
New England	29	24	28	21	13	11
Middle Atlantic	51	47	30	31	20	18
Midwest	55	53	41	37	27	22
East North Central	47	42	35	31	23	19
West North Central	29	31	20	21	14	13
South	85	81	62	60	41	38
South Atlantic	61	60	45	43	30	28
East South Central	29	32	27	23	18	14
West South Central	50	46	34	34	20	20
West	86	81	60	58	36	38
Mountain	33	34	22	26	17	17
Pacific	78	73	56	52	32	34
COUNTY TYPE						
Large Metro	136	140	83	91	48	52
Small Metro	103	97	66	60	41	33
250K – 1 Mil. Pop.	83	81	55	52	35	28
< 250K Pop.	60	57	40	31	21	19
Nonmetro	66	70	43	43	25	25
Urbanized	45	46	28	29	16	16
Less Urbanized	48	51	31	30	17	17
Completely Rural	21	21	16	14	9	8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.85D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.44	0.42	0.31	0.30	0.20	0.18
GEOGRAPHIC DIVISION						
Northeast	1.01	0.91	0.71	0.63	0.41	0.36
New England	1.91	1.59	1.81	1.35	0.83	0.70
Middle Atlantic	1.18	1.08	0.70	0.70	0.46	0.41
Midwest	0.74	0.73	0.55	0.51	0.37	0.31
East North Central	0.92	0.83	0.69	0.61	0.46	0.37
West North Central	1.26	1.36	0.86	0.91	0.61	0.56
South	0.73	0.70	0.53	0.51	0.35	0.32
South Atlantic	1.03	1.01	0.76	0.73	0.51	0.48
East South Central	1.55	1.69	1.42	1.25	0.97	0.75
West South Central	1.28	1.17	0.86	0.87	0.52	0.52
West	1.10	1.03	0.77	0.74	0.46	0.48
Mountain	1.41	1.43	0.93	1.09	0.73	0.72
Pacific	1.43	1.33	1.02	0.95	0.58	0.62
COUNTY TYPE						
Large Metro	0.63	0.59	0.43	0.44	0.26	0.27
Small Metro	0.74	0.71	0.55	0.50	0.38	0.31
250K – 1 Mil. Pop.	0.90	0.86	0.70	0.64	0.49	0.39
< 250K Pop.	1.31	1.23	1.02	0.77	0.58	0.52
Nonmetro	0.88	1.04	0.68	0.71	0.44	0.45
Urbanized	1.32	1.53	1.00	1.14	0.67	0.68
Less Urbanized	1.29	1.54	1.03	0.97	0.64	0.64
Completely Rural	2.95	3.93	2.55	3.11	1.74	1.87

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.85P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.2920		0.5073		0.2155	
GEOGRAPHIC DIVISION						
Northeast	0.3016		0.5889		0.3155	
New England	0.1203		0.2245		0.1727	
Middle Atlantic	0.7189		0.8033		0.7449	
Midwest	0.7315		0.0870		0.4193	
East North Central	0.5458		0.5651		0.5459	
West North Central	0.1446		0.0284		0.6042	
South	0.3507		0.6226		0.4175	
South Atlantic	0.4198		0.4718		0.7700	
East South Central	0.8311		0.9008		0.5909	
West South Central	0.6342		0.9251		0.4911	
West	0.9696		0.3923		0.9453	
Mountain	0.2865		0.9592		0.7111	
Pacific	0.5730		0.3400		0.9076	
COUNTY TYPE						
Large Metro	0.5995		0.8830		0.6350	
Small Metro	0.3240		0.0600		0.0021	
250K – 1 Mil. Pop.	0.8374		0.5973		0.0012	
< 250K Pop.	0.0550		0.0153		0.4515	
Nonmetro	0.6030		0.6079		0.8106	
Urbanized	0.8519		0.3273		0.4496	
Less Urbanized	0.5375		0.6966		0.6363	
Completely Rural	0.9436		0.7749		0.7537	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.86C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	497	526	272	290	199	200
GEOGRAPHIC DIVISION						
Northeast	214	209	117	102	90	70
New England	110	100	67	40	53	31
Middle Atlantic	184	182	96	93	72	63
Midwest	187	221	105	115	65	82
East North Central	153	178	88	102	59	74
West North Central	101	132	54	53	30	36
South	295	309	171	188	126	121
South Atlantic	218	222	128	128	88	81
East South Central	117	134	64	99	40	73
West South Central	156	159	90	92	70	50
West	292	296	153	158	110	118
Mountain	138	128	84	83	64	63
Pacific	256	269	127	134	88	100
COUNTY TYPE						
Large Metro	445	437	235	225	171	155
Small Metro	276	305	128	157	80	108
250K – 1 Mil. Pop.	222	251	108	110	65	75
< 250K Pop.	173	180	73	112	48	77
Nonmetro	165	204	98	106	66	70
Urbanized	102	140	60	70	32	39
Less Urbanized	121	142	62	82	40	61
Completely Rural	62	70	47	13	42	8

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.86D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.26	0.28	0.14	0.15	0.11	0.11
GEOGRAPHIC DIVISION						
Northeast	0.60	0.59	0.33	0.29	0.25	0.20
New England	1.18	1.07	0.72	0.43	0.57	0.34
Middle Atlantic	0.70	0.70	0.37	0.36	0.28	0.24
Midwest	0.45	0.53	0.25	0.27	0.16	0.20
East North Central	0.52	0.61	0.30	0.35	0.20	0.25
West North Central	0.81	1.05	0.44	0.42	0.24	0.29
South	0.43	0.45	0.25	0.27	0.19	0.18
South Atlantic	0.60	0.60	0.35	0.35	0.24	0.22
East South Central	1.04	1.18	0.57	0.87	0.36	0.64
West South Central	0.77	0.77	0.44	0.45	0.34	0.24
West	0.68	0.68	0.36	0.36	0.26	0.27
Mountain	1.08	0.97	0.66	0.63	0.50	0.48
Pacific	0.86	0.89	0.43	0.44	0.29	0.33
COUNTY TYPE						
Large Metro	0.40	0.40	0.23	0.22	0.17	0.15
Small Metro	0.49	0.48	0.24	0.27	0.15	0.19
250K – 1 Mil. Pop.	0.60	0.58	0.30	0.28	0.18	0.19
< 250K Pop.	0.89	0.88	0.40	0.59	0.27	0.41
Nonmetro	0.45	0.59	0.29	0.33	0.20	0.21
Urbanized	0.72	0.95	0.47	0.53	0.25	0.29
Less Urbanized	0.61	0.82	0.35	0.50	0.23	0.39
Completely Rural	1.63	1.80	1.24	0.34	1.14	0.23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.86P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.8074		0.9419		0.8313	
GEOGRAPHIC DIVISION						
Northeast	0.3456		0.4012		0.2449	
New England	0.2022		0.0977		0.1922	
Middle Atlantic	0.7379		0.9438		0.5683	
Midwest	0.1034		0.9292		0.4154	
East North Central	0.1859		0.8008		0.3629	
West North Central	0.3306		0.8220		0.9854	
South	0.4954		0.3824		0.7948	
South Atlantic	0.8241		0.8284		0.9919	
East South Central	0.0623		0.0231		0.0385	
West South Central	0.8540		0.9783		0.1350	
West	0.1086		0.7467		0.7306	
Mountain	0.4623		0.8843		0.9693	
Pacific	0.1504		0.6262		0.6932	
COUNTY TYPE						
Large Metro	0.4789		0.4700		0.3759	
Small Metro	0.2781		0.8552		0.2786	
250K – 1 Mil. Pop.	0.0203		0.0872		0.9343	
< 250K Pop.	0.1859		0.0153		0.1184	
Nonmetro	0.0111		0.1293		0.1876	
Urbanized	0.0667		0.5820		0.7737	
Less Urbanized	0.0455		0.0090		0.0175	
Completely Rural	0.8834		0.0780		0.1948	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.87C (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	533	558	298	312	211	211
GEOGRAPHIC DIVISION						
Northeast	223	217	125	114	93	72
New England	120	105	72	49	55	32
Middle Atlantic	189	189	101	102	76	64
Midwest	201	232	119	125	74	86
East North Central	165	186	101	109	66	76
West North Central	107	137	60	58	33	38
South	314	327	183	201	131	129
South Atlantic	227	240	138	140	93	89
East South Central	124	139	73	101	44	75
West South Central	170	168	97	100	73	54
West	317	317	170	173	116	126
Mountain	147	135	90	88	67	66
Pacific	279	288	144	148	94	107
COUNTY TYPE						
Large Metro	484	474	255	246	179	167
Small Metro	296	334	149	173	91	114
250K – 1 Mil. Pop.	242	276	126	126	75	79
< 250K Pop.	184	199	85	119	54	82
Nonmetro	192	225	111	122	71	77
Urbanized	121	158	72	83	36	45
Less Urbanized	141	159	72	92	44	66
Completely Rural	71	72	50	19	43	12

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.87D (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.24	0.25	0.14	0.14	0.10	0.09
GEOGRAPHIC DIVISION						
Northeast	0.54	0.52	0.30	0.27	0.23	0.17
New England	1.10	0.97	0.66	0.45	0.50	0.30
Middle Atlantic	0.62	0.62	0.33	0.33	0.25	0.21
Midwest	0.41	0.47	0.24	0.25	0.15	0.17
East North Central	0.48	0.54	0.29	0.32	0.19	0.22
West North Central	0.72	0.92	0.41	0.39	0.22	0.26
South	0.39	0.40	0.23	0.25	0.16	0.16
South Atlantic	0.54	0.56	0.32	0.33	0.22	0.21
East South Central	0.95	1.05	0.56	0.76	0.34	0.57
West South Central	0.70	0.68	0.40	0.41	0.30	0.22
West	0.63	0.62	0.34	0.34	0.23	0.25
Mountain	0.97	0.87	0.60	0.57	0.44	0.43
Pacific	0.79	0.81	0.41	0.42	0.27	0.30
COUNTY TYPE						
Large Metro	0.36	0.36	0.21	0.20	0.15	0.14
Small Metro	0.44	0.43	0.23	0.25	0.14	0.17
250K – 1 Mil. Pop.	0.54	0.52	0.29	0.27	0.17	0.17
< 250K Pop.	0.78	0.79	0.40	0.52	0.25	0.36
Nonmetro	0.43	0.54	0.28	0.31	0.19	0.20
Urbanized	0.69	0.85	0.46	0.51	0.24	0.28
Less Urbanized	0.58	0.75	0.34	0.47	0.22	0.35
Completely Rural	1.57	1.62	1.15	0.44	1.02	0.27

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 1.87P (REVISED) Nonmedical Use of Pain Relievers in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.5865		0.8237		0.8447	
GEOGRAPHIC DIVISION						
Northeast	0.2555		0.3464		0.1650	
New England	0.1224		0.0348		0.1070	
Middle Atlantic	0.6916		0.9817		0.5245	
Midwest	0.1557		0.5928		0.6345	
East North Central	0.1783		0.9838		0.5137	
West North Central	0.5485		0.3090		0.8396	
South	0.7127		0.5484		0.9786	
South Atlantic	0.6701		0.6399		0.9242	
East South Central	0.0941		0.0358		0.0711	
West South Central	0.9956		0.9852		0.0995	
West	0.1345		0.9781		0.7742	
Mountain	0.3241		0.9308		0.9456	
Pacific	0.2355		0.9295		0.6921	
COUNTY TYPE						
Large Metro	0.4167		0.5100		0.4639	
Small Metro	0.1490		0.5180		0.8966	
250K – 1 Mil. Pop.	0.0180		0.0467		0.2023	
< 250K Pop.	0.4348		0.1461		0.2123	
Nonmetro	0.0182		0.1020		0.1799	
Urbanized	0.0809		0.3663		0.5419	
Less Urbanized	0.0555		0.0119		0.0290	
Completely Rural	0.9287		0.0659		0.1821	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.