

Table 2.1A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	173,921	172,366	85,999	84,836	72,873	70,939
Cigarettes	162,991	161,806	71,676	70,569	61,565	60,069
Smokeless Tobacco	45,832	45,091	11,295	11,030	8,231	8,051
Cigars	88,737	88,362	26,844	26,565	13,708	13,263
Pipe Tobacco ²	38,411 ^b	35,402	--	--	2,321	2,046
ALCOHOL	203,368	203,976	162,415	162,876	125,309	126,760
Binge Alcohol Use ³	--	--	--	--	56,575	57,778
Heavy Alcohol Use ³	--	--	--	--	16,946	17,010

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.1B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	70.7 ^a	69.5	35.0	34.2	29.6 ^a	28.6
Cigarettes	66.3 ^a	65.3	29.1	28.5	25.0	24.2
Smokeless Tobacco	18.6	18.2	4.6	4.5	3.3	3.2
Cigars	36.1	35.7	10.9	10.7	5.6	5.4
Pipe Tobacco ²	15.6 ^b	14.3	--	--	0.9	0.8
ALCOHOL	82.7	82.3	66.0	65.7	50.9	51.1
Binge Alcohol Use ³	--	--	--	--	23.0	23.3
Heavy Alcohol Use ³	--	--	--	--	6.9	6.9

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.2A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	874	801	486	435	175	195
Cigarettes	721	645	407 ^a	330	136	145
Smokeless Tobacco	181	187	87	99	30	34
Cigars	254	246	136	157	36	56
Pipe Tobacco ²	68	53	--	--	11	15
ALCOHOL	1,230	1,169	845	792	311	283
Binge Alcohol Use ³	--	--	--	--	117	121
Heavy Alcohol Use ³	--	--	--	--	20 ^a	6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.2B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	10.9	9.9	6.0	5.3	2.2	2.4
Cigarettes	9.0	7.9	5.1 ^a	4.1	1.7	1.8
Smokeless Tobacco	2.3	2.3	1.1	1.2	0.4	0.4
Cigars	3.1	3.0	1.7	1.9	0.5	0.7
Pipe Tobacco ²	0.8	0.7	--	--	0.1	0.2
ALCOHOL	15.3	14.4	10.5	9.7	3.9	3.5
Binge Alcohol Use ³	--	--	--	--	1.5	1.5
Heavy Alcohol Use ³	--	--	--	--	0.2 ^a	0.1

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.3A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	2,593 ^b	2,261	1,755 ^a	1,585	1,041 ^a	921
Cigarettes	2,223 ^b	1,898	1,392 ^a	1,231	801	716
Smokeless Tobacco	583	588	388	412	178	185
Cigars	1,150 ^b	971	774	702	341	290
Pipe Tobacco ²	216	183	--	--	65	70
ALCOHOL	3,662 ^a	3,374	2,933 ^a	2,685	1,378	1,249
Binge Alcohol Use ³	--	--	--	--	789 ^a	664
Heavy Alcohol Use ³	--	--	--	--	108	123

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.3B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	29.4 ^b	26.5	19.9	18.6	11.8	10.8
Cigarettes	25.2 ^b	22.3	15.8	14.4	9.1	8.4
Smokeless Tobacco	6.6	6.9	4.4	4.8	2.0	2.2
Cigars	13.0 ^a	11.4	8.8	8.2	3.9	3.4
Pipe Tobacco ²	2.5	2.1	--	--	0.7	0.8
ALCOHOL	41.5	39.6	33.3	31.5	15.6	14.7
Binge Alcohol Use ³	--	--	--	--	8.9	7.8
Heavy Alcohol Use ³	--	--	--	--	1.2	1.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.4A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	4,085	3,950	3,041	2,972	2,058	2,012
Cigarettes	3,616	3,428	2,516	2,394	1,696	1,620
Smokeless Tobacco	1,047 ^a	1,180	739	790	392	398
Cigars	2,068	2,063	1,465	1,505	669	724
Pipe Tobacco ²	356	341	--	--	92	97
ALCOHOL	5,363	5,407	4,583	4,538	2,533	2,488
Binge Alcohol Use ³	--	--	--	--	1,708	1,663
Heavy Alcohol Use ³	--	--	--	--	475	461

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.4B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	47.9	46.0	35.7	34.6	24.2	23.4
Cigarettes	42.4 ^a	39.9	29.5	27.9	19.9	18.9
Smokeless Tobacco	12.3 ^a	13.7	8.7	9.2	4.6	4.6
Cigars	24.3	24.0	17.2	17.5	7.9	8.4
Pipe Tobacco ²	4.2	4.0	--	--	1.1	1.1
ALCOHOL	62.9	63.0	53.8	52.9	29.7	29.0
Binge Alcohol Use ³	--	--	--	--	20.0	19.4
Heavy Alcohol Use ³	--	--	--	--	5.6	5.4

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.5A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	8,605	8,597	6,806	6,717	5,297	5,164
Cigarettes	7,765	7,722	5,797	5,671	4,555	4,326
Smokeless Tobacco	2,375	2,422	1,353	1,441	726	792
Cigars	5,125	5,122	3,532	3,558	1,778	1,869
Pipe Tobacco ²	918	972	--	--	185	209
ALCOHOL	10,319	10,420	9,236	9,343	6,602	6,709
Binge Alcohol Use ³	--	--	--	--	4,625	4,725
Heavy Alcohol Use ³	--	--	--	--	1,756	1,722

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.5B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	67.3 ^a	65.0	53.2 ^a	50.8	41.4 ^a	39.0
Cigarettes	60.7 ^a	58.3	45.3 ^a	42.9	35.6 ^b	32.7
Smokeless Tobacco	18.6	18.3	10.6	10.9	5.7	6.0
Cigars	40.1	38.7	27.6	26.9	13.9	14.1
Pipe Tobacco ²	7.2	7.3	--	--	1.4	1.6
ALCOHOL	80.7 ^a	78.7	72.2	70.6	51.6	50.7
Binge Alcohol Use ³	--	--	--	--	36.2	35.7
Heavy Alcohol Use ³	--	--	--	--	13.7	13.0

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.6A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	14,909	14,410	10,877	10,415	9,087 ^a	8,530
Cigarettes	14,029	13,470	9,594 ^a	9,092	8,027 ^a	7,509
Smokeless Tobacco	4,231	4,145	1,492	1,487	980	929
Cigars	8,996	8,734	4,396	4,169	2,194	2,004
Pipe Tobacco ²	1,692	1,602	--	--	246	185
ALCOHOL	17,991	17,465	16,578	16,161	13,680	13,325
Binge Alcohol Use ³	--	--	--	--	9,189	8,950
Heavy Alcohol Use ³	--	--	--	--	3,336	3,097

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.6B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	74.7	73.9	54.5	53.4	45.6 ^a	43.8
Cigarettes	70.3	69.1	48.1	46.6	40.2 ^a	38.5
Smokeless Tobacco	21.2	21.3	7.5	7.6	4.9	4.8
Cigars	45.1	44.8	22.0	21.4	11.0	10.3
Pipe Tobacco ²	8.5	8.2	--	--	1.2	1.0
ALCOHOL	90.2	89.6	83.1	82.9	68.6	68.3
Binge Alcohol Use ³	--	--	--	--	46.1	45.9
Heavy Alcohol Use ³	--	--	--	--	16.7	15.9

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.7A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	26,678	26,228	16,473	16,104	13,960	13,619
Cigarettes	24,956	24,410	13,879	13,596	11,961	11,798
Smokeless Tobacco	9,046	8,385	2,283	2,149	1,700	1,606
Cigars	15,308	14,962	5,758	5,352	2,837	2,502
Pipe Tobacco ²	3,179	2,947	--	--	316	262
ALCOHOL	31,472	31,540	27,221	27,713	21,699	22,092
Binge Alcohol Use ³	--	--	--	--	12,000	12,381
Heavy Alcohol Use ³	--	--	--	--	3,528	3,418

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.7B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	76.0	74.3	46.9	45.6	39.8	38.6
Cigarettes	71.1	69.2	39.5	38.5	34.1	33.4
Smokeless Tobacco	25.8 ^a	23.8	6.5	6.1	4.8	4.5
Cigars	43.6	42.4	16.4	15.2	8.1	7.1
Pipe Tobacco ²	9.1	8.3	--	--	0.9	0.7
ALCOHOL	89.6	89.3	77.5	78.5	61.8	62.6
Binge Alcohol Use ³	--	--	--	--	34.2	35.1
Heavy Alcohol Use ³	--	--	--	--	10.0	9.7

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.8A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	116,177	116,119	46,562	46,607	41,254	40,499
Cigarettes	109,681	110,233	38,092	38,254	34,388	33,955
Smokeless Tobacco	28,369	28,186	4,953	4,653	4,224	4,107
Cigars	55,837	56,264	10,783	11,120	5,852	5,818
Pipe Tobacco ²	31,981 ^b	29,305	--	--	1,407	1,208
ALCOHOL	133,331	134,601	101,019	101,643	79,105	80,613
Binge Alcohol Use ³	--	--	--	--	28,147	29,275
Heavy Alcohol Use ³	--	--	--	--	7,723	8,182

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.8B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	76.0	75.1	30.5	30.2	27.0	26.2
Cigarettes	71.8	71.3	24.9	24.7	22.5	22.0
Smokeless Tobacco	18.6	18.2	3.2	3.0	2.8	2.7
Cigars	36.5	36.4	7.1	7.2	3.8	3.8
Pipe Tobacco ²	20.9 ^b	19.0	--	--	0.9	0.8
ALCOHOL	87.3	87.1	66.1	65.8	51.8	52.2
Binge Alcohol Use ³	--	--	--	--	18.4	18.9
Heavy Alcohol Use ³	--	--	--	--	5.1	5.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.9A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	93,500	92,802	51,249	50,742	43,389	42,369
Cigarettes	84,793	84,613	38,802	38,303	33,220	32,607
Smokeless Tobacco	38,579	38,204	10,469	10,117	7,843	7,589
Cigars	65,476	65,494	21,003	21,088	11,092	10,940
Pipe Tobacco ²	33,727 ^b	30,987	--	--	2,023	1,797
ALCOHOL	102,646	102,824	83,275	83,583	68,025	68,088
Binge Alcohol Use ³	--	--	--	--	37,298	38,128
Heavy Alcohol Use ³	--	--	--	--	12,775	12,786

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.9B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	78.3 ^a	77.2	42.9	42.2	36.4	35.2
Cigarettes	71.0	70.4	32.5	31.8	27.8	27.1
Smokeless Tobacco	32.3	31.8	8.8	8.4	6.6	6.3
Cigars	54.9	54.5	17.6	17.5	9.3	9.1
Pipe Tobacco ²	28.3 ^b	25.8	--	--	1.7	1.5
ALCOHOL	86.0	85.5	69.8	69.5	57.0	56.6
Binge Alcohol Use ³	--	--	--	--	31.2	31.7
Heavy Alcohol Use ³	--	--	--	--	10.7	10.6

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.10A Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	80,422	79,564	34,751	34,094	29,484	28,570
Cigarettes	78,198	77,193	32,875	32,266	28,345	27,462
Smokeless Tobacco	7,253	6,887	826	913	388	461
Cigars	23,261	22,869	5,841	5,477	2,616	2,323
Pipe Tobacco ²	4,684	4,415	--	--	298	249
ALCOHOL	100,722	101,152	79,140	79,293	57,283	58,672
Binge Alcohol Use ³	--	--	--	--	19,276	19,651
Heavy Alcohol Use ³	--	--	--	--	4,172	4,225

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.10B Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	63.5	62.4	27.4	26.7	23.3	22.4
Cigarettes	61.7	60.5	26.0	25.3	22.4	21.5
Smokeless Tobacco	5.7	5.4	0.7	0.7	0.3	0.4
Cigars	18.4	17.9	4.6	4.3	2.1	1.8
Pipe Tobacco ²	3.7	3.5	--	--	0.2	0.2
ALCOHOL	79.5	79.3	62.5	62.2	45.2	46.0
Binge Alcohol Use ³	--	--	--	--	15.2	15.4
Heavy Alcohol Use ³	--	--	--	--	3.3	3.3

*Low precision; no estimate reported.

-- Not available.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.