

Table 2.47A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	173,904	172,420	86,034	85,017	72,878	71,073
GEOGRAPHIC DIVISION						
Northeast	32,073	31,305	15,058	14,665	12,450	12,041
New England	8,767	8,594	3,777	3,803	3,168	3,046
Middle Atlantic	23,305	22,711	11,281	10,862	9,282	8,996
Midwest	39,911	40,434	20,752	20,747	17,628	17,541
East North Central	27,588	28,056	14,566	14,323	12,317	12,208
West North Central	12,323	12,377	6,186	6,424	5,311	5,333
South	63,589	63,153	32,901	31,976	28,698 ^a	27,253
South Atlantic	33,637	33,667	16,180	16,654	13,938	14,213
East South Central	11,126 ^a	10,690	6,216 ^a	5,537	5,518 ^a	4,873
West South Central	18,826	18,796	10,504	9,785	9,241 ^b	8,167
West	38,331	37,528	17,324	17,630	14,102	14,237
Mountain	11,975	11,959	5,850	5,882	4,767	4,969
Pacific	26,356	25,569	11,474	11,748	9,335	9,268
COUNTY TYPE						
Large Metro	91,736	89,251	43,844	43,191	36,330	35,139
Small Metro	50,556	52,864	25,534	25,912	21,768	21,953
250K – 1 Mil. Pop.	32,816	35,006	16,516	16,989	13,997	14,368
< 250K Pop.	17,740	17,859	9,018	8,923	7,771	7,585
Nonmetro	31,612	30,304	16,656	15,914	14,780	13,980
Urbanized	12,058	12,424	6,044	6,597	5,244	5,693
Less Urbanized	16,094	14,591	8,640	7,708	7,741	6,885
Completely Rural	3,461	3,289	1,972	1,609	1,796	1,402

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.47B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	70.7 ^a	69.6	35.0	34.3	29.6 ^a	28.7
GEOGRAPHIC DIVISION						
Northeast	69.9	68.2	32.8	32.0	27.2	26.2
New England	72.9	71.5	31.4	31.6	26.4	25.3
Middle Atlantic	68.9	67.1	33.3	32.1	27.4	26.6
Midwest	73.0	73.8	37.9	37.9	32.2	32.0
East North Central	72.1	73.3	38.1	37.4	32.2	31.9
West North Central	75.0	75.0	37.6	38.9	32.3	32.3
South	71.5	70.2	37.0	35.6	32.2 ^a	30.3
South Atlantic	71.5	70.8	34.4	35.0	29.6	29.9
East South Central	76.1 ^a	72.6	42.5 ^b	37.6	37.8 ^b	33.1
West South Central	68.9	68.0	38.4 ^a	35.4	33.8 ^b	29.6
West	67.9 ^a	65.6	30.7	30.8	25.0	24.9
Mountain	71.0	69.2	34.7	34.1	28.2	28.8
Pacific	66.6	64.0	29.0	29.4	23.6	23.2
COUNTY TYPE						
Large Metro	68.8 ^a	67.3	32.9	32.6	27.3	26.5
Small Metro	71.6	71.3	36.2	35.0	30.8	29.6
250K – 1 Mil. Pop.	70.7	70.6	35.6	34.3	30.2	29.0
< 250K Pop.	73.2	72.8	37.2	36.4	32.1	30.9
Nonmetro	75.0	73.8	39.5	38.7	35.1	34.0
Urbanized	73.5	73.9	36.9	39.3	32.0	33.9
Less Urbanized	76.1	74.2	40.8	39.2	36.6	35.0
Completely Rural	75.3	71.5	42.9 ^a	35.0	39.1 ^b	30.5

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.48A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	7,565 ^b	7,027	5,295 ^a	5,010	3,282	3,141
GEOGRAPHIC DIVISION						
Northeast	1,293 ^b	1,119	979 ^b	820	610 ^a	517
New England	334	315	254	227	169	146
Middle Atlantic	959 ^b	804	725 ^b	593	441	371
Midwest	1,718	1,683	1,199	1,235	784	783
East North Central	1,183	1,206	816	876	533	572
West North Central	536	476	382	358	251	211
South	2,935	2,774	1,999	1,941	1,206	1,247
South Atlantic	1,387	1,341	941	953	555	629
East South Central	602	547	410	369	264	242
West South Central	946	886	648	618	388	376
West	1,618 ^a	1,451	1,119	1,014	682	594
Mountain	551	507	394	353	236	222
Pacific	1,067	944	725	661	446	372
COUNTY TYPE						
Large Metro	3,739 ^b	3,369	2,631 ^a	2,406	1,623 ^a	1,439
Small Metro	2,303	2,176	1,626	1,527	1,012	949
250K – 1 Mil. Pop.	1,465	1,415	1,036	981	632	612
< 250K Pop.	838	761	591	546	379	337
Nonmetro	1,523	1,482	1,037	1,076	647 ^a	753
Urbanized	567	612	392	442	243 ^a	313
Less Urbanized	779	733	530	540	331	382
Completely Rural	177	137	115	95	74	58

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.48B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	29.8 ^b	27.8	20.9	19.8	12.9	12.4
GEOGRAPHIC DIVISION						
Northeast	28.7 ^b	25.1	21.7 ^b	18.4	13.5 ^a	11.6
New England	28.7	27.3	21.8	19.6	14.5	12.7
Middle Atlantic	28.6 ^b	24.3	21.6 ^b	18.0	13.2	11.2
Midwest	30.3	30.0	21.1	22.0	13.8	13.9
East North Central	29.7	30.6	20.5	22.2	13.4	14.5
West North Central	31.7	28.5	22.6	21.4	14.9	12.6
South	32.1	30.4	21.8	21.3	13.2	13.7
South Atlantic	29.8	28.9	20.2	20.6	11.9	13.6
East South Central	40.5	36.9	27.6	24.9	17.8	16.3
West South Central	31.4	29.5	21.5	20.6	12.8	12.5
West	26.7 ^a	24.0	18.5	16.8	11.3	9.8
Mountain	30.7	28.1	21.9	19.5	13.2	12.3
Pacific	25.0	22.3	17.0	15.6	10.5	8.8
COUNTY TYPE						
Large Metro	27.3 ^b	24.6	19.2 ^a	17.6	11.8 ^a	10.5
Small Metro	30.5	29.5	21.5	20.7	13.4	12.9
250K – 1 Mil. Pop.	28.9	28.4	20.4	19.7	12.5	12.3
< 250K Pop.	33.7	31.8	23.8	22.8	15.3	14.1
Nonmetro	36.9	35.5	25.1	25.8	15.7 ^a	18.1
Urbanized	34.0	35.6	23.5	25.7	14.6 ^a	18.2
Less Urbanized	39.2	36.2	26.7	26.6	16.6	18.9
Completely Rural	37.7	32.0	24.6	22.2	15.7	13.6

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.49A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	23,525 ^a	23,016	17,695 ^a	17,156	14,396 ^b	13,700
GEOGRAPHIC DIVISION						
Northeast	4,184	4,065	3,203 ^a	3,006	2,605 ^a	2,368
New England	1,148	1,138	900	853	732	659
Middle Atlantic	3,036	2,927	2,304	2,153	1,874	1,709
Midwest	5,635	5,501	4,405 ^a	4,226	3,680 ^b	3,441
East North Central	3,908	3,800	3,057	2,939	2,540 ^a	2,400
West North Central	1,727	1,700	1,348	1,287	1,140 ^a	1,042
South	8,373	8,167	6,276	6,075	5,225	4,995
South Atlantic	4,187	4,156	3,090	3,077	2,578	2,491
East South Central	1,457 ^b	1,331	1,161 ^a	1,054	981	905
West South Central	2,728	2,680	2,025	1,945	1,665	1,599
West	5,333	5,283	3,811	3,848	2,886	2,895
Mountain	1,677	1,635	1,225	1,222	981	985
Pacific	3,657	3,648	2,586	2,626	1,905	1,911
COUNTY TYPE						
Large Metro	12,193	11,733	8,987	8,575	7,220	6,758
Small Metro	7,399	7,409	5,609	5,566	4,519	4,443
250K – 1 Mil. Pop.	4,844	4,766	3,660	3,603	2,920	2,879
< 250K Pop.	2,555	2,644	1,949	1,963	1,600	1,564
Nonmetro	3,933	3,874	3,099	3,015	2,657	2,499
Urbanized	1,648	1,698	1,275	1,326	1,068	1,095
Less Urbanized	1,897	1,878	1,510	1,456	1,304	1,206
Completely Rural	388	298	314	233	285 ^a	197

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.49B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	71.9 ^a	70.3	54.0 ^a	52.4	44.0 ^b	41.9
GEOGRAPHIC DIVISION						
Northeast	71.4	68.9	54.6 ^a	50.9	44.4 ^b	40.1
New England	75.2	74.5	59.0	55.9	48.0	43.2
Middle Atlantic	70.0	66.9	53.1 ^a	49.2	43.2 ^a	39.1
Midwest	76.3	75.5	59.7	58.0	49.8 ^a	47.2
East North Central	76.6	75.6	59.9	58.5	49.8	47.7
West North Central	75.7	75.2	59.1	56.9	50.0	46.1
South	71.7	69.9	53.7	52.0	44.7	42.7
South Atlantic	71.3	70.4	52.6	52.1	43.9	42.2
East South Central	76.9 ^a	71.2	61.3	56.4	51.8	48.4
West South Central	69.7	68.5	51.7	49.7	42.5	40.9
West	68.3	67.3	48.8	49.0	36.9	36.9
Mountain	71.2	69.4	52.0	51.9	41.7	41.8
Pacific	67.0	66.4	47.4	47.8	34.9	34.8
COUNTY TYPE						
Large Metro	69.5	67.8	51.2	49.5	41.1 ^a	39.0
Small Metro	73.1	71.8	55.4	53.9	44.6	43.0
250K – 1 Mil. Pop.	72.6	70.4	54.8	53.2	43.7	42.5
< 250K Pop.	74.1	74.4	56.5	55.2	46.4	44.0
Nonmetro	77.7	76.0	61.2	59.1	52.5 ^a	49.0
Urbanized	77.0	74.5	59.6	58.2	49.9	48.0
Less Urbanized	78.5	76.7	62.5	59.5	54.0 ^a	49.3
Completely Rural	76.5	80.2	61.9	62.7	56.1	53.0

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.50A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	142,814	142,377	63,044	62,852	55,200	54,232
GEOGRAPHIC DIVISION						
Northeast	26,595	26,121	10,876	10,838	9,235	9,156
New England	7,285	7,141	2,623	2,723	2,267	2,240
Middle Atlantic	19,310	18,980	8,253	8,116	6,968	6,916
Midwest	32,558	33,250	15,149	15,286	13,164	13,317
East North Central	22,497	23,049	10,693	10,507	9,245	9,236
West North Central	10,061	10,201	4,456	4,779	3,919	4,081
South	52,281	52,212	24,625	23,960	22,267	21,011
South Atlantic	28,062	28,170	12,149	12,624	10,805	11,094
East South Central	9,068	8,813	4,645 ^a	4,113	4,273 ^a	3,725
West South Central	15,151	15,230	7,831	7,222	7,188 ^b	6,192
West	31,380	30,794	12,395	12,767	10,534	10,747
Mountain	9,747	9,816	4,231	4,307	3,550	3,762
Pacific	21,633	20,978	8,164	8,460	6,985	6,985
COUNTY TYPE						
Large Metro	75,805	74,149	32,226	32,210	27,487	26,942
Small Metro	40,854	43,279	18,299	18,819	16,237	16,562
250K – 1 Mil. Pop.	26,507 ^a	28,825	11,820	12,405	10,445	10,878
< 250K Pop.	14,347	14,454	6,479	6,414	5,792	5,684
Nonmetro	26,156	24,948	12,520	11,823	11,476	10,728
Urbanized	9,842	10,114	4,377	4,829	3,933	4,285
Less Urbanized	13,418	11,980	6,600 ^a	5,712	6,105 ^a	5,296
Completely Rural	2,896	2,854	1,543	1,281	1,438	1,147

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.50B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	76.0	75.0	33.6	33.1	29.4	28.6
GEOGRAPHIC DIVISION						
Northeast	75.0	73.5	30.7	30.5	26.0	25.8
New England	78.1	76.5	28.1	29.1	24.3	24.0
Middle Atlantic	73.9	72.5	31.6	31.0	26.7	26.4
Midwest	78.2	79.4	36.4	36.5	31.6	31.8
East North Central	77.1	78.6	36.6	35.8	31.7	31.5
West North Central	80.7	81.2	35.8	38.0	31.5	32.5
South	76.7	75.5	36.1	34.7	32.7 ^a	30.4
South Atlantic	76.8	76.1	33.3	34.1	29.6	30.0
East South Central	80.7	77.4	41.3 ^a	36.1	38.0 ^a	32.7
West South Central	74.3	73.5	38.4 ^a	34.9	35.2 ^b	29.9
West	73.6	71.1	29.1	29.5	24.7	24.8
Mountain	76.6	74.9	33.2	32.9	27.9	28.7
Pacific	72.4	69.4	27.3	28.0	23.4	23.1
COUNTY TYPE						
Large Metro	74.3	72.9	31.6	31.7	26.9	26.5
Small Metro	77.2	76.7	34.6	33.4	30.7	29.4
250K – 1 Mil. Pop.	76.5	76.3	34.1	32.8	30.1	28.8
< 250K Pop.	78.4	77.7	35.4	34.5	31.7	30.6
Nonmetro	79.4	78.4	38.0	37.2	34.8	33.7
Urbanized	78.2	78.9	34.8	37.7	31.2	33.4
Less Urbanized	80.1	78.8	39.4	37.6	36.5	34.9
Completely Rural	80.0	75.1	42.6 ^a	33.7	39.7 ^a	30.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.51A (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	166,339	165,393	80,739	80,008	69,596	67,932
GEOGRAPHIC DIVISION						
Northeast	30,779	30,186	14,079	13,845	11,840	11,524
New England	8,433	8,279	3,523	3,576	2,999	2,899
Middle Atlantic	22,346	21,907	10,556	10,269	8,842	8,625
Midwest	38,193	38,751	19,553	19,512	16,844	16,759
East North Central	26,405	26,850	13,750	13,446	11,785	11,636
West North Central	11,788	11,901	5,804	6,066	5,059	5,123
South	60,654	60,379	30,902	30,035	27,491 ^a	26,006
South Atlantic	32,250	32,326	15,239	15,701	13,383	13,585
East South Central	10,525	10,143	5,806 ^a	5,167	5,254 ^a	4,631
West South Central	17,879	17,910	9,856	9,167	8,854 ^b	7,791
West	36,714	36,077	16,206	16,616	13,420	13,642
Mountain	11,424	11,452	5,456	5,530	4,531	4,747
Pacific	25,290	24,625	10,749	11,086	8,889	8,895
COUNTY TYPE						
Large Metro	87,997	85,882	41,212	40,785	34,707	33,700
Small Metro	48,254	50,689	23,908	24,385	20,756	21,005
250K – 1 Mil. Pop.	31,351	33,591	15,481	16,008	13,365	13,756
< 250K Pop.	16,902	17,097	8,428	8,376	7,391	7,248
Nonmetro	30,089	28,822	15,619	14,838	14,133	13,227
Urbanized	11,490	11,812	5,652	6,155	5,001	5,380
Less Urbanized	15,314	13,858	8,110 ^a	7,168	7,410 ^a	6,503
Completely Rural	3,284	3,152	1,857	1,514	1,723	1,344

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.51B (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	75.4 ^a	74.3	36.6	35.9	31.5 ^a	30.5
GEOGRAPHIC DIVISION						
Northeast	74.5	72.9	34.1	33.4	28.6	27.8
New England	77.7	76.2	32.4	32.9	27.6	26.7
Middle Atlantic	73.3	71.7	34.6	33.6	29.0	28.2
Midwest	77.9	78.8	39.9	39.7	34.4	34.1
East North Central	77.0	78.1	40.1	39.1	34.4	33.9
West North Central	80.0	80.3	39.4	40.9	34.3	34.6
South	76.0	74.7	38.7	37.2	34.4 ^a	32.2
South Atlantic	76.1	75.3	35.9	36.6	31.6	31.6
East South Central	80.2 ^a	76.5	44.2 ^b	39.0	40.0 ^a	34.9
West South Central	73.5	72.7	40.5 ^a	37.2	36.4 ^b	31.6
West	72.8 ^a	70.5	32.1	32.5	26.6	26.7
Mountain	75.7	74.0	36.2	35.7	30.0	30.7
Pacific	71.6	68.9	30.4	31.0	25.2	24.9
COUNTY TYPE						
Large Metro	73.6 ^a	72.2	34.5	34.3	29.0	28.3
Small Metro	76.5	76.0	37.9	36.5	32.9	31.5
250K – 1 Mil. Pop.	75.9	75.4	37.5	35.9	32.3	30.9
< 250K Pop.	77.7	77.2	38.8	37.8	34.0	32.7
Nonmetro	79.1	78.1	41.1	40.2	37.2	35.8
Urbanized	78.0	78.3	38.4	40.8	33.9	35.7
Less Urbanized	79.9	78.5	42.3	40.6	38.7	36.9
Completely Rural	79.5	75.6	45.0 ^a	36.3	41.7 ^a	32.2

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.