

Table 2.11C Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	827	855	878	878	848	844
12	21	23	16	14	7	10
13	30	32	23	23	15	16
14	39	43	31	36	24	27
15	55	51	45	42	35	31
16	61	61	52	53	39	40
17	65	64	55	56	45	48
18	98	94	87	82	73	71
19	96	97	87	84	73	64
20	95	105	79	87	68	74
21	103	100	87	85	76	77
22	90	91	76	80	69	70
23	89	90	79	74	71	65
24	89	94	71	80	64	71
25	88	85	74	71	69	64
26-29	354	366	276	279	246	253
30-34	372	367	270	274	246	250
35-39	419	439	294	298	275	275
40-44	455	434	303	304	283	265
45-49	464	457	315	321	303	299
50-54	578	656	377	410	357	390
55-59	507	579	292	341	279	316
60-64	453	493	244	282	238	263
65 or Older	391	440	320	325	302	308

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.11D Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.34	0.34	0.36	0.35	0.34	0.34
12	0.52	0.55	0.40	0.35	0.18	0.24
13	0.67	0.72	0.55	0.55	0.36	0.39
14	0.83	0.93	0.68	0.80	0.55	0.62
15	0.96	0.96	0.86	0.84	0.71	0.66
16	1.04	1.08	1.01	1.03	0.82	0.80
17	1.05	1.09	1.02	1.09	0.91	1.03
18	1.18	1.14	1.21	1.15	1.18	1.13
19	1.17	1.30	1.26	1.32	1.25	1.21
20	1.21	1.17	1.36	1.36	1.33	1.34
21	1.09	1.23	1.21	1.33	1.23	1.34
22	1.11	1.24	1.31	1.30	1.29	1.29
23	1.11	1.10	1.31	1.25	1.27	1.24
24	1.15	1.21	1.26	1.33	1.20	1.33
25	1.09	1.14	1.25	1.30	1.28	1.31
26-29	1.06	1.04	1.14	1.21	1.10	1.21
30-34	1.00	1.00	1.07	1.08	1.04	1.09
35-39	1.10	1.03	1.17	1.10	1.14	1.06
40-44	1.00	1.04	1.08	1.14	1.04	1.05
45-49	0.93	0.91	1.13	1.08	1.09	1.06
50-54	1.36	1.38	1.44	1.51	1.41	1.47
55-59	1.34	1.56	1.40	1.56	1.36	1.48
60-64	1.56	1.58	1.59	1.80	1.57	1.70
65 or Older	1.10	1.21	0.90	0.90	0.85	0.85

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.11P Tobacco Product Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.0123		0.1309		0.0306
12	0.3229		0.4965		0.2938
13	0.2363		0.2125		0.7707
14	0.1035		0.2488		0.3400
15	0.0070		0.2616		0.2124
16	0.3926		0.4972		0.4089
17	0.0999		0.4810		0.8005
18	0.0041		0.0358		0.4062
19	0.4047		0.1521		0.0276
20	0.8186		0.5616		0.3011
21	0.0633		0.1957		0.2493
22	0.5900		0.8847		0.6186
23	0.4491		0.6386		0.1650
24	0.6502		0.3977		0.3845
25	0.6962		0.0476		0.0047
26-29	0.5649		0.4358		0.4628
30-34	0.0864		0.4247		0.4681
35-39	0.8266		0.9011		0.9660
40-44	0.2941		0.7567		0.1141
45-49	0.8092		0.6368		0.5663
50-54	0.8379		0.6906		0.9594
55-59	0.4627		0.4898		0.3932
60-64	0.5328		0.6326		1.0000
65 or Older	0.0591		0.5909		0.7146

*Low precision; no estimate reported.

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.12C Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	863	879	840	848	804	808
12	19	20	14	13	6	8
13	28	29	22	20	14	15
14	36	38	29	29	22	22
15	51	48	40	38	30	28
16	58	57	50	46	36	35
17	61	57	50	49	41	43
18	88	89	74	75	65	63
19	88	92	76	78	66	60
20	90	97	74	78	65	68
21	95	93	79	76	69	68
22	87	87	69	75	64	65
23	85	87	74	71	67	64
24	84	92	66	72	61	66
25	84	80	67	63	64	58
26-29	342	351	256	255	229	236
30-34	360	350	243	246	223	234
35-39	392	416	263	272	249	252
40-44	434	420	275	266	265	236
45-49	446	443	288	288	283	274
50-54	565	648	354	377	334	357
55-59	499	570	268	320	253	299
60-64	446	483	224	255	220	246
65 or Older	416	439	288	289	266	255

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.12D Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.35	0.35	0.34	0.34	0.33	0.33
12	0.49	0.49	0.36	0.31	0.15	0.19
13	0.63	0.66	0.52	0.48	0.32	0.35
14	0.78	0.84	0.65	0.67	0.51	0.51
15	0.92	0.93	0.79	0.77	0.61	0.60
16	1.05	1.04	1.00	0.91	0.76	0.73
17	1.04	1.06	0.98	1.00	0.84	0.93
18	1.19	1.19	1.13	1.18	1.11	1.09
19	1.20	1.32	1.25	1.32	1.25	1.16
20	1.37	1.26	1.41	1.35	1.31	1.29
21	1.08	1.25	1.19	1.30	1.21	1.26
22	1.20	1.28	1.32	1.32	1.28	1.30
23	1.15	1.15	1.30	1.27	1.25	1.25
24	1.19	1.26	1.26	1.28	1.20	1.30
25	1.14	1.16	1.23	1.25	1.27	1.22
26-29	1.13	1.08	1.15	1.21	1.09	1.17
30-34	1.06	1.08	1.05	1.06	0.99	1.07
35-39	1.13	1.08	1.11	1.05	1.06	0.99
40-44	1.03	1.09	1.03	1.04	1.01	0.95
45-49	0.96	0.97	1.08	1.04	1.06	1.01
50-54	1.44	1.41	1.41	1.46	1.36	1.40
55-59	1.43	1.60	1.34	1.50	1.28	1.43
60-64	1.61	1.64	1.51	1.67	1.48	1.62
65 or Older	1.17	1.21	0.81	0.80	0.75	0.70

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.12P Cigarette Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0438		0.1568		0.0749	
12	0.4692		0.3443		0.4666	
13	0.0976		0.0294		0.9232	
14	0.0250		0.0806		0.2995	
15	0.0123		0.3443		0.4622	
16	0.5328		0.2065		0.1538	
17	0.0061		0.2923		0.7337	
18	0.0175		0.0329		0.0808	
19	0.3628		0.4132		0.0286	
20	0.4921		0.2352		0.2202	
21	0.0164		0.0240		0.1121	
22	0.1538		1.0000		0.6221	
23	0.9711		0.6155		0.3096	
24	0.9142		0.4340		0.5605	
25	0.9205		0.0313		0.0121	
26-29	0.2798		0.2790		0.6254	
30-34	0.1641		0.8570		0.7351	
35-39	0.6479		0.3393		0.7428	
40-44	0.1757		0.2614		0.0390	
45-49	0.5557		0.8887		0.9906	
50-54	0.1970		0.8769		1.0000	
55-59	0.7652		0.7452		0.7320	
60-64	0.6422		0.8694		0.8910	
65 or Older	0.1395		0.7731		0.6335	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.13C Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	688	703	325	312	284	281
12	10	10	6	7	4	4
13	13	14	10	11	5	7
14	18	21	14	19	9	15
15	25	25	20	21	14	14
16	28	35	24	30	17	20
17	32	34	27	28	19	19
18	51	42	42	33	29	24
19	46	45	32	37	24	26
20	46	53	30	39	20	29
21	50	46	30	30	23	21
22	44	45	25	28	21	19
23	44	44	25	26	20	19
24	45	50	26	26	20	20
25	45	41	24	24	21	18
26-29	187	186	95	96	80	82
30-34	207	196	93	90	77	80
35-39	244	235	112	115	106	108
40-44	279	229	108	107	100	92
45-49	213	215	84	80	80	78
50-54	259	297	99	89	93	88
55-59	231	233	110	46	91	46
60-64	180	197	49	62	49	53
65 or Older	310	346	138	142	126	142

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.13D Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.28	0.28	0.13	0.13	0.12	0.11
12	0.25	0.26	0.16	0.16	0.11	0.10
13	0.32	0.34	0.23	0.27	0.12	0.16
14	0.41	0.50	0.33	0.45	0.21	0.36
15	0.53	0.54	0.44	0.47	0.30	0.31
16	0.61	0.76	0.54	0.64	0.38	0.44
17	0.69	0.76	0.60	0.65	0.44	0.46
18	0.96	0.78	0.82	0.65	0.61	0.49
19	0.95	0.92	0.70	0.77	0.55	0.58
20	1.00	1.07	0.70	0.85	0.48	0.67
21	1.01	0.99	0.67	0.69	0.53	0.51
22	1.01	1.04	0.63	0.70	0.52	0.50
23	1.01	1.00	0.62	0.65	0.49	0.47
24	1.02	1.08	0.63	0.62	0.49	0.50
25	1.05	1.00	0.62	0.63	0.52	0.46
26-29	0.98	0.99	0.56	0.57	0.47	0.49
30-34	0.95	0.89	0.48	0.45	0.40	0.41
35-39	1.05	0.94	0.52	0.53	0.50	0.50
40-44	1.06	0.92	0.47	0.49	0.44	0.43
45-49	0.85	0.84	0.38	0.34	0.36	0.34
50-54	1.11	1.26	0.47	0.41	0.44	0.40
55-59	1.19	1.19	0.62	0.25	0.51	0.25
60-64	1.25	1.37	0.36	0.46	0.36	0.39
65 or Older	0.87	0.95	0.39	0.39	0.35	0.39

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.13P Smokeless Tobacco Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD				
	Lifetime		Past Year		Past Month
	2006	2007	2006	2007	2006 2007
TOTAL	0.2734		0.4303		0.5340
12	0.4674		0.7884		0.4050
13	0.4419		0.3397		0.2880
14	0.5024		0.2026		0.2479
15	0.8270		0.7780		0.7075
16	0.1822		0.7951		0.6231
17	0.0929		0.3137		0.7003
18	0.0239		0.0655		0.4538
19	0.8903		0.1872		0.4846
20	0.1250		0.1108		0.2033
21	0.8813		0.6668		0.6847
22	0.3744		0.1510		0.4640
23	0.4185		0.6609		0.4650
24	0.4742		0.5390		0.7314
25	0.5906		0.9389		0.3247
26-29	0.4905		0.5535		0.4670
30-34	0.0241		0.5835		0.8359
35-39	0.9910		0.5446		0.5115
40-44	0.6070		0.1481		0.3218
45-49	0.7706		0.3389		0.3734
50-54	0.4814		0.4360		0.7888
55-59	0.3938		0.0050		0.0217
60-64	0.3341		0.3318		0.8548
65 or Older	0.8384		0.4061		0.9494

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.14C Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	872	872	468	484	352	337
12	13	9	10	8	2	5
13	16	19	12	15	7	9
14	26	26	21	22	15	13
15	37	34	30	29	19	19
16	42	41	33	35	22	22
17	43	47	38	38	25	28
18	73	69	63	60	41	43
19	75	72	62	57	42	41
20	72	81	49	61	34	43
21	80	78	61	57	39	42
22	68	67	55	48	37	34
23	66	70	44	48	31	28
24	70	75	43	46	31	31
25	66	64	45	43	34	27
26-29	274	282	167	154	123	105
30-34	274	268	153	157	105	105
35-39	317	299	160	153	112	109
40-44	336	289	160	149	108	99
45-49	314	314	154	168	111	116
50-54	414	435	175	213	122	146
55-59	357	386	148	132	115	108
60-64	305	328	99	133	84	74
65 or Older	381	400	124	122	100	109

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.14D Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.35	0.35	0.19	0.20	0.14	0.14
12	0.33	0.23	0.25	0.20	0.06	0.13
13	0.38	0.44	0.29	0.36	0.17	0.22
14	0.58	0.61	0.49	0.51	0.34	0.31
15	0.76	0.71	0.63	0.62	0.41	0.43
16	0.87	0.81	0.74	0.72	0.49	0.48
17	0.90	0.95	0.82	0.82	0.56	0.63
18	1.18	1.09	1.10	1.00	0.78	0.81
19	1.28	1.23	1.16	1.09	0.89	0.86
20	1.24	1.30	1.03	1.18	0.81	0.95
21	1.25	1.27	1.14	1.12	0.80	0.90
22	1.34	1.28	1.18	1.07	0.87	0.82
23	1.20	1.28	0.97	1.09	0.74	0.67
24	1.26	1.33	0.97	1.04	0.74	0.75
25	1.24	1.25	1.04	1.02	0.82	0.70
26-29	1.19	1.18	0.90	0.82	0.70	0.61
30-34	1.12	1.05	0.75	0.76	0.54	0.53
35-39	1.17	1.05	0.74	0.68	0.53	0.50
40-44	1.17	1.04	0.68	0.65	0.47	0.46
45-49	1.08	1.04	0.67	0.70	0.49	0.50
50-54	1.49	1.51	0.81	0.95	0.57	0.66
55-59	1.55	1.71	0.81	0.70	0.64	0.58
60-64	1.76	1.90	0.72	0.97	0.62	0.55
65 or Older	1.07	1.10	0.35	0.34	0.28	0.30

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.14P Cigar Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.4074		0.4718		0.2695	
12	0.3139		0.8716		0.0522	
13	0.7413		0.3121		0.4489	
14	0.1700		0.2359		0.1172	
15	0.0539		0.7821		0.6953	
16	0.3839		0.8468		0.6338	
17	0.5797		0.6100		0.2729	
18	0.0562		0.1723		0.5186	
19	0.1457		0.2775		0.9023	
20	0.3065		0.2598		0.9928	
21	0.4406		0.8416		0.3281	
22	0.1893		0.7472		0.9657	
23	0.3898		0.9268		0.1741	
24	0.5842		0.4622		0.3632	
25	0.5182		0.1490		0.0176	
26-29	0.6897		0.3029		0.1870	
30-34	0.2737		0.2640		0.3203	
35-39	0.9704		0.8405		0.9282	
40-44	0.5610		0.4014		0.9729	
45-49	0.3528		0.5924		0.6779	
50-54	0.8169		0.4300		0.9581	
55-59	0.5716		0.1159		0.1817	
60-64	0.5742		0.5560		0.5160	
65 or Older	0.7123		0.9820		0.5244	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.15C Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	672	719	914	959	969	1,018
12	24	27	18	22	10	13
13	39	38	34	32	21	19
14	51	51	42	46	29	27
15	67	66	59	60	42	43
16	71	72	64	64	47	45
17	73	76	66	71	50	54
18	109	112	104	106	85	83
19	111	108	105	105	89	93
20	117	125	112	120	95	101
21	116	124	113	121	104	109
22	110	100	106	98	99	89
23	103	104	100	99	91	91
24	98	105	93	99	85	88
25	98	94	93	89	83	82
26-29	399	395	369	374	324	339
30-34	407	425	372	393	325	354
35-39	475	499	429	455	386	385
40-44	491	477	467	437	392	394
45-49	502	491	449	432	394	394
50-54	625	687	555	615	493	551
55-59	527	637	462	561	418	472
60-64	501	516	409	440	352	388
65 or Older	370	404	474	488	453	474

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.15D Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.29	0.37	0.39	0.39	0.41
12	0.58	0.62	0.45	0.53	0.26	0.31
13	0.85	0.81	0.77	0.72	0.49	0.44
14	0.97	1.02	0.87	0.97	0.65	0.63
15	1.08	1.09	1.02	1.08	0.83	0.88
16	1.02	1.09	1.04	1.10	0.92	0.89
17	1.05	1.02	1.05	1.11	0.94	1.04
18	1.06	1.08	1.16	1.22	1.23	1.19
19	1.03	1.13	1.17	1.22	1.29	1.36
20	0.99	0.97	1.18	1.15	1.36	1.36
21	0.78	0.85	0.97	1.00	1.19	1.23
22	0.82	0.89	0.95	1.06	1.16	1.32
23	0.74	0.88	0.93	1.01	1.16	1.21
24	0.94	0.90	1.13	1.04	1.25	1.21
25	0.80	0.75	1.09	1.01	1.30	1.24
26-29	0.74	0.77	1.00	0.96	1.16	1.19
30-34	0.73	0.72	0.90	0.92	1.04	1.08
35-39	0.70	0.78	1.09	1.02	1.25	1.11
40-44	0.71	0.74	1.09	1.02	1.17	1.15
45-49	0.60	0.64	0.96	0.96	1.12	1.09
50-54	1.02	1.15	1.52	1.51	1.57	1.58
55-59	1.23	1.22	1.60	1.78	1.64	1.80
60-64	1.22	1.28	1.83	2.05	1.89	2.10
65 or Older	1.04	1.11	1.33	1.34	1.27	1.31

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.15P Alcohol Use in Lifetime, Past Year, and Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3363		0.5599		0.6941	
12	0.5126		0.3622		0.4666	
13	0.0678		0.0633		0.1392	
14	0.0396		0.0392		0.0552	
15	0.5336		0.6008		0.8838	
16	0.6682		0.1299		0.0098	
17	0.5567		0.6835		0.1005	
18	0.0113		0.0012		0.0116	
19	0.9538		0.3865		0.4667	
20	0.1903		0.8407		0.6388	
21	0.7241		0.2285		0.3470	
22	0.1867		0.1717		0.0149	
23	0.1085		0.2497		0.4471	
24	0.7051		0.8920		0.6012	
25	0.6752		0.4947		0.3033	
26-29	0.2982		0.5126		0.8271	
30-34	0.6768		0.3878		0.2278	
35-39	0.1271		0.7888		1.0000	
40-44	0.8511		0.5826		0.0126	
45-49	0.1124		0.0623		0.6928	
50-54	0.3244		0.4055		0.6235	
55-59	0.4897		0.5829		0.6590	
60-64	0.8723		0.7275		0.8766	
65 or Older	0.2656		0.4982		0.8495	

*Low precision; no estimate reported.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.16C Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Numbers in Thousands, 2006 and 2007

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	969	1,018	746	767	427	429
12	10	13	6	8	*	1
13	21	19	13	13	6	2
14	29	27	22	17	7	4
15	42	43	33	32	11	18
16	47	45	40	35	19	19
17	50	54	41	44	23	25
18	85	83	70	71	48	35
19	89	93	72	79	45	42
20	95	101	75	81	38	48
21	104	109	85	84	52	46
22	99	89	78	73	43	40
23	91	91	74	71	42	44
24	85	88	67	70	40	40
25	83	82	66	64	39	35
26-29	324	339	240	253	132	131
30-34	325	354	223	235	115	120
35-39	386	385	260	257	129	140
40-44	392	394	256	250	126	130
45-49	394	394	256	237	135	141
50-54	493	551	300	333	161	194
55-59	418	472	198	236	122	119
60-64	352	388	185	210	75	98
65 or Older	453	474	241	248	137	101

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.16D Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: Standard Errors of Percentages, 2006 and 2007

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.39	0.41	0.30	0.31	0.17	0.17
12	0.26	0.31	0.15	0.20	*	0.03
13	0.49	0.44	0.32	0.30	0.15	0.04
14	0.65	0.63	0.50	0.42	0.16	0.11
15	0.83	0.88	0.68	0.68	0.24	0.40
16	0.92	0.89	0.85	0.74	0.44	0.42
17	0.94	1.04	0.85	0.92	0.53	0.59
18	1.23	1.19	1.15	1.16	0.93	0.65
19	1.29	1.36	1.25	1.34	0.91	0.87
20	1.36	1.36	1.33	1.35	0.89	0.96
21	1.19	1.23	1.31	1.31	1.00	0.99
22	1.16	1.32	1.31	1.32	0.99	0.96
23	1.16	1.21	1.22	1.25	0.93	1.00
24	1.25	1.21	1.28	1.29	0.93	0.94
25	1.30	1.24	1.31	1.28	0.93	0.86
26-29	1.16	1.19	1.15	1.19	0.77	0.74
30-34	1.04	1.08	0.98	1.00	0.59	0.61
35-39	1.25	1.11	1.06	0.99	0.59	0.62
40-44	1.17	1.15	1.00	0.98	0.54	0.59
45-49	1.12	1.09	1.00	0.89	0.60	0.59
50-54	1.57	1.58	1.29	1.36	0.76	0.86
55-59	1.64	1.80	1.06	1.18	0.69	0.63
60-64	1.89	2.10	1.27	1.44	0.55	0.72
65 or Older	1.27	1.31	0.68	0.68	0.39	0.28

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.16P Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month, by Detailed Age Category: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Age Category	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.6941		0.4697		0.9193	
12	0.4666		0.1988		*	
13	0.1392		0.6056		0.0173	
14	0.0552		0.0102		0.1407	
15	0.8838		0.5348		0.1297	
16	0.0098		0.0070		0.2221	
17	0.1005		0.1041		0.5789	
18	0.0116		0.0211		0.0046	
19	0.4667		0.3477		0.7794	
20	0.6388		0.5182		0.1540	
21	0.3470		0.6798		0.2497	
22	0.0149		0.2891		0.7908	
23	0.4471		0.2869		0.4382	
24	0.6012		0.5884		0.7119	
25	0.3033		0.3931		0.6285	
26-29	0.8271		0.8118		0.2034	
30-34	0.2278		0.1383		0.5165	
35-39	1.0000		0.5624		0.1930	
40-44	0.0126		0.0704		0.3954	
45-49	0.6928		0.9478		0.3415	
50-54	0.6235		0.7883		0.7243	
55-59	0.6590		0.1900		0.8615	
60-64	0.8766		0.7216		0.8010	
65 or Older	0.8495		0.9959		0.5966	

*Low precision; no estimate reported.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.