

Table 2.1C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	827	855	878	878	848	844
Cigarettes	863	879	840	848	804	808
Smokeless Tobacco	688	703	325	312	284	281
Cigars	872	872	468	484	352	337
Pipe Tobacco ²	740	735	--	--	166	170
ALCOHOL	672	719	914	959	969	1,018
Binge Alcohol Use ³	--	--	--	--	746	767
Heavy Alcohol Use ³	--	--	--	--	427	429

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.1D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.34	0.34	0.36	0.35	0.34	0.34
Cigarettes	0.35	0.35	0.34	0.34	0.33	0.33
Smokeless Tobacco	0.28	0.28	0.13	0.13	0.12	0.11
Cigars	0.35	0.35	0.19	0.20	0.14	0.14
Pipe Tobacco ²	0.30	0.30	--	--	0.07	0.07
ALCOHOL	0.27	0.29	0.37	0.39	0.39	0.41
Binge Alcohol Use ³	--	--	--	--	0.30	0.31
Heavy Alcohol Use ³	--	--	--	--	0.17	0.17

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.1P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0123		0.1309		0.0306	
Cigarettes	0.0438		0.1568		0.0749	
Smokeless Tobacco	0.2734		0.4303		0.5340	
Cigars	0.4074		0.4718		0.2695	
Pipe Tobacco ²	0.0018		--		0.2149	
ALCOHOL	0.3363		0.5599		0.6941	
Binge Alcohol Use ³	--		--		0.4697	
Heavy Alcohol Use ³	--		--		0.9193	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.2C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	37	41	28	28	17	19
Cigarettes	34	37	27	25	15	17
Smokeless Tobacco	17	18	12	13	7	8
Cigars	21	21	16	17	8	11
Pipe Tobacco ²	12	10	--	--	5	4
ALCOHOL	46	48	38	40	23	23
Binge Alcohol Use ³	--	--	--	--	15	15
Heavy Alcohol Use ³	--	--	--	--	6	2

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.2D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.43	0.47	0.34	0.34	0.21	0.23
Cigarettes	0.41	0.43	0.33	0.30	0.18	0.21
Smokeless Tobacco	0.21	0.22	0.14	0.16	0.08	0.10
Cigars	0.25	0.25	0.19	0.21	0.10	0.13
Pipe Tobacco ²	0.15	0.12	--	--	0.06	0.05
ALCOHOL	0.52	0.53	0.45	0.46	0.29	0.27
Binge Alcohol Use ³	--	--	--	--	0.18	0.18
Heavy Alcohol Use ³	--	--	--	--	0.08	0.03

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.2P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or 13: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.1070		0.1428		0.4819	
Cigarettes	0.0658		0.0192		0.7317	
Smokeless Tobacco	0.8946		0.5264		0.7253	
Cigars	0.7305		0.3881		0.1410	
Pipe Tobacco ²	0.3159		--		0.5761	
ALCOHOL	0.2159		0.2502		0.3351	
Binge Alcohol Use ³	--		--		0.8896	
Heavy Alcohol Use ³	--		--		0.0433	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.3C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	69	68	55	57	43	41
Cigarettes	64	62	51	49	37	36
Smokeless Tobacco	31	33	25	29	17	20
Cigars	46	44	37	37	24	23
Pipe Tobacco ²	19	18	--	--	11	12
ALCOHOL	84	84	71	76	50	50
Binge Alcohol Use ³	--	--	--	--	39	36
Heavy Alcohol Use ³	--	--	--	--	13	19

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.3D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.66	0.68	0.57	0.60	0.46	0.45
Cigarettes	0.64	0.65	0.54	0.53	0.40	0.40
Smokeless Tobacco	0.34	0.37	0.27	0.33	0.19	0.23
Cigars	0.49	0.48	0.40	0.41	0.27	0.26
Pipe Tobacco ²	0.22	0.21	--	--	0.13	0.14
ALCOHOL	0.73	0.77	0.66	0.74	0.52	0.55
Binge Alcohol Use ³	--	--	--	--	0.42	0.41
Heavy Alcohol Use ³	--	--	--	--	0.15	0.22

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.3P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 14 or 15: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0023		0.1198		0.1160	
Cigarettes	0.0013		0.0789		0.2284	
Smokeless Tobacco	0.5488		0.2929		0.6045	
Cigars	0.0199		0.3584		0.2130	
Pipe Tobacco ²	0.3178		--		0.6801	
ALCOHOL	0.0702		0.0793		0.2023	
Binge Alcohol Use ³	--		--		0.0534	
Heavy Alcohol Use ³	--		--		0.3916	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.4C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	93	94	79	80	62	63
Cigarettes	87	85	72	71	55	56
Smokeless Tobacco	43	50	37	41	25	29
Cigars	62	61	51	52	33	36
Pipe Tobacco ²	27	26	--	--	13	14
ALCOHOL	104	110	95	100	70	73
Binge Alcohol Use ³	--	--	--	--	58	57
Heavy Alcohol Use ³	--	--	--	--	30	32

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.4D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.76	0.78	0.73	0.74	0.62	0.64
Cigarettes	0.75	0.75	0.71	0.69	0.58	0.59
Smokeless Tobacco	0.46	0.54	0.41	0.45	0.29	0.32
Cigars	0.63	0.60	0.55	0.54	0.37	0.40
Pipe Tobacco ²	0.31	0.30	--	--	0.16	0.17
ALCOHOL	0.71	0.75	0.73	0.78	0.66	0.71
Binge Alcohol Use ³	--	--	--	--	0.60	0.60
Heavy Alcohol Use ³	--	--	--	--	0.35	0.36

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.4P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 16 or 17: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0736		0.2977		0.4151	
Cigarettes	0.0184		0.0989		0.2077	
Smokeless Tobacco	0.0414		0.3863		0.9486	
Cigars	0.7766		0.6593		0.2819	
Pipe Tobacco ²	0.6439		--		0.7922	
ALCOHOL	0.9805		0.3794		0.4468	
Binge Alcohol Use ³	--		--		0.4173	
Heavy Alcohol Use ³	--		--		0.6849	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.5C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	184	206	161	172	134	134
Cigarettes	167	190	139	153	120	121
Smokeless Tobacco	87	93	63	70	45	47
Cigars	135	149	110	114	70	78
Pipe Tobacco ²	52	52	--	--	22	24
ALCOHOL	230	250	216	242	180	202
Binge Alcohol Use ³	--	--	--	--	142	163
Heavy Alcohol Use ³	--	--	--	--	86	83

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.5D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.71	0.75	0.75	0.77	0.75	0.72
Cigarettes	0.75	0.75	0.75	0.75	0.74	0.70
Smokeless Tobacco	0.56	0.57	0.44	0.46	0.33	0.33
Cigars	0.70	0.74	0.65	0.64	0.48	0.52
Pipe Tobacco ²	0.37	0.36	--	--	0.17	0.18
ALCOHOL	0.63	0.67	0.71	0.76	0.78	0.84
Binge Alcohol Use ³	--	--	--	--	0.76	0.80
Heavy Alcohol Use ³	--	--	--	--	0.58	0.51

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.5P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 20: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0214		0.0198		0.0218	
Cigarettes	0.0222		0.0179		0.0032	
Smokeless Tobacco	0.7341		0.6205		0.5194	
Cigars	0.1496		0.3951		0.7504	
Pipe Tobacco ²	0.7465		--		0.5804	
ALCOHOL	0.0291		0.1079		0.3849	
Binge Alcohol Use ³	--		--		0.6598	
Heavy Alcohol Use ³	--		--		0.3072	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.6C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	253	244	206	199	176	173
Cigarettes	236	229	185	183	163	159
Smokeless Tobacco	115	107	63	63	50	47
Cigars	187	181	122	117	79	76
Pipe Tobacco ²	70	68	--	--	25	22
ALCOHOL	297	298	285	285	260	251
Binge Alcohol Use ³	--	--	--	--	195	192
Heavy Alcohol Use ³	--	--	--	--	109	100

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.6D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.56	0.54	0.62	0.61	0.59	0.62
Cigarettes	0.55	0.56	0.59	0.62	0.58	0.61
Smokeless Tobacco	0.50	0.45	0.31	0.31	0.24	0.23
Cigars	0.62	0.61	0.48	0.48	0.35	0.34
Pipe Tobacco ²	0.33	0.32	--	--	0.12	0.11
ALCOHOL	0.39	0.41	0.49	0.50	0.61	0.59
Binge Alcohol Use ³	--	--	--	--	0.62	0.60
Heavy Alcohol Use ³	--	--	--	--	0.47	0.43

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.6P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 21 to 25: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.2698		0.2013		0.0323	
Cigarettes	0.1080		0.0817		0.0343	
Smokeless Tobacco	0.9426		0.7353		0.6724	
Cigars	0.7280		0.3397		0.1470	
Pipe Tobacco ²	0.5727		--		0.0885	
ALCOHOL	0.2478		0.7415		0.7695	
Binge Alcohol Use ³	--		--		0.8422	
Heavy Alcohol Use ³	--		--		0.1795	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.7C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	260	268	274	290	269	291
Cigarettes	270	283	270	289	257	279
Smokeless Tobacco	247	239	126	129	107	112
Cigars	292	287	204	203	159	143
Pipe Tobacco ²	167	143	--	--	56	43
ALCOHOL	184	197	242	244	284	294
Binge Alcohol Use ³	--	--	--	--	272	285
Heavy Alcohol Use ³	--	--	--	--	172	178

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.7D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.74	0.76	0.78	0.82	0.77	0.82
Cigarettes	0.77	0.80	0.77	0.82	0.73	0.79
Smokeless Tobacco	0.70	0.68	0.36	0.37	0.30	0.32
Cigars	0.83	0.81	0.58	0.57	0.45	0.40
Pipe Tobacco ²	0.48	0.41	--	--	0.16	0.12
ALCOHOL	0.52	0.56	0.69	0.69	0.81	0.83
Binge Alcohol Use ³	--	--	--	--	0.77	0.81
Heavy Alcohol Use ³	--	--	--	--	0.49	0.50

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.7P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 to 34: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.1092		0.2604		0.3001	
Cigarettes	0.0873		0.3653		0.5449	
Smokeless Tobacco	0.0388		0.4179		0.5010	
Cigars	0.2863		0.1306		0.1065	
Pipe Tobacco ²	0.2643		--		0.4329	
ALCOHOL	0.6909		0.2984		0.4951	
Binge Alcohol Use ³	--		--		0.4255	
Heavy Alcohol Use ³	--		--		0.6039	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.8C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	714	754	748	750	723	719
Cigarettes	749	780	705	723	683	687
Smokeless Tobacco	615	644	275	249	250	236
Cigars	774	774	378	393	288	282
Pipe Tobacco ²	683	693	--	--	149	159
ALCOHOL	583	647	827	901	873	935
Binge Alcohol Use ³	--	--	--	--	628	641
Heavy Alcohol Use ³	--	--	--	--	346	357

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.8D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.47	0.49	0.49	0.49	0.47	0.47
Cigarettes	0.49	0.50	0.46	0.47	0.45	0.44
Smokeless Tobacco	0.40	0.42	0.18	0.16	0.16	0.15
Cigars	0.51	0.50	0.25	0.25	0.19	0.18
Pipe Tobacco ²	0.45	0.45	--	--	0.10	0.10
ALCOHOL	0.38	0.42	0.54	0.58	0.57	0.60
Binge Alcohol Use ³	--	--	--	--	0.41	0.41
Heavy Alcohol Use ³	--	--	--	--	0.23	0.23

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.8P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 35 or Older: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.1574		0.6333		0.2168	
Cigarettes	0.4879		0.7794		0.3842	
Smokeless Tobacco	0.5567		0.3311		0.6175	
Cigars	0.8394		0.6934		0.8034	
Pipe Tobacco ²	0.0020		--		0.3181	
ALCOHOL	0.7256		0.6327		0.6337	
Binge Alcohol Use ³	--		--		0.3862	
Heavy Alcohol Use ³	--		--		0.4664	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.9C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	446	510	596	620	584	595
Cigarettes	528	560	559	605	543	574
Smokeless Tobacco	581	599	309	282	273	251
Cigars	618	628	426	429	319	311
Pipe Tobacco ²	646	634	--	--	160	168
ALCOHOL	352	401	569	604	611	667
Binge Alcohol Use ³	--	--	--	--	559	585
Heavy Alcohol Use ³	--	--	--	--	355	375

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.9D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.37	0.42	0.50	0.52	0.49	0.50
Cigarettes	0.44	0.47	0.47	0.50	0.45	0.48
Smokeless Tobacco	0.49	0.50	0.26	0.23	0.23	0.21
Cigars	0.52	0.52	0.36	0.36	0.27	0.26
Pipe Tobacco ²	0.54	0.53	--	--	0.13	0.14
ALCOHOL	0.30	0.33	0.48	0.50	0.51	0.55
Binge Alcohol Use ³	--	--	--	--	0.47	0.49
Heavy Alcohol Use ³	--	--	--	--	0.30	0.31

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.9P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Males: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0377		0.2824		0.0928	
Cigarettes	0.2850		0.3268		0.2536	
Smokeless Tobacco	0.4281		0.2962		0.3864	
Cigars	0.5834		0.9011		0.6016	
Pipe Tobacco ²	0.0009		--		0.2984	
ALCOHOL	0.2526		0.6997		0.6120	
Binge Alcohol Use ³	--		--		0.5079	
Heavy Alcohol Use ³	--		--		0.8720	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.10C Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	614	629	543	545	511	511
Cigarettes	616	639	529	531	502	493
Smokeless Tobacco	275	280	88	119	76	113
Cigars	466	461	194	185	131	116
Pipe Tobacco ²	232	246	--	--	47	30
ALCOHOL	524	558	630	671	644	674
Binge Alcohol Use ³	--	--	--	--	399	394
Heavy Alcohol Use ³	--	--	--	--	168	171

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.10D Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: Standard Errors of Percentages, 2006 and 2007

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.49	0.49	0.43	0.43	0.40	0.40
Cigarettes	0.49	0.50	0.42	0.42	0.40	0.39
Smokeless Tobacco	0.22	0.22	0.07	0.09	0.06	0.09
Cigars	0.37	0.36	0.15	0.14	0.10	0.09
Pipe Tobacco ²	0.18	0.19	--	--	0.04	0.02
ALCOHOL	0.41	0.44	0.50	0.53	0.51	0.53
Binge Alcohol Use ³	--	--	--	--	0.32	0.31
Heavy Alcohol Use ³	--	--	--	--	0.13	0.13

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.10P Tobacco Product and Alcohol Use in Lifetime, Past Year, and Past Month among Females: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS¹	0.0820		0.2208		0.1053	
Cigarettes	0.0638		0.2417		0.1099	
Smokeless Tobacco	0.2884		0.5733		0.5855	
Cigars	0.3789		0.1328		0.0798	
Pipe Tobacco ²	0.3635		--		0.3523	
ALCOHOL	0.6821		0.6293		0.2724	
Binge Alcohol Use ³	--		--		0.6754	
Heavy Alcohol Use ³	--		--		0.9213	

*Low precision; no estimate reported.

-- Not available.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

² Information about past year use of pipe tobacco was not collected.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.