

Table 2.47C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	828	856	880	876	850	844
GEOGRAPHIC DIVISION						
Northeast	355	359	347	337	322	330
New England	180	175	165	160	163	145
Middle Atlantic	304	312	306	294	277	296
Midwest	299	287	367	367	343	352
East North Central	255	245	285	297	273	281
West North Central	152	152	225	206	206	203
South	489	542	556	564	550	532
South Atlantic	368	422	434	431	430	420
East South Central	168	163	202	197	205	190
West South Central	267	286	276	290	276	268
West	431	468	431	437	403	407
Mountain	163	200	178	222	171	215
Pacific	397	423	393	377	366	347
COUNTY TYPE						
Large Metro	1,507	1,487	857	871	767	783
Small Metro	1,045	1,167	625	666	583	597
250K – 1 Mil. Pop.	816	992	472	588	429	534
< 250K Pop.	703	702	406	380	383	338
Nonmetro	880	841	533	504	497	470
Urbanized	551	577	314	350	289	317
Less Urbanized	733	649	439	377	411	352
Completely Rural	351	358	219	209	205	190

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.47D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.34	0.35	0.36	0.35	0.35	0.34
GEOGRAPHIC DIVISION						
Northeast	0.77	0.78	0.76	0.73	0.70	0.72
New England	1.50	1.45	1.37	1.33	1.35	1.20
Middle Atlantic	0.90	0.92	0.90	0.87	0.82	0.87
Midwest	0.55	0.52	0.67	0.67	0.63	0.64
East North Central	0.67	0.64	0.74	0.78	0.71	0.73
West North Central	0.93	0.92	1.37	1.25	1.25	1.23
South	0.55	0.60	0.62	0.63	0.62	0.59
South Atlantic	0.78	0.89	0.92	0.91	0.91	0.88
East South Central	1.15	1.11	1.38	1.34	1.40	1.29
West South Central	0.98	1.03	1.01	1.05	1.01	0.97
West	0.76	0.82	0.76	0.76	0.71	0.71
Mountain	0.96	1.16	1.05	1.29	1.02	1.24
Pacific	1.00	1.06	0.99	0.94	0.92	0.87
COUNTY TYPE						
Large Metro	0.51	0.50	0.49	0.50	0.47	0.48
Small Metro	0.56	0.62	0.65	0.63	0.62	0.59
250K – 1 Mil. Pop.	0.69	0.77	0.77	0.78	0.72	0.74
< 250K Pop.	0.93	1.03	1.12	1.13	1.11	1.05
Nonmetro	0.66	0.69	0.84	0.79	0.83	0.78
Urbanized	1.15	1.01	1.27	1.13	1.30	1.14
Less Urbanized	0.87	0.96	1.18	1.27	1.14	1.22
Completely Rural	2.14	2.41	2.36	2.58	2.37	2.38

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.47P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0150		0.1660		0.0407	
GEOGRAPHIC DIVISION						
Northeast	0.1098		0.4037		0.3445	
New England	0.4373		0.9141		0.5479	
Middle Atlantic	0.1600		0.3042		0.4504	
Midwest	0.2919		0.9325		0.8023	
East North Central	0.2276		0.5356		0.7634	
West North Central	0.9840		0.4494		0.9971	
South	0.1032		0.1014		0.0182	
South Atlantic	0.5082		0.6214		0.8365	
East South Central	0.0154		0.0066		0.0091	
West South Central	0.5161		0.0295		0.0011	
West	0.0322		0.8933		0.9216	
Mountain	0.2503		0.7125		0.7464	
Pacific	0.0634		0.7261		0.7428	
COUNTY TYPE						
Large Metro	0.0184		0.5955		0.2209	
Small Metro	0.7668		0.1752		0.1448	
250K – 1 Mil. Pop.	0.9296		0.2136		0.2287	
< 250K Pop.	0.7412		0.5807		0.4305	
Nonmetro	0.1815		0.4766		0.3295	
Urbanized	0.7819		0.1541		0.2677	
Less Urbanized	0.1374		0.2979		0.2883	
Completely Rural	0.2620		0.0241		0.0089	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.48C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	106	109	90	95	72	76
GEOGRAPHIC DIVISION						
Northeast	38	40	37	35	32	28
New England	18	20	19	18	16	15
Middle Atlantic	34	34	31	30	27	23
Midwest	42	42	35	38	30	32
East North Central	34	35	28	31	24	26
West North Central	25	23	22	21	19	17
South	68	68	58	61	46	51
South Atlantic	49	48	41	45	32	39
East South Central	26	26	22	21	19	17
West South Central	35	39	33	35	25	28
West	57	56	49	49	39	39
Mountain	28	26	25	24	19	19
Pacific	50	49	42	42	34	34
COUNTY TYPE						
Large Metro	100	104	79	82	60	58
Small Metro	75	78	61	63	45	46
250K – 1 Mil. Pop.	59	61	48	47	34	35
< 250K Pop.	48	50	38	41	30	29
Nonmetro	58	59	45	49	34	39
Urbanized	37	40	29	31	20	25
Less Urbanized	45	43	35	37	27	30
Completely Rural	21	18	14	14	10	10

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.48D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.42	0.43	0.36	0.38	0.29	0.30
GEOGRAPHIC DIVISION						
Northeast	0.85	0.89	0.81	0.78	0.71	0.62
New England	1.56	1.71	1.63	1.53	1.40	1.26
Middle Atlantic	1.01	1.03	0.93	0.90	0.82	0.71
Midwest	0.75	0.75	0.62	0.68	0.53	0.57
East North Central	0.85	0.88	0.71	0.80	0.60	0.67
West North Central	1.45	1.39	1.28	1.25	1.10	1.03
South	0.74	0.74	0.64	0.67	0.51	0.56
South Atlantic	1.06	1.04	0.88	0.96	0.69	0.83
East South Central	1.77	1.73	1.51	1.41	1.26	1.14
West South Central	1.17	1.31	1.09	1.15	0.83	0.93
West	0.95	0.92	0.80	0.81	0.64	0.65
Mountain	1.58	1.44	1.38	1.30	1.08	1.02
Pacific	1.16	1.15	0.98	1.00	0.79	0.81
COUNTY TYPE						
Large Metro	0.59	0.56	0.50	0.49	0.40	0.38
Small Metro	0.74	0.84	0.65	0.72	0.52	0.57
250K – 1 Mil. Pop.	0.91	1.01	0.79	0.83	0.60	0.67
< 250K Pop.	1.22	1.40	1.06	1.28	0.98	0.99
Nonmetro	0.89	0.97	0.80	0.90	0.68	0.79
Urbanized	1.40	1.38	1.29	1.26	1.00	1.15
Less Urbanized	1.38	1.37	1.20	1.32	1.07	1.19
Completely Rural	2.55	3.10	2.35	2.64	1.93	2.07

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.48P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0011		0.0538		0.2320	
GEOGRAPHIC DIVISION						
Northeast	0.0024		0.0021		0.0408	
New England	0.5512		0.3310		0.3150	
Middle Atlantic	0.0013		0.0024		0.0733	
Midwest	0.7433		0.3391		0.8813	
East North Central	0.4429		0.0950		0.1906	
West North Central	0.0881		0.4675		0.1247	
South	0.1045		0.5259		0.5171	
South Atlantic	0.5119		0.8045		0.1209	
East South Central	0.1387		0.1941		0.4182	
West South Central	0.2909		0.5578		0.7793	
West	0.0476		0.1594		0.1198	
Mountain	0.2109		0.2242		0.5541	
Pacific	0.1067		0.3435		0.1432	
COUNTY TYPE						
Large Metro	0.0011		0.0212		0.0134	
Small Metro	0.3766		0.4004		0.4883	
250K – 1 Mil. Pop.	0.7235		0.5223		0.8386	
< 250K Pop.	0.2677		0.5584		0.3782	
Nonmetro	0.2844		0.5856		0.0287	
Urbanized	0.3853		0.2244		0.0237	
Less Urbanized	0.1318		0.9790		0.1680	
Completely Rural	0.1220		0.4622		0.4765	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.49C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	153	149	162	159	160	162
GEOGRAPHIC DIVISION						
Northeast	65	65	68	71	70	72
New England	26	31	30	33	30	33
Middle Atlantic	60	56	61	62	63	63
Midwest	56	53	62	60	64	62
East North Central	45	40	50	45	52	45
West North Central	34	35	38	41	38	43
South	87	93	93	101	95	101
South Atlantic	64	64	69	69	70	70
East South Central	31	34	33	39	38	40
West South Central	48	57	53	61	54	58
West	88	89	96	84	88	88
Mountain	36	47	41	44	38	43
Pacific	80	76	86	72	78	76
COUNTY TYPE						
Large Metro	277	267	219	215	183	182
Small Metro	195	206	161	171	138	140
250K – 1 Mil. Pop.	158	163	128	135	108	111
< 250K Pop.	118	136	99	110	86	89
Nonmetro	135	143	115	115	102	95
Urbanized	102	91	83	73	70	61
Less Urbanized	101	110	87	88	78	71
Completely Rural	39	42	37	38	35	33

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.49D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.47	0.45	0.49	0.49	0.49	0.50
GEOGRAPHIC DIVISION						
Northeast	1.11	1.10	1.17	1.19	1.19	1.21
New England	1.73	2.05	1.94	2.15	1.97	2.15
Middle Atlantic	1.37	1.29	1.41	1.41	1.44	1.45
Midwest	0.76	0.73	0.84	0.83	0.86	0.85
East North Central	0.88	0.79	0.98	0.90	1.01	0.89
West North Central	1.47	1.56	1.66	1.80	1.66	1.89
South	0.75	0.80	0.80	0.86	0.82	0.86
South Atlantic	1.09	1.09	1.18	1.17	1.18	1.18
East South Central	1.65	1.80	1.73	2.07	2.02	2.16
West South Central	1.22	1.46	1.36	1.57	1.38	1.49
West	1.12	1.14	1.22	1.07	1.12	1.12
Mountain	1.54	1.98	1.74	1.87	1.61	1.84
Pacific	1.46	1.39	1.57	1.31	1.43	1.39
COUNTY TYPE						
Large Metro	0.71	0.67	0.73	0.70	0.71	0.72
Small Metro	0.76	0.86	0.80	0.90	0.84	0.88
250K – 1 Mil. Pop.	0.91	1.07	0.97	1.11	0.99	1.07
< 250K Pop.	1.43	1.37	1.42	1.47	1.45	1.47
Nonmetro	0.90	0.98	1.09	1.10	1.16	1.25
Urbanized	1.39	1.66	1.66	1.75	1.85	1.97
Less Urbanized	1.33	1.20	1.44	1.49	1.48	1.78
Completely Rural	2.72	2.41	3.47	3.92	3.58	3.94

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.49P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0138		0.0166		0.0019	
GEOGRAPHIC DIVISION						
Northeast	0.1090		0.0198		0.0079	
New England	0.7522		0.2613		0.0823	
Middle Atlantic	0.1034		0.0387		0.0334	
Midwest	0.3995		0.1480		0.0236	
East North Central	0.4014		0.2925		0.1405	
West North Central	0.7798		0.3278		0.0655	
South	0.0874		0.1302		0.0923	
South Atlantic	0.5265		0.7505		0.3188	
East South Central	0.0200		0.0795		0.2590	
West South Central	0.4961		0.2888		0.3654	
West	0.5093		0.8898		0.9596	
Mountain	0.4308		0.9494		0.9560	
Pacific	0.7335		0.8482		0.9372	
COUNTY TYPE						
Large Metro	0.0770		0.1065		0.0371	
Small Metro	0.2240		0.2089		0.1857	
250K – 1 Mil. Pop.	0.1092		0.2438		0.3771	
< 250K Pop.	0.8561		0.5501		0.2708	
Nonmetro	0.1550		0.1418		0.0257	
Urbanized	0.1609		0.4964		0.3979	
Less Urbanized	0.2484		0.1209		0.0400	
Completely Rural	0.3251		0.8754		0.5506	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.50C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	762	816	826	827	799	801
GEOGRAPHIC DIVISION						
Northeast	322	347	325	321	298	313
New England	157	171	148	153	148	137
Middle Atlantic	280	301	289	281	258	281
Midwest	278	269	338	349	322	337
East North Central	235	234	265	284	259	270
West North Central	145	140	207	195	193	192
South	459	512	530	536	527	508
South Atlantic	348	401	414	410	413	402
East South Central	161	154	199	187	196	182
West South Central	245	266	262	273	268	253
West	401	443	417	415	387	390
Mountain	146	183	171	208	161	201
Pacific	369	403	379	359	351	336
COUNTY TYPE						
Large Metro	1,418	1,408	773	788	701	715
Small Metro	965	1,088	566	595	535	540
250K – 1 Mil. Pop.	754	921	424	522	393	482
< 250K Pop.	639	650	363	340	347	307
Nonmetro	804	775	478	439	455	417
Urbanized	493	516	267	308	255	281
Less Urbanized	654	597	383	317	367	306
Completely Rural	319	318	196	176	185	164

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.50D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.41	0.43	0.44	0.44	0.43	0.42
GEOGRAPHIC DIVISION						
Northeast	0.91	0.98	0.92	0.90	0.84	0.88
New England	1.68	1.83	1.58	1.64	1.59	1.47
Middle Atlantic	1.07	1.15	1.11	1.07	0.99	1.08
Midwest	0.67	0.64	0.81	0.83	0.77	0.80
East North Central	0.80	0.80	0.91	0.97	0.89	0.92
West North Central	1.16	1.12	1.66	1.55	1.55	1.53
South	0.67	0.74	0.78	0.78	0.77	0.74
South Atlantic	0.95	1.08	1.13	1.11	1.13	1.09
East South Central	1.43	1.36	1.77	1.64	1.75	1.60
West South Central	1.20	1.29	1.29	1.32	1.31	1.22
West	0.94	1.02	0.98	0.96	0.91	0.90
Mountain	1.14	1.39	1.34	1.58	1.27	1.53
Pacific	1.23	1.33	1.27	1.19	1.17	1.11
COUNTY TYPE						
Large Metro	0.61	0.63	0.60	0.62	0.57	0.59
Small Metro	0.69	0.74	0.83	0.78	0.80	0.74
250K – 1 Mil. Pop.	0.84	0.91	0.98	0.97	0.92	0.93
< 250K Pop.	1.16	1.27	1.42	1.40	1.41	1.33
Nonmetro	0.82	0.86	1.04	0.97	1.02	0.96
Urbanized	1.43	1.28	1.55	1.45	1.58	1.42
Less Urbanized	1.07	1.17	1.47	1.51	1.42	1.50
Completely Rural	2.68	2.98	3.01	3.05	2.99	2.87

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.50P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0723		0.4554		0.1573	
GEOGRAPHIC DIVISION						
Northeast	0.2788		0.9129		0.8309	
New England	0.4761		0.6638		0.8809	
Middle Atlantic	0.3907		0.7106		0.8704	
Midwest	0.2188		0.9201		0.8684	
East North Central	0.2088		0.5347		0.8830	
West North Central	0.7740		0.2728		0.5939	
South	0.2065		0.1799		0.0288	
South Atlantic	0.5750		0.5910		0.8017	
East South Central	0.0726		0.0264		0.0192	
West South Central	0.6484		0.0476		0.0017	
West	0.0562		0.7690		0.9488	
Mountain	0.3501		0.8514		0.6878	
Pacific	0.0881		0.6734		0.8602	
COUNTY TYPE						
Large Metro	0.0928		0.9155		0.5634	
Small Metro	0.6650		0.2908		0.2197	
250K – 1 Mil. Pop.	0.8325		0.3442		0.2816	
< 250K Pop.	0.6814		0.6370		0.5593	
Nonmetro	0.4172		0.5480		0.4049	
Urbanized	0.6633		0.1690		0.2935	
Less Urbanized	0.4075		0.3552		0.3920	
Completely Rural	0.2495		0.0430		0.0219	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.51C (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	793	830	868	868	838	838
GEOGRAPHIC DIVISION						
Northeast	344	350	345	334	319	325
New England	167	171	160	158	159	140
Middle Atlantic	299	304	305	293	276	294
Midwest	288	276	362	365	341	349
East North Central	244	239	280	296	272	281
West North Central	149	145	222	205	202	199
South	470	524	552	558	544	527
South Atlantic	356	407	430	428	426	416
East South Central	165	163	203	197	201	191
West South Central	252	275	271	285	273	264
West	416	452	430	433	401	407
Mountain	154	189	175	218	170	212
Pacific	382	410	392	375	364	349
COUNTY TYPE						
Large Metro	1,484	1,461	841	858	757	777
Small Metro	1,016	1,138	608	646	571	584
250K – 1 Mil. Pop.	796	970	461	571	421	525
< 250K Pop.	680	679	393	366	372	328
Nonmetro	855	820	518	489	488	458
Urbanized	530	561	302	340	282	309
Less Urbanized	711	630	423	361	402	340
Completely Rural	339	346	213	201	200	185

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.51D (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.36	0.37	0.39	0.39	0.38	0.38
GEOGRAPHIC DIVISION						
Northeast	0.83	0.85	0.83	0.81	0.77	0.79
New England	1.54	1.57	1.47	1.46	1.46	1.29
Middle Atlantic	0.98	1.00	1.00	0.96	0.90	0.96
Midwest	0.59	0.56	0.74	0.74	0.69	0.71
East North Central	0.71	0.70	0.82	0.86	0.79	0.82
West North Central	1.01	0.98	1.51	1.38	1.37	1.34
South	0.59	0.65	0.69	0.69	0.68	0.65
South Atlantic	0.84	0.95	1.01	1.00	1.01	0.97
East South Central	1.25	1.23	1.55	1.49	1.53	1.44
West South Central	1.04	1.12	1.12	1.16	1.12	1.07
West	0.83	0.88	0.85	0.85	0.80	0.80
Mountain	1.02	1.22	1.16	1.41	1.12	1.37
Pacific	1.08	1.15	1.11	1.05	1.03	0.98
COUNTY TYPE						
Large Metro	0.55	0.55	0.54	0.55	0.52	0.54
Small Metro	0.59	0.65	0.72	0.69	0.69	0.65
250K – 1 Mil. Pop.	0.72	0.79	0.85	0.85	0.79	0.81
< 250K Pop.	1.01	1.12	1.24	1.26	1.23	1.17
Nonmetro	0.73	0.76	0.94	0.89	0.92	0.87
Urbanized	1.25	1.10	1.41	1.26	1.44	1.26
Less Urbanized	0.96	1.03	1.31	1.40	1.26	1.35
Completely Rural	2.37	2.73	2.67	2.93	2.65	2.75

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.51P (REVISED) Tobacco Product Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0267		0.2212		0.0448	
GEOGRAPHIC DIVISION						
Northeast	0.1730		0.5835		0.4379	
New England	0.4458		0.8323		0.6138	
Middle Atlantic	0.2504		0.4504		0.5395	
Midwest	0.2933		0.8358		0.7685	
East North Central	0.2810		0.4164		0.6625	
West North Central	0.8183		0.4076		0.8911	
South	0.1231		0.1110		0.0136	
South Atlantic	0.5125		0.6501		0.9546	
East South Central	0.0253		0.0100		0.0106	
West South Central	0.5708		0.0326		0.0011	
West	0.0431		0.7704		0.9733	
Mountain	0.2868		0.8157		0.7146	
Pacific	0.0773		0.6523		0.8454	
COUNTY TYPE						
Large Metro	0.0467		0.7861		0.3141	
Small Metro	0.5274		0.1631		0.1182	
250K – 1 Mil. Pop.	0.6299		0.1872		0.1726	
< 250K Pop.	0.7198		0.5907		0.4420	
Nonmetro	0.3000		0.4747		0.2570	
Urbanized	0.8605		0.1944		0.3620	
Less Urbanized	0.3193		0.3401		0.2755	
Completely Rural	0.2899		0.0293		0.0109	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.