

Table 2.57C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	688	706	324	313	284	283
GEOGRAPHIC DIVISION						
Northeast	250	290	84	95	67	83
New England	122	147	35	55	28	49
Middle Atlantic	220	250	76	77	62	66
Midwest	282	259	134	144	116	129
East North Central	217	222	98	108	86	98
West North Central	176	145	86	90	75	79
South	455	455	254	207	224	191
South Atlantic	345	365	185	152	168	140
East South Central	188	176	97	84	88	79
West South Central	221	205	135	110	115	100
West	331	378	136	160	114	138
Mountain	159	168	68	86	58	77
Pacific	297	329	118	133	99	114
COUNTY TYPE						
Large Metro	601	589	202	208	166	190
Small Metro	424	458	178	196	157	166
250K – 1 Mil. Pop.	312	326	131	144	123	123
< 250K Pop.	285	332	117	128	96	107
Nonmetro	432	434	212	176	192	161
Urbanized	257	292	108	117	100	107
Less Urbanized	354	306	176	123	156	109
Completely Rural	158	160	91	63	83	61

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.57D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.28	0.28	0.13	0.13	0.12	0.11
GEOGRAPHIC DIVISION						
Northeast	0.55	0.63	0.18	0.21	0.15	0.18
New England	1.02	1.23	0.29	0.46	0.23	0.41
Middle Atlantic	0.65	0.74	0.22	0.23	0.18	0.20
Midwest	0.52	0.47	0.24	0.26	0.21	0.24
East North Central	0.57	0.58	0.26	0.28	0.22	0.26
West North Central	1.07	0.88	0.52	0.54	0.46	0.48
South	0.51	0.51	0.29	0.23	0.25	0.21
South Atlantic	0.73	0.77	0.39	0.32	0.36	0.29
East South Central	1.29	1.19	0.67	0.57	0.60	0.53
West South Central	0.81	0.74	0.49	0.40	0.42	0.36
West	0.59	0.66	0.24	0.28	0.20	0.24
Mountain	0.94	0.97	0.40	0.50	0.35	0.45
Pacific	0.75	0.82	0.30	0.33	0.25	0.28
COUNTY TYPE						
Large Metro	0.39	0.39	0.15	0.15	0.12	0.14
Small Metro	0.49	0.49	0.24	0.24	0.22	0.21
250K – 1 Mil. Pop.	0.57	0.54	0.28	0.26	0.26	0.23
< 250K Pop.	0.90	0.95	0.44	0.49	0.36	0.41
Nonmetro	0.70	0.74	0.45	0.38	0.42	0.36
Urbanized	1.10	1.16	0.57	0.61	0.55	0.57
Less Urbanized	1.03	0.99	0.71	0.55	0.65	0.50
Completely Rural	2.24	2.72	1.67	1.32	1.58	1.30

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.57P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.2745		0.4459		0.5581	
GEOGRAPHIC DIVISION						
Northeast	0.7352		0.9248		0.7094	
New England	0.7503		0.5440		0.6416	
Middle Atlantic	0.8280		0.7904		0.8837	
Midwest	0.3567		0.3602		0.5060	
East North Central	0.2376		0.4127		0.3638	
West North Central	0.9740		0.6305		0.9929	
South	0.0406		0.0880		0.1097	
South Atlantic	0.5300		0.4669		0.3041	
East South Central	0.1867		0.2533		0.2776	
West South Central	0.0231		0.1797		0.4463	
West	0.9657		0.9234		0.6755	
Mountain	0.9123		0.6577		0.8119	
Pacific	0.9359		0.7161		0.7500	
COUNTY TYPE						
Large Metro	0.3279		0.7041		0.6503	
Small Metro	0.4171		0.7463		0.4035	
250K – 1 Mil. Pop.	0.0635		0.6422		0.4243	
< 250K Pop.	0.3410		0.8913		0.7983	
Nonmetro	0.8722		0.5655		0.5033	
Urbanized	0.2635		0.1456		0.1837	
Less Urbanized	0.7303		0.2887		0.2783	
Completely Rural	0.4756		0.0871		0.0928	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.58C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	52	62	44	53	31	36
GEOGRAPHIC DIVISION						
Northeast	21	19	18	16	11	12
New England	8	10	8	8	6	6
Middle Atlantic	19	16	16	14	10	11
Midwest	24	27	18	22	13	14
East North Central	17	22	14	18	10	11
West North Central	17	15	12	12	9	9
South	37	42	30	37	21	27
South Atlantic	21	33	18	27	12	21
East South Central	17	17	15	15	11	10
West South Central	23	22	19	19	14	13
West	29	29	25	23	16	13
Mountain	15	15	14	11	7	6
Pacific	24	25	20	21	15	12
COUNTY TYPE						
Large Metro	40	42	31	35	19	23
Small Metro	33	39	27	30	18	18
250K – 1 Mil. Pop.	23	28	19	22	14	12
< 250K Pop.	25	27	19	20	12	13
Nonmetro	30	37	24	31	19	23
Urbanized	17	23	15	18	12	13
Less Urbanized	24	28	18	23	15	18
Completely Rural	11	12	7	9	5	7

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.58D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.21	0.24	0.17	0.21	0.12	0.14
GEOGRAPHIC DIVISION						
Northeast	0.46	0.42	0.40	0.36	0.25	0.27
New England	0.70	0.86	0.68	0.72	0.50	0.48
Middle Atlantic	0.57	0.48	0.48	0.42	0.29	0.32
Midwest	0.42	0.47	0.32	0.39	0.23	0.25
East North Central	0.43	0.55	0.35	0.46	0.24	0.28
West North Central	0.99	0.88	0.69	0.70	0.51	0.53
South	0.40	0.46	0.32	0.40	0.23	0.30
South Atlantic	0.45	0.71	0.38	0.59	0.25	0.45
East South Central	1.17	1.11	0.98	1.03	0.77	0.71
West South Central	0.78	0.73	0.62	0.63	0.46	0.45
West	0.48	0.48	0.42	0.39	0.27	0.22
Mountain	0.86	0.85	0.81	0.58	0.41	0.33
Pacific	0.57	0.58	0.48	0.49	0.34	0.28
COUNTY TYPE						
Large Metro	0.27	0.28	0.22	0.24	0.14	0.16
Small Metro	0.42	0.47	0.34	0.37	0.24	0.23
250K – 1 Mil. Pop.	0.43	0.51	0.36	0.41	0.27	0.25
< 250K Pop.	0.92	0.97	0.72	0.76	0.46	0.49
Nonmetro	0.63	0.75	0.53	0.66	0.44	0.52
Urbanized	0.92	1.08	0.83	0.94	0.68	0.70
Less Urbanized	1.00	1.16	0.81	1.03	0.68	0.83
Completely Rural	1.83	2.39	1.32	2.00	1.10	1.50

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.58P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0482		0.1521		0.6417	
GEOGRAPHIC DIVISION						
Northeast	0.7098		0.4005		1.0000	
New England	0.6848		0.5174		0.5432	
Middle Atlantic	0.5020		0.5631		0.7366	
Midwest	0.0140		0.0279		0.3432	
East North Central	0.0036		0.0147		0.1565	
West North Central	0.8047		0.6827		0.8753	
South	0.1553		0.0776		0.4244	
South Atlantic	0.0019		0.0035		0.0340	
East South Central	0.6451		0.8807		0.6473	
West South Central	0.4279		0.5963		0.4014	
West	0.9802		0.3818		0.2176	
Mountain	0.5344		0.1792		0.7691	
Pacific	0.6873		0.8623		0.2101	
COUNTY TYPE						
Large Metro	0.3298		0.4212		0.8811	
Small Metro	0.8042		0.8915		0.5874	
250K – 1 Mil. Pop.	0.8807		0.7688		0.2545	
< 250K Pop.	0.8191		0.8908		0.6561	
Nonmetro	0.0218		0.0608		0.1565	
Urbanized	0.0150		0.1529		0.2852	
Less Urbanized	0.2137		0.2301		0.3851	
Completely Rural	0.8930		0.6748		0.6367	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.59C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	126	119	86	85	65	63
GEOGRAPHIC DIVISION						
Northeast	49	45	34	28	26	23
New England	25	26	18	15	15	9
Middle Atlantic	42	37	29	25	21	21
Midwest	54	51	37	37	29	30
East North Central	43	40	29	29	22	22
West North Central	33	33	22	24	20	19
South	78	73	53	51	39	39
South Atlantic	58	55	40	38	30	27
East South Central	29	29	21	20	16	17
West South Central	48	40	28	29	21	24
West	66	64	46	47	31	31
Mountain	30	31	21	19	16	14
Pacific	57	56	41	43	26	28
COUNTY TYPE						
Large Metro	110	107	63	66	42	44
Small Metro	85	85	54	57	43	39
250K – 1 Mil. Pop.	64	68	38	45	28	31
< 250K Pop.	58	54	37	36	32	24
Nonmetro	67	70	43	45	33	32
Urbanized	45	44	27	28	19	20
Less Urbanized	50	54	35	33	27	24
Completely Rural	24	23	14	16	12	10

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.59D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.38	0.36	0.26	0.26	0.20	0.19
GEOGRAPHIC DIVISION						
Northeast	0.83	0.76	0.59	0.48	0.44	0.40
New England	1.64	1.68	1.21	0.95	0.99	0.62
Middle Atlantic	0.96	0.84	0.67	0.56	0.48	0.49
Midwest	0.73	0.71	0.50	0.51	0.40	0.41
East North Central	0.85	0.81	0.57	0.57	0.43	0.44
West North Central	1.43	1.47	0.97	1.05	0.87	0.86
South	0.66	0.62	0.45	0.44	0.34	0.33
South Atlantic	0.99	0.93	0.68	0.65	0.51	0.47
East South Central	1.54	1.55	1.11	1.06	0.85	0.90
West South Central	1.22	1.02	0.71	0.74	0.55	0.61
West	0.84	0.81	0.59	0.60	0.39	0.40
Mountain	1.29	1.30	0.88	0.82	0.66	0.60
Pacific	1.05	1.02	0.74	0.79	0.48	0.51
COUNTY TYPE						
Large Metro	0.54	0.49	0.34	0.34	0.23	0.24
Small Metro	0.65	0.66	0.48	0.47	0.39	0.35
250K – 1 Mil. Pop.	0.76	0.80	0.53	0.57	0.40	0.43
< 250K Pop.	1.28	1.18	0.94	0.84	0.86	0.62
Nonmetro	0.98	0.93	0.72	0.67	0.58	0.55
Urbanized	1.42	1.40	0.94	0.97	0.82	0.74
Less Urbanized	1.47	1.33	1.17	1.00	0.92	0.84
Completely Rural	3.28	3.81	2.29	2.99	1.96	2.66

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.59P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.7558		0.5300		0.9671	
GEOGRAPHIC DIVISION						
Northeast	0.6999		0.1917		0.2700	
New England	0.7866		0.3819		0.1378	
Middle Atlantic	0.5351		0.3232		0.6962	
Midwest	0.1824		0.2265		0.6326	
East North Central	0.1799		0.1135		0.1982	
West North Central	0.6337		0.9745		0.4555	
South	0.1961		0.3859		0.7669	
South Atlantic	0.6733		0.8107		0.7209	
East South Central	0.3085		0.6169		0.6699	
West South Central	0.3371		0.3373		0.4825	
West	0.7125		0.8744		0.8357	
Mountain	0.2914		0.1215		0.5294	
Pacific	0.8111		0.3210		0.5547	
COUNTY TYPE						
Large Metro	0.4471		0.8387		0.4285	
Small Metro	0.8338		0.7149		0.4439	
250K – 1 Mil. Pop.	0.8208		0.2122		0.2926	
< 250K Pop.	0.5244		0.3735		0.0260	
Nonmetro	0.5127		0.6410		0.7814	
Urbanized	0.2381		0.5992		0.6956	
Less Urbanized	0.6764		0.9658		0.9166	
Completely Rural	0.5264		0.4009		0.4610	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.60C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	668	674	301	284	267	268
GEOGRAPHIC DIVISION						
Northeast	240	282	76	85	63	75
New England	116	144	29	53	25	47
Middle Atlantic	212	243	71	66	57	58
Midwest	271	250	125	133	109	123
East North Central	208	217	93	102	82	95
West North Central	170	140	79	83	69	75
South	450	429	241	189	216	181
South Atlantic	343	346	177	140	163	132
East South Central	183	168	90	77	82	73
West South Central	215	195	129	102	113	96
West	323	370	121	144	106	134
Mountain	150	162	64	83	55	75
Pacific	292	323	104	116	93	110
COUNTY TYPE						
Large Metro	565	559	182	186	156	178
Small Metro	406	427	163	172	147	153
250K – 1 Mil. Pop.	300	304	122	124	117	113
< 250K Pop.	267	306	104	117	86	101
Nonmetro	408	406	197	156	182	149
Urbanized	241	270	101	103	96	98
Less Urbanized	329	285	161	106	145	101
Completely Rural	143	145	84	56	79	56

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.60D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.36	0.35	0.16	0.15	0.14	0.14
GEOGRAPHIC DIVISION						
Northeast	0.68	0.79	0.22	0.24	0.18	0.21
New England	1.25	1.54	0.31	0.57	0.26	0.51
Middle Atlantic	0.81	0.93	0.27	0.25	0.22	0.22
Midwest	0.65	0.60	0.30	0.32	0.26	0.29
East North Central	0.71	0.74	0.32	0.35	0.28	0.32
West North Central	1.36	1.11	0.64	0.66	0.55	0.60
South	0.66	0.62	0.35	0.27	0.32	0.26
South Atlantic	0.94	0.93	0.48	0.38	0.45	0.36
East South Central	1.63	1.48	0.80	0.68	0.73	0.64
West South Central	1.05	0.94	0.63	0.49	0.55	0.46
West	0.76	0.85	0.28	0.33	0.25	0.31
Mountain	1.18	1.24	0.50	0.63	0.44	0.57
Pacific	0.98	1.07	0.35	0.38	0.31	0.36
COUNTY TYPE						
Large Metro	0.48	0.49	0.17	0.18	0.15	0.17
Small Metro	0.65	0.61	0.30	0.29	0.27	0.26
250K – 1 Mil. Pop.	0.76	0.68	0.36	0.31	0.34	0.29
< 250K Pop.	1.18	1.16	0.54	0.60	0.44	0.51
Nonmetro	0.88	0.92	0.55	0.46	0.51	0.44
Urbanized	1.38	1.50	0.73	0.75	0.70	0.72
Less Urbanized	1.28	1.24	0.84	0.64	0.78	0.61
Completely Rural	2.77	3.23	2.05	1.48	1.96	1.47

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.60P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.1985		0.2327		0.4942	
GEOGRAPHIC DIVISION						
Northeast	0.6807		0.4310		0.4155	
New England	0.7458		0.2472		0.2732	
Middle Atlantic	0.7659		0.8326		0.8097	
Midwest	0.6495		0.7167		0.6172	
East North Central	0.4889		0.9568		0.6329	
West North Central	0.8637		0.6219		0.8286	
South	0.0418		0.0236		0.0676	
South Atlantic	0.4133		0.2068		0.2038	
East South Central	0.2477		0.1826		0.2444	
West South Central	0.0392		0.1313		0.4134	
West	0.9368		0.8157		0.5993	
Mountain	0.8824		0.8108		0.6701	
Pacific	0.9894		0.9253		0.7495	
COUNTY TYPE						
Large Metro	0.3431		0.5502		0.7572	
Small Metro	0.3392		0.7462		0.5633	
250K – 1 Mil. Pop.	0.0379		0.5105		0.3641	
< 250K Pop.	0.3036		0.7123		0.7668	
Nonmetro	0.8988		0.2898		0.3898	
Urbanized	0.4496		0.2508		0.2439	
Less Urbanized	0.6329		0.1431		0.1990	
Completely Rural	0.5441		0.0650		0.1113	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.61C (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	685	695	320	304	281	280
GEOGRAPHIC DIVISION						
Northeast	248	289	82	92	66	82
New England	122	146	35	54	28	49
Middle Atlantic	217	250	75	75	61	65
Midwest	279	255	131	140	114	128
East North Central	215	218	97	106	85	96
West North Central	174	143	83	88	74	79
South	457	446	252	201	222	188
South Atlantic	348	357	183	147	166	136
East South Central	187	175	95	82	86	77
West South Central	221	200	135	108	115	99
West	332	376	131	157	112	139
Mountain	157	164	67	86	58	77
Pacific	298	328	114	130	97	114
COUNTY TYPE						
Large Metro	592	582	197	204	163	188
Small Metro	419	446	174	187	155	162
250K – 1 Mil. Pop.	309	320	128	137	121	121
< 250K Pop.	281	321	115	124	94	103
Nonmetro	426	425	208	170	190	157
Urbanized	251	286	105	113	99	105
Less Urbanized	348	299	172	117	152	106
Completely Rural	153	154	90	59	84	58

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.61D (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.31	0.31	0.14	0.14	0.13	0.13
GEOGRAPHIC DIVISION						
Northeast	0.60	0.70	0.20	0.22	0.16	0.20
New England	1.12	1.34	0.32	0.49	0.25	0.45
Middle Atlantic	0.71	0.82	0.25	0.24	0.20	0.21
Midwest	0.57	0.52	0.27	0.29	0.23	0.26
East North Central	0.63	0.63	0.28	0.31	0.25	0.28
West North Central	1.18	0.96	0.56	0.59	0.50	0.53
South	0.57	0.55	0.32	0.25	0.28	0.23
South Atlantic	0.82	0.83	0.43	0.34	0.39	0.32
East South Central	1.42	1.32	0.72	0.62	0.65	0.58
West South Central	0.91	0.81	0.56	0.44	0.47	0.40
West	0.66	0.73	0.26	0.31	0.22	0.27
Mountain	1.04	1.06	0.44	0.56	0.38	0.50
Pacific	0.84	0.92	0.32	0.36	0.27	0.32
COUNTY TYPE						
Large Metro	0.43	0.43	0.16	0.17	0.13	0.15
Small Metro	0.55	0.53	0.27	0.26	0.24	0.23
250K – 1 Mil. Pop.	0.64	0.59	0.31	0.28	0.29	0.26
< 250K Pop.	1.01	1.02	0.50	0.53	0.40	0.44
Nonmetro	0.78	0.81	0.50	0.42	0.46	0.39
Urbanized	1.21	1.29	0.63	0.68	0.61	0.63
Less Urbanized	1.13	1.09	0.77	0.59	0.70	0.54
Completely Rural	2.48	2.95	1.89	1.39	1.79	1.39

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.61P (REVISED) Smokeless Tobacco Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.1913		0.3214		0.5101	
GEOGRAPHIC DIVISION						
Northeast	0.7423		0.7881		0.7056	
New England	0.7204		0.4422		0.5739	
Middle Atlantic	0.8522		0.8801		0.9275	
Midwest	0.4933		0.5391		0.5714	
East North Central	0.3616		0.6568		0.4474	
West North Central	0.9402		0.6587		0.9966	
South	0.0276		0.0483		0.0843	
South Atlantic	0.3852		0.2475		0.1893	
East South Central	0.1957		0.2421		0.2952	
West South Central	0.0260		0.2082		0.5096	
West	0.9887		0.8059		0.5791	
Mountain	0.9433		0.8292		0.7929	
Pacific	0.9773		0.6874		0.6407	
COUNTY TYPE						
Large Metro	0.2939		0.6064		0.6335	
Small Metro	0.3259		0.7631		0.4154	
250K – 1 Mil. Pop.	0.0441		0.6782		0.4851	
< 250K Pop.	0.3748		0.9090		0.7400	
Nonmetro	0.9927		0.3922		0.3970	
Urbanized	0.3485		0.2036		0.2211	
Less Urbanized	0.6965		0.2047		0.2291	
Completely Rural	0.4374		0.0723		0.0786	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.