

Table 2.67C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	671	721	914	957	968	1,019
GEOGRAPHIC DIVISION						
Northeast	263	299	356	375	399	421
New England	106	121	165	175	187	191
Middle Atlantic	241	270	313	330	354	372
Midwest	227	232	332	353	369	372
East North Central	186	186	267	285	290	311
West North Central	131	136	203	197	223	207
South	422	465	606	634	647	631
South Atlantic	301	341	466	481	504	495
East South Central	173	169	211	218	200	213
West South Central	236	250	313	335	319	306
West	353	386	462	454	501	504
Mountain	136	168	203	202	228	205
Pacific	326	350	412	410	442	462
COUNTY TYPE						
Large Metro	1,663	1,745	1,463	1,514	1,247	1,308
Small Metro	1,155	1,339	957	1,114	792	931
250K – 1 Mil. Pop.	905	1,119	776	926	648	785
< 250K Pop.	770	818	600	672	491	548
Nonmetro	933	882	735	667	619	550
Urbanized	597	617	462	485	370	383
Less Urbanized	751	660	585	504	489	416
Completely Rural	400	410	314	289	258	243

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.67D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.29	0.37	0.39	0.39	0.41
GEOGRAPHIC DIVISION						
Northeast	0.57	0.65	0.78	0.82	0.87	0.92
New England	0.88	1.01	1.37	1.45	1.56	1.59
Middle Atlantic	0.71	0.80	0.93	0.98	1.05	1.10
Midwest	0.41	0.42	0.61	0.64	0.67	0.68
East North Central	0.49	0.48	0.70	0.74	0.76	0.81
West North Central	0.80	0.82	1.24	1.19	1.36	1.26
South	0.47	0.52	0.68	0.71	0.73	0.70
South Atlantic	0.64	0.72	0.99	1.01	1.07	1.04
East South Central	1.18	1.15	1.44	1.48	1.37	1.45
West South Central	0.87	0.90	1.15	1.21	1.17	1.11
West	0.62	0.67	0.82	0.79	0.89	0.88
Mountain	0.81	0.98	1.21	1.17	1.35	1.18
Pacific	0.82	0.88	1.04	1.03	1.12	1.16
COUNTY TYPE						
Large Metro	0.38	0.41	0.52	0.52	0.55	0.59
Small Metro	0.45	0.50	0.66	0.65	0.71	0.66
250K – 1 Mil. Pop.	0.56	0.60	0.75	0.78	0.80	0.80
< 250K Pop.	0.83	0.88	1.25	1.16	1.34	1.15
Nonmetro	0.64	0.63	0.86	0.86	0.86	0.85
Urbanized	0.96	0.92	1.36	1.50	1.33	1.40
Less Urbanized	0.98	0.94	1.29	1.17	1.30	1.24
Completely Rural	1.96	2.31	2.55	2.90	2.76	2.75

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.67P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3036		0.5949		0.6650	
GEOGRAPHIC DIVISION						
Northeast	0.3251		0.1787		0.8574	
New England	0.5399		0.4284		0.4246	
Middle Atlantic	0.4114		0.2638		0.5388	
Midwest	0.1504		0.2390		0.2243	
East North Central	0.0804		0.2927		0.3741	
West North Central	0.9112		0.5621		0.3834	
South	0.5328		0.4566		0.8762	
South Atlantic	0.3233		0.4390		0.1632	
East South Central	0.1966		0.2086		0.3853	
West South Central	0.1625		0.1244		0.0766	
West	0.1759		0.8754		0.7452	
Mountain	0.1805		0.9266		0.4538	
Pacific	0.3786		0.8911		0.9716	
COUNTY TYPE						
Large Metro	0.4381		0.6206		0.9699	
Small Metro	0.8894		0.6671		0.1585	
250K – 1 Mil. Pop.	0.8254		0.6958		0.6262	
< 250K Pop.	0.9662		0.2559		0.1105	
Nonmetro	0.1785		0.2771		0.4019	
Urbanized	0.5403		0.4767		0.1869	
Less Urbanized	0.0600		0.1201		0.0609	
Completely Rural	0.1869		0.1522		0.3199	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.68C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	112	112	106	107	82	87
GEOGRAPHIC DIVISION						
Northeast	46	45	43	44	38	35
New England	21	19	19	18	18	16
Middle Atlantic	41	40	39	40	33	31
Midwest	43	44	41	40	32	32
East North Central	35	35	33	33	25	26
West North Central	25	27	24	21	20	18
South	69	71	66	66	51	51
South Atlantic	51	54	48	51	39	40
East South Central	26	23	24	22	16	17
West South Central	39	41	40	37	28	29
West	64	60	59	57	43	49
Mountain	27	27	25	27	21	19
Pacific	58	54	53	51	38	44
COUNTY TYPE						
Large Metro	119	135	104	116	71	76
Small Metro	91	92	80	79	51	53
250K – 1 Mil. Pop.	70	73	62	63	41	42
< 250K Pop.	58	57	51	47	30	32
Nonmetro	63	64	54	53	35	39
Urbanized	41	41	35	34	21	25
Less Urbanized	48	45	41	41	27	27
Completely Rural	22	29	17	20	11	15

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.68D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.44	0.44	0.42	0.42	0.32	0.34
GEOGRAPHIC DIVISION						
Northeast	1.01	1.00	0.96	0.98	0.84	0.78
New England	1.79	1.67	1.65	1.54	1.53	1.35
Middle Atlantic	1.21	1.22	1.15	1.21	0.99	0.95
Midwest	0.76	0.79	0.71	0.70	0.57	0.57
East North Central	0.88	0.88	0.83	0.83	0.64	0.66
West North Central	1.50	1.59	1.40	1.28	1.21	1.09
South	0.75	0.78	0.73	0.73	0.56	0.56
South Atlantic	1.09	1.17	1.03	1.10	0.85	0.86
East South Central	1.73	1.56	1.61	1.46	1.07	1.12
West South Central	1.30	1.35	1.33	1.24	0.93	0.95
West	1.06	1.00	0.98	0.95	0.72	0.80
Mountain	1.52	1.50	1.42	1.48	1.15	1.08
Pacific	1.35	1.27	1.24	1.20	0.89	1.04
COUNTY TYPE						
Large Metro	0.64	0.63	0.60	0.59	0.47	0.48
Small Metro	0.81	0.82	0.78	0.75	0.60	0.63
250K – 1 Mil. Pop.	0.99	1.03	0.94	0.93	0.72	0.75
< 250K Pop.	1.31	1.23	1.30	1.15	1.03	1.06
Nonmetro	0.94	1.00	0.88	0.92	0.68	0.77
Urbanized	1.40	1.39	1.31	1.29	1.01	1.13
Less Urbanized	1.43	1.41	1.31	1.41	1.00	1.10
Completely Rural	2.48	3.78	2.29	2.53	1.77	2.56

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.68P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.1339		0.0576		0.1687	
GEOGRAPHIC DIVISION						
Northeast	0.0001		0.0001		0.1321	
New England	0.1653		0.0441		0.7271	
Middle Atlantic	0.0003		0.0007		0.1274	
Midwest	0.8701		0.6798		0.5239	
East North Central	0.4920		0.3214		0.1799	
West North Central	0.2077		0.0217		0.4583	
South	0.6744		0.9685		0.4101	
South Atlantic	0.6007		0.9287		0.9170	
East South Central	0.9384		0.9038		0.5834	
West South Central	0.9858		0.9775		0.3213	
West	0.7475		0.7665		0.3912	
Mountain	0.1531		0.3370		0.0947	
Pacific	0.7326		0.8817		0.8526	
COUNTY TYPE						
Large Metro	0.1128		0.0476		0.1129	
Small Metro	0.9297		0.7836		0.8476	
250K – 1 Mil. Pop.	0.8383		0.9616		0.6033	
< 250K Pop.	0.6101		0.6622		0.6826	
Nonmetro	0.5260		0.5302		0.8883	
Urbanized	0.7085		0.6233		0.7342	
Less Urbanized	0.4907		0.5586		0.6001	
Completely Rural	0.8290		0.8195		0.9489	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.69C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	116	125	140	147	168	170
GEOGRAPHIC DIVISION						
Northeast	47	50	57	58	69	69
New England	22	21	27	24	31	32
Middle Atlantic	42	45	50	53	62	60
Midwest	41	37	50	45	66	63
East North Central	35	30	43	36	58	51
West North Central	22	22	26	27	32	36
South	71	74	89	91	104	104
South Atlantic	57	53	66	66	76	77
East South Central	24	31	27	34	37	37
West South Central	36	43	50	54	60	56
West	65	83	77	91	88	96
Mountain	33	54	37	55	41	51
Pacific	55	62	66	71	78	81
COUNTY TYPE						
Large Metro	343	325	323	316	282	261
Small Metro	227	242	218	233	196	208
250K – 1 Mil. Pop.	189	189	182	184	165	165
< 250K Pop.	130	161	124	152	111	134
Nonmetro	157	167	145	156	122	133
Urbanized	127	119	117	107	96	86
Less Urbanized	111	123	103	118	82	103
Completely Rural	45	46	42	41	33	35

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.69D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.35	0.38	0.43	0.45	0.51	0.52
GEOGRAPHIC DIVISION						
Northeast	0.81	0.84	0.97	0.98	1.18	1.16
New England	1.45	1.40	1.80	1.57	2.05	2.08
Middle Atlantic	0.96	1.02	1.14	1.20	1.42	1.37
Midwest	0.55	0.51	0.68	0.62	0.90	0.86
East North Central	0.69	0.60	0.84	0.72	1.13	1.02
West North Central	0.95	0.96	1.14	1.20	1.42	1.60
South	0.61	0.63	0.76	0.78	0.89	0.89
South Atlantic	0.98	0.89	1.13	1.12	1.30	1.30
East South Central	1.26	1.66	1.45	1.80	1.98	1.98
West South Central	0.91	1.09	1.29	1.37	1.52	1.44
West	0.83	1.06	0.99	1.16	1.13	1.22
Mountain	1.38	2.28	1.57	2.34	1.73	2.18
Pacific	1.01	1.13	1.22	1.29	1.42	1.47
COUNTY TYPE						
Large Metro	0.52	0.49	0.62	0.60	0.71	0.70
Small Metro	0.61	0.81	0.73	0.89	0.94	0.97
250K – 1 Mil. Pop.	0.78	1.07	0.91	1.13	1.19	1.19
< 250K Pop.	0.96	1.10	1.17	1.42	1.43	1.65
Nonmetro	0.76	0.85	0.99	0.96	1.17	1.18
Urbanized	1.14	1.16	1.38	1.31	1.71	1.50
Less Urbanized	1.11	1.40	1.50	1.61	1.66	2.03
Completely Rural	2.28	3.20	2.66	3.30	3.56	4.00

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.69P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.0090		0.1022		0.2782	
GEOGRAPHIC DIVISION						
Northeast	0.4260		0.2345		0.3067	
New England	0.9326		0.8248		0.6889	
Middle Atlantic	0.3541		0.1402		0.3443	
Midwest	0.4476		0.5647		0.2004	
East North Central	0.5934		0.9282		0.6308	
West North Central	0.5842		0.2474		0.1043	
South	0.5651		0.9970		0.2541	
South Atlantic	0.4664		0.3126		0.0422	
East South Central	0.2551		0.0699		0.2663	
West South Central	0.2199		0.8459		0.8837	
West	0.0045		0.1019		0.1016	
Mountain	0.0894		0.3168		0.4967	
Pacific	0.0216		0.1880		0.1320	
COUNTY TYPE						
Large Metro	0.0142		0.1237		0.4933	
Small Metro	0.3567		0.4641		0.3687	
250K – 1 Mil. Pop.	0.1946		0.1610		0.3705	
< 250K Pop.	0.7071		0.4646		0.7447	
Nonmetro	0.5438		0.9123		0.8400	
Urbanized	0.9746		0.6886		0.6743	
Less Urbanized	0.4260		0.4274		0.9898	
Completely Rural	0.7904		0.0947		0.9621	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.70C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	614	679	873	933	914	986
GEOGRAPHIC DIVISION						
Northeast	243	289	340	362	371	407
New England	82	115	152	173	172	191
Middle Atlantic	228	261	302	316	332	357
Midwest	208	212	324	346	355	358
East North Central	169	169	255	282	271	301
West North Central	121	121	197	189	218	196
South	396	442	587	625	624	613
South Atlantic	282	323	455	485	489	484
East South Central	166	156	208	203	199	200
West South Central	221	242	295	323	304	295
West	330	356	439	434	467	487
Mountain	124	149	191	190	214	198
Pacific	305	326	391	392	412	445
COUNTY TYPE						
Large Metro	1,555	1,621	1,358	1,400	1,169	1,229
Small Metro	1,066	1,241	875	1,023	728	865
250K – 1 Mil. Pop.	833	1,027	713	841	596	721
< 250K Pop.	700	754	534	616	446	510
Nonmetro	851	798	661	593	569	499
Urbanized	530	537	398	416	326	336
Less Urbanized	665	597	512	448	440	374
Completely Rural	367	363	286	250	245	218

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.70D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.33	0.36	0.46	0.49	0.49	0.52
GEOGRAPHIC DIVISION						
Northeast	0.68	0.81	0.96	1.02	1.05	1.15
New England	0.88	1.23	1.63	1.85	1.84	2.04
Middle Atlantic	0.87	1.00	1.16	1.21	1.27	1.36
Midwest	0.50	0.51	0.78	0.82	0.85	0.86
East North Central	0.58	0.58	0.87	0.96	0.93	1.03
West North Central	0.97	0.97	1.58	1.51	1.75	1.56
South	0.58	0.64	0.86	0.90	0.92	0.89
South Atlantic	0.77	0.87	1.24	1.31	1.34	1.31
East South Central	1.48	1.37	1.85	1.78	1.77	1.76
West South Central	1.08	1.17	1.45	1.56	1.49	1.42
West	0.77	0.82	1.03	1.00	1.10	1.12
Mountain	0.98	1.13	1.50	1.45	1.68	1.51
Pacific	1.02	1.08	1.31	1.30	1.38	1.47
COUNTY TYPE						
Large Metro	0.45	0.50	0.64	0.66	0.67	0.74
Small Metro	0.56	0.60	0.85	0.82	0.89	0.86
250K – 1 Mil. Pop.	0.67	0.71	0.96	0.99	1.01	1.03
< 250K Pop.	1.08	1.10	1.60	1.49	1.69	1.46
Nonmetro	0.78	0.77	1.06	1.07	1.06	1.05
Urbanized	1.20	1.10	1.70	1.88	1.64	1.76
Less Urbanized	1.20	1.15	1.56	1.45	1.58	1.50
Completely Rural	2.32	2.71	3.20	3.44	3.44	3.25

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.70P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 26 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.5899		0.8865		0.4951	
GEOGRAPHIC DIVISION						
Northeast	0.7671		0.4444		0.9220	
New England	0.6895		0.5265		0.3614	
Middle Atlantic	0.8658		0.5856		0.7165	
Midwest	0.1127		0.1888		0.1611	
East North Central	0.0841		0.3553		0.3872	
West North Central	0.7018		0.3097		0.2139	
South	0.4711		0.4318		0.7498	
South Atlantic	0.4731		0.5417		0.2799	
East South Central	0.2388		0.3071		0.5206	
West South Central	0.1889		0.1123		0.0753	
West	0.3551		0.6454		0.5055	
Mountain	0.4513		0.6345		0.2853	
Pacific	0.5012		0.7617		0.7896	
COUNTY TYPE						
Large Metro	0.9440		0.9860		0.7959	
Small Metro	0.7995		0.7233		0.1679	
250K – 1 Mil. Pop.	0.8941		0.6914		0.6472	
< 250K Pop.	0.8044		0.3183		0.1132	
Nonmetro	0.3069		0.3204		0.4501	
Urbanized	0.4722		0.4345		0.1721	
Less Urbanized	0.1552		0.1038		0.0717	
Completely Rural	0.1353		0.2176		0.3222	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.71C (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	640	694	897	955	954	1,011
GEOGRAPHIC DIVISION						
Northeast	250	292	351	371	390	415
New England	89	117	159	174	183	192
Middle Atlantic	233	264	311	325	347	365
Midwest	215	217	332	352	367	373
East North Central	176	173	264	285	287	311
West North Central	123	125	202	195	224	207
South	408	456	599	639	644	630
South Atlantic	292	331	462	488	502	498
East South Central	168	165	210	214	203	209
West South Central	224	248	300	335	312	301
West	343	365	457	446	492	498
Mountain	135	161	203	198	230	203
Pacific	315	330	406	403	432	456
COUNTY TYPE						
Large Metro	1,635	1,698	1,438	1,476	1,233	1,287
Small Metro	1,123	1,301	932	1,083	780	916
250K – 1 Mil. Pop.	883	1,087	759	900	640	774
< 250K Pop.	744	789	578	650	480	536
Nonmetro	908	853	712	645	608	540
Urbanized	576	596	444	470	363	375
Less Urbanized	727	636	565	486	477	407
Completely Rural	387	391	304	274	254	234

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 2.71D (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
Standard Errors of Percentages, 2006 and 2007**

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.29	0.31	0.41	0.43	0.43	0.45
GEOGRAPHIC DIVISION						
Northeast	0.61	0.71	0.85	0.89	0.94	1.00
New England	0.82	1.08	1.47	1.60	1.68	1.77
Middle Atlantic	0.77	0.87	1.02	1.06	1.14	1.20
Midwest	0.44	0.44	0.68	0.72	0.75	0.76
East North Central	0.51	0.50	0.77	0.83	0.84	0.91
West North Central	0.84	0.85	1.37	1.32	1.52	1.39
South	0.51	0.56	0.75	0.79	0.81	0.78
South Atlantic	0.69	0.77	1.09	1.14	1.18	1.16
East South Central	1.28	1.24	1.60	1.61	1.55	1.58
West South Central	0.92	1.01	1.24	1.36	1.28	1.22
West	0.68	0.71	0.91	0.87	0.98	0.97
Mountain	0.89	1.04	1.34	1.28	1.52	1.31
Pacific	0.89	0.92	1.15	1.13	1.22	1.28
COUNTY TYPE						
Large Metro	0.40	0.44	0.57	0.58	0.60	0.65
Small Metro	0.49	0.53	0.73	0.71	0.79	0.73
250K – 1 Mil. Pop.	0.58	0.63	0.81	0.86	0.88	0.89
< 250K Pop.	0.94	0.94	1.42	1.27	1.50	1.26
Nonmetro	0.69	0.68	0.94	0.96	0.95	0.95
Urbanized	1.04	0.96	1.51	1.64	1.48	1.55
Less Urbanized	1.07	1.02	1.41	1.31	1.43	1.38
Completely Rural	2.09	2.53	2.79	3.23	3.01	3.03

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.71P (REVISED) Alcohol Use in Lifetime, Past Year, and Past Month among Persons Aged 18 or Older, by Geographic Characteristics:
P Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TIME PERIOD					
	Lifetime		Past Year		Past Month	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3294		0.6820		0.6451	
GEOGRAPHIC DIVISION						
Northeast	0.6662		0.3561		0.9352	
New England	0.7098		0.5637		0.4091	
Middle Atlantic	0.7453		0.4527		0.6066	
Midwest	0.1500		0.2397		0.2698	
East North Central	0.1090		0.3759		0.4703	
West North Central	0.7697		0.4123		0.3607	
South	0.4196		0.4275		0.8900	
South Atlantic	0.4001		0.4542		0.1683	
East South Central	0.1763		0.1982		0.4011	
West South Central	0.1334		0.1110		0.0827	
West	0.1502		0.8893		0.7200	
Mountain	0.2532		0.8410		0.3809	
Pacific	0.2945		0.9429		0.9914	
COUNTY TYPE						
Large Metro	0.6221		0.8120		0.8980	
Small Metro	0.6726		0.8725		0.2598	
250K – 1 Mil. Pop.	0.6979		0.4753		0.8241	
< 250K Pop.	0.8652		0.2783		0.1430	
Nonmetro	0.2751		0.3570		0.4670	
Urbanized	0.4757		0.4435		0.1984	
Less Urbanized	0.1349		0.1818		0.0925	
Completely Rural	0.1290		0.1280		0.2799	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.72C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	968	1,019	746	770	427	429
GEOGRAPHIC DIVISION						
Northeast	399	421	301	321	172	174
New England	187	191	133	155	80	91
Middle Atlantic	354	372	270	279	153	149
Midwest	369	372	318	325	193	174
East North Central	290	311	259	254	145	149
West North Central	223	207	183	184	129	93
South	647	631	480	467	246	264
South Atlantic	504	495	369	365	194	206
East South Central	200	213	155	143	87	83
West South Central	319	306	257	252	148	136
West	501	504	370	387	207	224
Mountain	228	205	170	171	97	121
Pacific	442	462	327	344	183	187
COUNTY TYPE						
Large Metro	1,247	1,308	713	685	339	340
Small Metro	792	931	425	523	222	261
250K – 1 Mil. Pop.	648	785	347	434	184	227
< 250K Pop.	491	548	269	305	135	130
Nonmetro	619	550	359	353	185	165
Urbanized	370	383	191	263	100	130
Less Urbanized	489	416	299	253	146	102
Completely Rural	258	243	136	130	57	35

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.72D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.39	0.41	0.30	0.31	0.17	0.17
GEOGRAPHIC DIVISION						
Northeast	0.87	0.92	0.66	0.70	0.37	0.38
New England	1.56	1.59	1.11	1.29	0.67	0.75
Middle Atlantic	1.05	1.10	0.80	0.83	0.45	0.44
Midwest	0.67	0.68	0.58	0.59	0.35	0.32
East North Central	0.76	0.81	0.68	0.66	0.38	0.39
West North Central	1.36	1.26	1.12	1.11	0.78	0.56
South	0.73	0.70	0.54	0.52	0.28	0.29
South Atlantic	1.07	1.04	0.78	0.77	0.41	0.43
East South Central	1.37	1.45	1.06	0.97	0.59	0.56
West South Central	1.17	1.11	0.94	0.91	0.54	0.49
West	0.89	0.88	0.66	0.68	0.37	0.39
Mountain	1.35	1.18	1.01	0.99	0.57	0.70
Pacific	1.12	1.16	0.83	0.86	0.46	0.47
COUNTY TYPE						
Large Metro	0.55	0.59	0.45	0.44	0.24	0.25
Small Metro	0.71	0.66	0.52	0.54	0.31	0.31
250K – 1 Mil. Pop.	0.80	0.80	0.63	0.62	0.38	0.39
< 250K Pop.	1.34	1.15	0.95	0.98	0.53	0.50
Nonmetro	0.86	0.85	0.67	0.69	0.40	0.37
Urbanized	1.33	1.40	0.95	1.06	0.56	0.67
Less Urbanized	1.30	1.24	0.99	1.01	0.59	0.48
Completely Rural	2.76	2.75	2.23	2.02	1.09	0.65

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.72P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.6650		0.4861		0.9200	
GEOGRAPHIC DIVISION						
Northeast	0.8574		0.5892		0.1042	
New England	0.4246		0.8325		0.6801	
Middle Atlantic	0.5388		0.6181		0.1005	
Midwest	0.2243		0.3096		0.9522	
East North Central	0.3741		0.5678		0.5516	
West North Central	0.3834		0.3320		0.4572	
South	0.8762		0.6217		0.8848	
South Atlantic	0.1632		0.2261		0.4093	
East South Central	0.3853		0.8066		0.2605	
West South Central	0.0766		0.4120		0.1262	
West	0.7452		0.6988		0.3960	
Mountain	0.4538		0.6243		0.4205	
Pacific	0.9716		0.8553		0.6092	
COUNTY TYPE						
Large Metro	0.9699		0.8394		0.5390	
Small Metro	0.1585		0.3519		0.4385	
250K – 1 Mil. Pop.	0.6262		0.5412		0.2286	
< 250K Pop.	0.1105		0.4662		0.6405	
Nonmetro	0.4019		0.3073		0.8948	
Urbanized	0.1869		0.0038		0.0348	
Less Urbanized	0.0609		0.7270		0.2187	
Completely Rural	0.3199		0.1891		0.0139	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.73C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	82	87	69	69	33	37
GEOGRAPHIC DIVISION						
Northeast	38	35	31	28	17	14
New England	18	16	14	12	6	6
Middle Atlantic	33	31	27	25	16	12
Midwest	32	32	26	27	13	18
East North Central	25	26	20	21	10	13
West North Central	20	18	17	16	8	12
South	51	51	43	42	20	22
South Atlantic	39	40	32	32	14	16
East South Central	16	17	13	14	5	6
West South Central	28	29	24	24	12	13
West	43	49	38	38	19	19
Mountain	21	19	17	16	8	7
Pacific	38	44	34	34	17	17
COUNTY TYPE						
Large Metro	71	76	58	55	28	26
Small Metro	51	53	40	39	19	18
250K – 1 Mil. Pop.	41	42	33	31	15	14
< 250K Pop.	30	32	22	23	11	11
Nonmetro	35	39	28	32	11	18
Urbanized	21	25	17	18	7	9
Less Urbanized	27	27	21	23	7	12
Completely Rural	11	15	8	14	4	12

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.73D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.32	0.34	0.27	0.27	0.13	0.15
GEOGRAPHIC DIVISION						
Northeast	0.84	0.78	0.68	0.63	0.37	0.31
New England	1.53	1.35	1.21	1.07	0.50	0.56
Middle Atlantic	0.99	0.95	0.82	0.76	0.47	0.37
Midwest	0.57	0.57	0.46	0.47	0.23	0.31
East North Central	0.64	0.66	0.51	0.54	0.26	0.33
West North Central	1.21	1.09	0.98	0.95	0.47	0.71
South	0.56	0.56	0.47	0.47	0.22	0.24
South Atlantic	0.85	0.86	0.70	0.70	0.31	0.35
East South Central	1.07	1.12	0.88	0.96	0.36	0.40
West South Central	0.93	0.95	0.80	0.80	0.40	0.42
West	0.72	0.80	0.62	0.63	0.31	0.31
Mountain	1.15	1.08	0.97	0.87	0.45	0.41
Pacific	0.89	1.04	0.79	0.79	0.40	0.41
COUNTY TYPE						
Large Metro	0.47	0.48	0.39	0.36	0.20	0.19
Small Metro	0.60	0.63	0.49	0.49	0.24	0.24
250K – 1 Mil. Pop.	0.72	0.75	0.60	0.58	0.29	0.28
< 250K Pop.	1.03	1.06	0.82	0.84	0.44	0.46
Nonmetro	0.68	0.77	0.61	0.66	0.26	0.43
Urbanized	1.01	1.13	0.94	0.87	0.42	0.51
Less Urbanized	1.00	1.10	0.91	1.00	0.37	0.59
Completely Rural	1.77	2.56	1.51	2.62	0.82	2.63

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.73P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 12 to 17, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.1687		0.1229		0.8667	
GEOGRAPHIC DIVISION						
Northeast	0.1321		0.0370		0.5591	
New England	0.7271		0.5119		0.8231	
Middle Atlantic	0.1274		0.0452		0.4543	
Midwest	0.5239		0.3878		0.1362	
East North Central	0.1799		0.1048		0.2546	
West North Central	0.4583		0.5044		0.3301	
South	0.4101		0.9791		0.8499	
South Atlantic	0.9170		0.9559		0.9161	
East South Central	0.5834		0.6764		0.9159	
West South Central	0.3213		0.8278		0.8093	
West	0.3912		0.0824		0.3964	
Mountain	0.0947		0.3807		0.5473	
Pacific	0.8526		0.1265		0.5065	
COUNTY TYPE						
Large Metro	0.1129		0.0514		0.3742	
Small Metro	0.8476		0.6001		0.2226	
250K – 1 Mil. Pop.	0.6033		0.6076		0.1059	
< 250K Pop.	0.6826		0.8389		0.9485	
Nonmetro	0.8883		0.6471		0.0082	
Urbanized	0.7342		0.7178		0.1344	
Less Urbanized	0.6001		0.6949		0.0582	
Completely Rural	0.9489		0.9080		0.3166	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.74C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	168	170	169	169	125	114
GEOGRAPHIC DIVISION						
Northeast	69	69	71	73	59	50
New England	31	32	33	35	30	27
Middle Atlantic	62	60	63	64	50	42
Midwest	66	63	71	62	53	46
East North Central	58	51	61	52	44	39
West North Central	32	36	37	34	31	27
South	104	104	102	99	72	66
South Atlantic	76	77	73	74	53	50
East South Central	37	37	35	36	27	22
West South Central	60	56	60	56	43	36
West	88	96	90	92	62	64
Mountain	41	51	38	42	27	27
Pacific	78	81	81	82	55	58
COUNTY TYPE						
Large Metro	282	261	209	205	110	101
Small Metro	196	208	149	159	88	83
250K – 1 Mil. Pop.	165	165	121	124	71	68
< 250K Pop.	111	134	88	102	53	47
Nonmetro	122	133	98	107	55	47
Urbanized	96	86	75	70	40	35
Less Urbanized	82	103	65	83	33	34
Completely Rural	33	35	30	26	16	11

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.74D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.51	0.52	0.52	0.52	0.38	0.35
GEOGRAPHIC DIVISION						
Northeast	1.18	1.16	1.21	1.24	1.00	0.85
New England	2.05	2.08	2.13	2.28	1.94	1.78
Middle Atlantic	1.42	1.37	1.45	1.47	1.16	0.96
Midwest	0.90	0.86	0.96	0.85	0.72	0.63
East North Central	1.13	1.02	1.19	1.04	0.86	0.77
West North Central	1.42	1.60	1.64	1.52	1.34	1.18
South	0.89	0.89	0.88	0.85	0.62	0.56
South Atlantic	1.30	1.30	1.25	1.25	0.90	0.85
East South Central	1.98	1.98	1.86	1.93	1.42	1.19
West South Central	1.52	1.44	1.54	1.43	1.10	0.91
West	1.13	1.22	1.15	1.17	0.79	0.82
Mountain	1.73	2.18	1.60	1.78	1.13	1.13
Pacific	1.42	1.47	1.49	1.48	1.02	1.06
COUNTY TYPE						
Large Metro	0.71	0.70	0.75	0.72	0.53	0.49
Small Metro	0.94	0.97	0.89	0.94	0.71	0.62
250K – 1 Mil. Pop.	1.19	1.19	1.12	1.15	0.88	0.78
< 250K Pop.	1.43	1.65	1.44	1.53	1.23	1.03
Nonmetro	1.17	1.18	1.14	1.16	0.82	0.72
Urbanized	1.71	1.50	1.84	1.54	1.31	1.09
Less Urbanized	1.66	2.03	1.55	1.93	1.03	1.07
Completely Rural	3.56	4.00	3.49	3.39	2.35	2.70

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.74P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 to 25, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.2782		0.5757		0.0997	
GEOGRAPHIC DIVISION						
Northeast	0.3067		0.0623		0.0653	
New England	0.6889		0.4593		0.1730	
Middle Atlantic	0.3443		0.0835		0.1882	
Midwest	0.2004		0.2906		0.2763	
East North Central	0.6308		0.4626		0.8363	
West North Central	0.1043		0.4319		0.1239	
South	0.2541		0.2145		0.3653	
South Atlantic	0.0422		0.0557		0.9889	
East South Central	0.2663		0.2514		0.0287	
West South Central	0.8837		0.5657		0.8080	
West	0.1016		0.9262		0.7047	
Mountain	0.4967		0.7795		0.6882	
Pacific	0.1320		0.9719		0.8201	
COUNTY TYPE						
Large Metro	0.4933		0.8353		0.4778	
Small Metro	0.3687		0.3707		0.3878	
250K – 1 Mil. Pop.	0.3705		0.3987		0.8262	
< 250K Pop.	0.7447		0.6857		0.2704	
Nonmetro	0.8400		0.8672		0.0521	
Urbanized	0.6743		0.7433		0.3073	
Less Urbanized	0.9898		0.9827		0.1866	
Completely Rural	0.9621		0.7227		0.0795	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.75C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	914	986	697	714	392	402
GEOGRAPHIC DIVISION						
Northeast	371	407	280	300	159	160
New England	172	191	123	147	75	84
Middle Atlantic	332	357	252	260	142	137
Midwest	355	358	295	305	178	164
East North Central	271	301	237	244	130	141
West North Central	218	196	170	169	122	86
South	624	613	455	436	229	250
South Atlantic	489	484	351	342	182	196
East South Central	199	200	149	131	75	81
West South Central	304	295	243	233	138	128
West	467	487	348	353	194	208
Mountain	214	198	163	160	88	116
Pacific	412	445	306	313	172	173
COUNTY TYPE						
Large Metro	1,169	1,229	637	616	311	317
Small Metro	728	865	380	469	197	237
250K – 1 Mil. Pop.	596	721	309	394	163	209
< 250K Pop.	446	510	238	261	120	115
Nonmetro	569	499	318	314	164	151
Urbanized	326	336	155	227	85	118
Less Urbanized	440	374	268	223	133	93
Completely Rural	245	218	126	110	54	28

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.75D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.49	0.52	0.37	0.38	0.21	0.21
GEOGRAPHIC DIVISION						
Northeast	1.05	1.15	0.79	0.84	0.45	0.45
New England	1.84	2.04	1.32	1.58	0.81	0.90
Middle Atlantic	1.27	1.36	0.96	0.99	0.54	0.52
Midwest	0.85	0.86	0.71	0.73	0.43	0.39
East North Central	0.93	1.03	0.81	0.83	0.44	0.48
West North Central	1.75	1.56	1.37	1.34	0.98	0.68
South	0.92	0.89	0.67	0.63	0.34	0.36
South Atlantic	1.34	1.31	0.96	0.92	0.50	0.53
East South Central	1.77	1.76	1.32	1.15	0.67	0.71
West South Central	1.49	1.42	1.19	1.13	0.67	0.62
West	1.10	1.12	0.82	0.81	0.46	0.48
Mountain	1.68	1.51	1.28	1.22	0.69	0.88
Pacific	1.38	1.47	1.03	1.04	0.58	0.57
COUNTY TYPE						
Large Metro	0.67	0.74	0.54	0.53	0.29	0.31
Small Metro	0.89	0.86	0.66	0.67	0.37	0.38
250K – 1 Mil. Pop.	1.01	1.03	0.79	0.79	0.46	0.49
< 250K Pop.	1.69	1.46	1.18	1.17	0.64	0.58
Nonmetro	1.06	1.05	0.80	0.83	0.47	0.45
Urbanized	1.64	1.76	1.09	1.32	0.64	0.83
Less Urbanized	1.58	1.50	1.21	1.19	0.71	0.57
Completely Rural	3.44	3.25	2.81	2.28	1.36	0.73

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.75P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 26 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.4951		0.3048		0.6901	
GEOGRAPHIC DIVISION						
Northeast	0.9220		0.9540		0.2776	
New England	0.3614		0.9646		0.9865	
Middle Atlantic	0.7165		0.9669		0.2089	
Midwest	0.1611		0.2014		0.7681	
East North Central	0.3872		0.5034		0.5193	
West North Central	0.2139		0.1978		0.6962	
South	0.7498		0.7741		0.6709	
South Atlantic	0.2799		0.3487		0.3842	
East South Central	0.5206		0.9652		0.0569	
West South Central	0.0753		0.3201		0.1320	
West	0.5055		0.5341		0.3820	
Mountain	0.2853		0.4946		0.4199	
Pacific	0.7896		0.7346		0.5927	
COUNTY TYPE						
Large Metro	0.7959		0.9503		0.7419	
Small Metro	0.1679		0.2200		0.2130	
250K – 1 Mil. Pop.	0.6472		0.3638		0.1259	
< 250K Pop.	0.1132		0.4144		0.8621	
Nonmetro	0.4501		0.3486		0.9599	
Urbanized	0.1721		0.0044		0.0183	
Less Urbanized	0.0717		0.6106		0.2103	
Completely Rural	0.3222		0.2743		0.0455	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.76C (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	954	1,011	742	766	425	426
GEOGRAPHIC DIVISION						
Northeast	390	415	302	317	171	173
New England	183	192	134	152	80	90
Middle Atlantic	347	365	270	277	153	148
Midwest	367	373	317	326	193	173
East North Central	287	311	259	254	144	148
West North Central	224	207	182	184	128	92
South	644	630	481	463	247	262
South Atlantic	502	498	369	364	194	206
East South Central	203	209	156	142	86	82
West South Central	312	301	256	247	149	134
West	492	498	372	384	206	222
Mountain	230	203	171	172	96	120
Pacific	432	456	329	342	182	185
COUNTY TYPE						
Large Metro	1,233	1,287	705	674	337	339
Small Metro	780	916	419	514	221	259
250K – 1 Mil. Pop.	640	774	343	427	184	226
< 250K Pop.	480	536	263	298	134	129
Nonmetro	608	540	355	346	184	162
Urbanized	363	375	188	256	100	129
Less Urbanized	477	407	294	248	144	100
Completely Rural	254	234	133	123	58	32

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.76D (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.43	0.45	0.34	0.34	0.19	0.19
GEOGRAPHIC DIVISION						
Northeast	0.94	1.00	0.73	0.77	0.41	0.42
New England	1.68	1.77	1.23	1.40	0.74	0.83
Middle Atlantic	1.14	1.20	0.89	0.91	0.50	0.49
Midwest	0.75	0.76	0.65	0.66	0.39	0.35
East North Central	0.84	0.91	0.75	0.74	0.42	0.43
West North Central	1.52	1.39	1.24	1.24	0.87	0.62
South	0.81	0.78	0.60	0.57	0.31	0.32
South Atlantic	1.18	1.16	0.87	0.85	0.46	0.48
East South Central	1.55	1.58	1.19	1.07	0.65	0.62
West South Central	1.28	1.22	1.05	1.00	0.61	0.55
West	0.98	0.97	0.74	0.75	0.41	0.43
Mountain	1.52	1.31	1.13	1.11	0.64	0.78
Pacific	1.22	1.28	0.93	0.96	0.51	0.52
COUNTY TYPE						
Large Metro	0.60	0.65	0.50	0.48	0.27	0.28
Small Metro	0.79	0.73	0.59	0.59	0.34	0.34
250K – 1 Mil. Pop.	0.88	0.89	0.70	0.69	0.43	0.44
< 250K Pop.	1.50	1.26	1.07	1.08	0.59	0.55
Nonmetro	0.95	0.95	0.75	0.76	0.45	0.40
Urbanized	1.48	1.55	1.07	1.17	0.62	0.73
Less Urbanized	1.43	1.38	1.09	1.11	0.65	0.53
Completely Rural	3.01	3.03	2.53	2.20	1.25	0.69

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 2.76P (REVISED) Alcohol Use, Binge Alcohol Use, and Heavy Alcohol Use in the Past Month among Persons Aged 18 or Older, by Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Geographic Characteristic	TYPE OF ALCOHOL USE					
	Alcohol Use		Binge Alcohol Use		Heavy Alcohol Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.6451		0.4266		0.9082	
GEOGRAPHIC DIVISION						
Northeast	0.9352		0.7200		0.1126	
New England	0.4091		0.8790		0.6655	
Middle Atlantic	0.6066		0.7454		0.1130	
Midwest	0.2698		0.3611		0.9331	
East North Central	0.4703		0.6820		0.6196	
West North Central	0.3607		0.3116		0.4014	
South	0.8900		0.6341		0.8850	
South Atlantic	0.1683		0.2275		0.4105	
East South Central	0.4011		0.7682		0.2615	
West South Central	0.0827		0.4093		0.1281	
West	0.7200		0.5891		0.3624	
Mountain	0.3809		0.5809		0.4007	
Pacific	0.9914		0.7492		0.5744	
COUNTY TYPE						
Large Metro	0.8980		0.9822		0.5923	
Small Metro	0.2598		0.4025		0.4381	
250K – 1 Mil. Pop.	0.8241		0.6077		0.2177	
< 250K Pop.	0.1430		0.4864		0.6118	
Nonmetro	0.4670		0.3096		0.7395	
Urbanized	0.1984		0.0043		0.0453	
Less Urbanized	0.0925		0.7528		0.1741	
Completely Rural	0.2799		0.1829		0.0081	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.