

Table 3.22C Past Month Use of Selected Substances, by Measures of Youths' Exposure to Prevention Programs and Messages and Youths' Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Numbers in Thousands, 2007

| Prevention Programs/Messages and Delinquency ¹ | SUBSTANCE USE IN THE PAST MONTH | | | | | | | |
|--|---------------------------------|----|-----------|----|------------|----|----------------------------|----|
| | Illicit Drugs ² | | Marijuana | | Cigarettes | | Binge Alcohol ³ | |
| | Yes | No | Yes | No | Yes | No | Yes | No |
| EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR | | | | | | | | |
| Participated in Problem-Solving, Communication, or Self-Esteem Group | 28 | 66 | 24 | 53 | 29 | 66 | 28 | 66 |
| Participated in Violence Prevention Program | 25 | 67 | 20 | 54 | 24 | 67 | 24 | 67 |
| Participated in Pregnancy or Sexually Transmitted Disease Prevention Program | 25 | 65 | 19 | 54 | 23 | 67 | 21 | 68 |
| Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School | 24 | 67 | 20 | 54 | 21 | 67 | 22 | 69 |
| Participated in Program to Help Deal with Drug or Alcohol Use | 20 | 68 | 19 | 53 | 19 | 68 | 19 | 69 |
| Saw or Heard Drug or Alcohol Prevention Message from Sources Outside School | 62 | 37 | 49 | 30 | 63 | 33 | 63 | 33 |
| Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use | 50 | 50 | 41 | 39 | 51 | 45 | 53 | 44 |
| DELINQUENT BEHAVIOR IN PAST YEAR⁴ | | | | | | | | |
| Got into a Serious Fight at School or Work | 43 | 54 | 35 | 43 | 40 | 56 | 42 | 56 |
| Took Part in a Group-Against-Group Fight | 38 | 58 | 31 | 47 | 37 | 60 | 40 | 60 |
| Attacked Someone with the Intent to Harm or Seriously Hurt Them | 32 | 63 | 25 | 52 | 28 | 66 | 28 | 65 |
| Carried a Handgun | 21 | 69 | 20 | 55 | 22 | 67 | 21 | 68 |
| Sold Illegal Drugs | 30 | 63 | 29 | 49 | 27 | 66 | 25 | 67 |
| Stole or Tried to Steal Anything Worth More Than \$50 | 29 | 64 | 26 | 50 | 27 | 67 | 26 | 68 |

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as "Yes" if they indicated participating in a particular behavior at least one time and "No" if they did not.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007.

Table 3.22D Past Month Use of Selected Substances, by Measures of Youths' Exposure to Prevention Programs and Messages and Youths' Delinquency in the Past Year among Persons Aged 12 to 17: Standard Errors of Percentages, 2007

| Prevention Programs/Messages and Delinquency ¹ | SUBSTANCE USE IN THE PAST MONTH | | | | | | | |
|--|---------------------------------|------|-----------|------|------------|------|----------------------------|------|
| | Illicit Drugs ² | | Marijuana | | Cigarettes | | Binge Alcohol ³ | |
| | Yes | No | Yes | No | Yes | No | Yes | No |
| EXPOSURE TO PREVENTION PROGRAMS AND MESSAGES IN PAST YEAR | | | | | | | | |
| Participated in Problem-Solving, Communication, or Self-Esteem Group | 0.49 | 0.32 | 0.42 | 0.27 | 0.50 | 0.32 | 0.49 | 0.32 |
| Participated in Violence Prevention Program | 0.62 | 0.30 | 0.51 | 0.25 | 0.60 | 0.29 | 0.59 | 0.29 |
| Participated in Pregnancy or Sexually Transmitted Disease Prevention Program | 0.81 | 0.28 | 0.65 | 0.24 | 0.78 | 0.29 | 0.72 | 0.29 |
| Participated in Alcohol, Tobacco, or Drug Prevention Program Outside of School | 0.80 | 0.29 | 0.67 | 0.23 | 0.70 | 0.28 | 0.72 | 0.29 |
| Participated in Program to Help Deal with Drug or Alcohol Use | 1.58 | 0.27 | 1.46 | 0.22 | 1.47 | 0.27 | 1.51 | 0.27 |
| Saw or Heard Drug or Alcohol Prevention Message from Sources Outside School | 0.30 | 0.62 | 0.24 | 0.52 | 0.30 | 0.56 | 0.30 | 0.56 |
| Talked to Parents about Dangers of Tobacco, Alcohol, or Drug Use | 0.32 | 0.47 | 0.27 | 0.38 | 0.33 | 0.43 | 0.33 | 0.42 |
| DELINQUENT BEHAVIOR IN PAST YEAR⁴ | | | | | | | | |
| Got into a Serious Fight at School or Work | 0.68 | 0.26 | 0.58 | 0.22 | 0.66 | 0.27 | 0.67 | 0.28 |
| Took Part in a Group-Against-Group Fight | 0.89 | 0.26 | 0.77 | 0.21 | 0.86 | 0.26 | 0.93 | 0.27 |
| Attacked Someone with the Intent to Harm or Seriously Hurt Them | 1.49 | 0.26 | 1.27 | 0.21 | 1.36 | 0.26 | 1.37 | 0.26 |
| Carried a Handgun | 2.17 | 0.26 | 2.12 | 0.22 | 2.30 | 0.26 | 2.16 | 0.26 |
| Sold Illegal Drugs | 2.34 | 0.24 | 2.40 | 0.19 | 2.40 | 0.25 | 2.43 | 0.26 |
| Stole or Tried to Steal Anything Worth More Than \$50 | 2.14 | 0.25 | 2.04 | 0.20 | 2.07 | 0.26 | 1.89 | 0.27 |

*Low precision; no estimate reported.

¹ Respondents with unknown data were excluded.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

⁴ Respondents were asked how many times they participated in a delinquent behavior in the past year and were categorized as "Yes" if they indicated participating in a particular behavior at least one time and "No" if they did not.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2007.