

Table 6.11C Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
Outpatient Mental Health Clinic or Center	154	137	24	24	117	100	94	89
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	194	193	44	41	126	131	127	136
Doctor's Office - Not Part of a Clinic	141	138	18	18	86	81	106	105
Outpatient Medical Clinic	84	83	12	9	52	58	64	57
Partial Day Hospital or Day Treatment Program	21	51	7	6	16	33	11	*
School or University Setting/Clinic/Center ²	12	9	10	9	6	*	*	*
Some Other Place ³	47	31	8	6	22	21	*	22

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.11D Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: Standard Errors of Percentages, 2006 and 2007

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
Outpatient Mental Health Clinic or Center	1.94	1.79	2.04	2.13	2.53	2.22	4.13	4.27
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	2.11	2.06	2.56	2.50	2.59	2.48	4.86	5.29
Doctor's Office - Not Part of a Clinic	1.77	1.80	1.62	1.67	2.05	1.95	4.39	4.82
Outpatient Medical Clinic	1.19	1.19	1.17	0.91	1.37	1.45	3.02	2.96
Partial Day Hospital or Day Treatment Program	0.31	0.76	0.67	0.62	0.43	0.89	0.54	*
School or University Setting/Clinic/Center ²	0.18	0.14	0.99	0.88	0.16	*	*	*
Some Other Place ³	0.69	0.47	0.79	0.59	0.58	0.56	*	1.16

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.11P Location of Outpatient Mental Health Treatment/Counseling among Persons Aged 18 or Older with Past Year Serious Psychological Distress Who Received Outpatient Mental Health Treatment in the Past Year, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Location of Treatment/Counseling ¹	Total		AGE GROUP					
			18-25		26-49		50 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
Outpatient Mental Health Clinic or Center	0.4205		0.8931		0.8465		0.2865	
Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or Counselor - Not Part of a Clinic	0.4291		0.2787		0.5025		0.9073	
Doctor's Office - Not Part of a Clinic	0.6963		0.6378		0.2214		0.6207	
Outpatient Medical Clinic	0.3812		0.3940		0.1662		0.9025	
Partial Day Hospital or Day Treatment Program	0.0932		0.7341		0.2136		*	
School or University Setting/Clinic/Center ²	0.1817		0.3150		*		*	
Some Other Place ³	0.6471		0.2741		0.8927		*	

*Low precision; no estimate reported.

NOTE: Outpatient Mental Health Treatment/Counseling is defined as having received outpatient care for problems with emotions, nerves, or mental health. Respondents were not to include treatment for drug or alcohol use. Respondents with unknown treatment/counseling information were excluded. Estimates were based only on responses to items in the Adult Mental Health Service Utilization module.

NOTE: Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

¹ Respondents could indicate multiple locations; thus, these response categories are not mutually exclusive.

² Respondents were permitted to specify other locations for receiving outpatient mental health treatment/counseling. This location is one of the most commonly reported other locations for receiving outpatient treatment/counseling.

³ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Treatment/Counseling were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.