

Table 6.3C Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2006 and 2007

| Drug | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--|-------|------|---|------|------|------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| ILLCIT DRUGS² | 547 | 592 | 236 | 238 | 493 | 542 |
| Marijuana and Hashish | 465 | 484 | 194 | 205 | 412 | 424 |
| Cocaine | 228 | 239 | 108 | 117 | 200 | 202 |
| Crack | 133 | 123 | 74 | 88 | 111 | 86 |
| Heroin | 85 | 52 | 35 | 35 | 80 | 38 |
| Hallucinogens | 142 | 134 | 67 | 54 | 119 | 118 |
| LSD | 55 | 53 | 17 | 16 | 51 | 50 |
| PCP | 37 | 16 | 8 | 12 | 36 | 10 |
| Ecstasy | 101 | 104 | 55 | 44 | 85 | 93 |
| Inhalants | 93 | 96 | 40 | 44 | 85 | 85 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 353 | 377 | 167 | 157 | 319 | 343 |
| Pain Relievers | 298 | 312 | 143 | 134 | 274 | 283 |
| OxyContin [®] | 90 | 93 | 40 | 66 | 80 | 64 |
| Tranquilizers | 193 | 219 | 95 | 111 | 173 | 187 |
| Stimulants ⁴ | 177 | 150 | 95 | 76 | 139 | 124 |
| Methamphetamine ⁴ | 136 | 113 | 67 | 61 | 106 | 88 |
| Sedatives | 98 | 92 | 61 | 43 | 76 | 80 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 404 | 443 | 187 | 186 | 359 | 403 |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.3D Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2006 and 2007

| Drug | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--|-------|------|---|------|------|------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| ILLCIT DRUGS² | 0.25 | 0.27 | 0.87 | 0.87 | 0.24 | 0.26 |
| Marijuana and Hashish | 0.21 | 0.22 | 0.75 | 0.78 | 0.20 | 0.21 |
| Cocaine | 0.10 | 0.11 | 0.43 | 0.46 | 0.10 | 0.10 |
| Crack | 0.06 | 0.06 | 0.30 | 0.35 | 0.06 | 0.04 |
| Heroin | 0.04 | 0.02 | 0.14 | 0.15 | 0.04 | 0.02 |
| Hallucinogens | 0.06 | 0.06 | 0.27 | 0.22 | 0.06 | 0.06 |
| LSD | 0.02 | 0.02 | 0.07 | 0.07 | 0.03 | 0.03 |
| PCP | 0.02 | 0.01 | 0.03 | 0.05 | 0.02 | 0.01 |
| Ecstasy | 0.05 | 0.05 | 0.22 | 0.18 | 0.04 | 0.05 |
| Inhalants | 0.04 | 0.04 | 0.16 | 0.18 | 0.04 | 0.04 |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.16 | 0.17 | 0.65 | 0.62 | 0.16 | 0.17 |
| Pain Relievers | 0.13 | 0.14 | 0.56 | 0.54 | 0.14 | 0.14 |
| OxyContin [®] | 0.04 | 0.04 | 0.16 | 0.27 | 0.04 | 0.03 |
| Tranquilizers | 0.09 | 0.10 | 0.38 | 0.44 | 0.09 | 0.09 |
| Stimulants ⁴ | 0.08 | 0.07 | 0.38 | 0.31 | 0.07 | 0.06 |
| Methamphetamine ⁴ | 0.06 | 0.05 | 0.27 | 0.25 | 0.05 | 0.04 |
| Sedatives | 0.04 | 0.04 | 0.24 | 0.17 | 0.04 | 0.04 |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.18 | 0.20 | 0.72 | 0.72 | 0.18 | 0.20 |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.3P Types of Illicit Drug Use in the Past Year among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

| Drug | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--|--------|------|---|------|--------|------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| ILLCIT DRUGS² | 0.8825 | | 0.4955 | | 0.7660 | |
| Marijuana and Hashish | 0.6597 | | 0.1549 | | 0.2892 | |
| Cocaine | 0.2764 | | 0.4781 | | 0.1119 | |
| Crack | 0.8534 | | 0.2604 | | 0.2696 | |
| Heroin | 0.0563 | | 0.8809 | | 0.0505 | |
| Hallucinogens | 0.3007 | | 0.7074 | | 0.3671 | |
| LSD | 0.3635 | | 0.8112 | | 0.3082 | |
| PCP | 0.2450 | | 0.3201 | | 0.1094 | |
| Ecstasy | 0.8050 | | 0.9835 | | 0.8435 | |
| Inhalants | 0.8192 | | 0.9022 | | 0.7874 | |
| Nonmedical Use of Psychotherapeutics ^{3,4} | 0.7134 | | 0.8391 | | 0.8998 | |
| Pain Relievers | 0.7301 | | 0.4638 | | 0.5582 | |
| OxyContin [®] | 0.6674 | | 0.0344 | | 0.3409 | |
| Tranquilizers | 0.5913 | | 0.3540 | | 0.8017 | |
| Stimulants ⁴ | 0.0034 | | 0.0490 | | 0.0234 | |
| Methamphetamine ⁴ | 0.0058 | | 0.3063 | | 0.0058 | |
| Sedatives | 0.5924 | | 0.5216 | | 0.8820 | |
| ILLCIT DRUGS OTHER THAN MARIJUANA² | 0.9448 | | 0.9152 | | 0.9585 | |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.4C Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Numbers in Thousands, 2006 and 2007

| Substance | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--------------------------------|-------|-------|---|------|-------|-------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| PAST YEAR | | | | | | |
| Tobacco Products ² | 865 | 870 | 357 | 355 | 958 | 1,009 |
| Cigarettes | 830 | 840 | 349 | 344 | 855 | 916 |
| Smokeless Tobacco | 320 | 303 | 108 | 87 | 314 | 300 |
| Cigars | 461 | 475 | 156 | 189 | 461 | 455 |
| Alcohol | 897 | 957 | 428 | 426 | 1,562 | 1,704 |
| PAST MONTH | | | | | | |
| Tobacco Products ² | 837 | 838 | 345 | 339 | 893 | 921 |
| Cigarettes | 796 | 802 | 336 | 328 | 798 | 835 |
| Smokeless Tobacco | 281 | 278 | 85 | 75 | 278 | 275 |
| Cigars | 347 | 331 | 107 | 117 | 338 | 320 |
| Pipe Tobacco | 164 | 169 | 52 | 97 | 160 | 140 |
| Alcohol | 955 | 1,010 | 373 | 376 | 1,385 | 1,492 |
| Binge Alcohol Use ³ | 741 | 763 | 260 | 268 | 773 | 825 |
| Heavy Alcohol Use ³ | 425 | 426 | 129 | 168 | 407 | 413 |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.4D Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: Standard Errors of Percentages, 2006 and 2007

| Substance | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--------------------------------|-------|------|---|------|------|------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| PAST YEAR | | | | | | |
| Tobacco Products ² | 0.39 | 0.39 | 1.01 | 1.06 | 0.41 | 0.41 |
| Cigarettes | 0.38 | 0.38 | 1.02 | 1.05 | 0.39 | 0.39 |
| Smokeless Tobacco | 0.14 | 0.14 | 0.42 | 0.35 | 0.16 | 0.15 |
| Cigars | 0.21 | 0.21 | 0.61 | 0.73 | 0.22 | 0.22 |
| Alcohol | 0.41 | 0.43 | 1.01 | 1.06 | 0.43 | 0.46 |
| PAST MONTH | | | | | | |
| Tobacco Products ² | 0.38 | 0.38 | 1.02 | 1.06 | 0.40 | 0.39 |
| Cigarettes | 0.36 | 0.36 | 1.03 | 1.04 | 0.37 | 0.37 |
| Smokeless Tobacco | 0.13 | 0.13 | 0.33 | 0.31 | 0.14 | 0.14 |
| Cigars | 0.16 | 0.15 | 0.42 | 0.47 | 0.16 | 0.15 |
| Pipe Tobacco | 0.07 | 0.08 | 0.21 | 0.39 | 0.08 | 0.07 |
| Alcohol | 0.43 | 0.45 | 1.09 | 1.13 | 0.46 | 0.49 |
| Binge Alcohol Use ³ | 0.34 | 0.34 | 0.91 | 0.95 | 0.35 | 0.37 |
| Heavy Alcohol Use ³ | 0.19 | 0.19 | 0.49 | 0.66 | 0.20 | 0.20 |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 6.4P Tobacco Product and Alcohol Use in the Past Year and Past Month among Persons Aged 18 or Older, by Past Year Serious Psychological Distress: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

| Substance | Total | | SERIOUS PSYCHOLOGICAL DISTRESS ¹ | | | |
|--------------------------------|--------|------|---|------|--------|------|
| | | | Yes | | No | |
| | 2006 | 2007 | 2006 | 2007 | 2006 | 2007 |
| PAST YEAR | | | | | | |
| Tobacco Products ² | 0.1786 | | 0.1774 | | 0.3909 | |
| Cigarettes | 0.2416 | | 0.1713 | | 0.5156 | |
| Smokeless Tobacco | 0.3091 | | 0.9839 | | 0.2876 | |
| Cigars | 0.4542 | | 0.3203 | | 0.2433 | |
| Alcohol | 0.6511 | | 0.6535 | | 0.7511 | |
| PAST MONTH | | | | | | |
| Tobacco Products ² | 0.0342 | | 0.1039 | | 0.1145 | |
| Cigarettes | 0.0861 | | 0.1537 | | 0.2260 | |
| Smokeless Tobacco | 0.4896 | | 0.8935 | | 0.4476 | |
| Cigars | 0.2331 | | 0.8129 | | 0.2391 | |
| Pipe Tobacco | 0.1883 | | 0.3900 | | 0.0595 | |
| Alcohol | 0.6667 | | 0.3945 | | 0.8592 | |
| Binge Alcohol Use ³ | 0.4100 | | 0.0100 | | 0.9215 | |
| Heavy Alcohol Use ³ | 0.9076 | | 0.0612 | | 0.4635 | |

*Low precision; no estimate reported.

¹ Serious Psychological Distress (SPD) is defined as having a score of 13 or higher on the K6 scale. See Section B.4.4 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco. Tobacco Product use in the past year excludes past year pipe tobacco use, but includes past month pipe tobacco use.

³ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.