

Table 7.11A Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	5,533	5,330	6,342	6,547	4,414 ^a	3,809	4,069	4,171
Marijuana and Hashish	4,536	4,268	5,037	5,071	2,992	2,782	2,246	2,327
Cocaine	1,179	932	693	611	373	263	176	270
Crack	236	204	211	135	199	107	56 ^a	164
Heroin	156 ^a	29	32	17	71	65	79	42
Hallucinogens	386	367	335	363	146	126	140	139
LSD	48	51	33	55	37	20	12	18
PCP	10	4	14	15	*	4	*	18
Ecstasy	222	181	174	180	75	92	57	51
Inhalants	128	149	165	159	227 ^b	77	241	231
Nonmedical Use of Psychotherapeutics ^{3,4}	1,916	1,780	1,868	1,884	1,503	1,324	1,807	1,908
Pain Relievers	1,399	1,452	1,386	1,348	1,004	894	1,431	1,480
OxyContin [®]	90	108	94	118	33	36	60	106
Tranquilizers	511	580	544	500	380	372	330	382
Stimulants ⁴	423	315	398	302	318	191	246	244
Methamphetamine ⁴	230	154	209	152	148	93	143	130
Sedatives	102	77	42	117	129	57	112	95
ILLICIT DRUGS OTHER THAN MARIJUANA²	2,789	2,615	2,626	2,657	1,983	1,634	2,217	2,365

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.11B Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	32.6	31.3	16.0	16.1	6.4 ^a	5.5	3.4	3.4
Marijuana and Hashish	26.8	25.1	12.7	12.4	4.4	4.0	1.9	1.9
Cocaine	7.0	5.5	1.7	1.5	0.5	0.4	0.1	0.2
Crack	1.4	1.2	0.5	0.3	0.3	0.2	0.0 ^a	0.1
Heroin	0.9 ^a	0.2	0.1	0.0	0.1	0.1	0.1	0.0
Hallucinogens	2.3	2.2	0.8	0.9	0.2	0.2	0.1	0.1
LSD	0.3	0.3	0.1	0.1	0.1	0.0	0.0	0.0
PCP	0.1	0.0	0.0	0.0	*	0.0	*	0.0
Ecstasy	1.3	1.1	0.4	0.4	0.1	0.1	0.0	0.0
Inhalants	0.8	0.9	0.4	0.4	0.3 ^b	0.1	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	11.3	10.5	4.7	4.6	2.2	1.9	1.5	1.6
Pain Relievers	8.3	8.5	3.5	3.3	1.5	1.3	1.2	1.2
OxyContin [®]	0.5	0.6	0.2	0.3	0.0	0.1	0.0	0.1
Tranquilizers	3.0	3.4	1.4	1.2	0.6	0.5	0.3	0.3
Stimulants ⁴	2.5	1.9	1.0	0.7	0.5	0.3	0.2	0.2
Methamphetamine ⁴	1.4	0.9	0.5	0.4	0.2	0.1	0.1	0.1
Sedatives	0.6	0.5	0.1	0.3	0.2	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	16.5	15.4	6.6	6.5	2.9	2.4	1.8	2.0

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.12A Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	347	355	762	704	358	296	1,014	1,053
Marijuana and Hashish	298	317	633	593	251	226	507	550
Cocaine	31	33	44	37	16	9	19	24
Crack	5	6	4	4	3	1	*	5
Heroin	6	1	4	*	2	1	4	*
Hallucinogens	56	44	53	74	18	18	54	39
LSD	11	6	11	12	3	3	7	8
PCP	5	1	3	4	*	1	4	3
Ecstasy	31	20	25	24	8	8	18	13
Inhalants	26	36	75	60	63 ^b	27	176	172
Nonmedical Use of Psychotherapeutics ^{3,4}	129	144	208	194	79	90	427	402
Pain Relievers	111	126	177	154	60	83	342	306
OxyContin [®]	6 ^a	18	8	13	3	7	11	9
Tranquilizers	31	54	46	57	15	12	34	59
Stimulants ⁴	28	26	48	36	16	9	79	56
Methamphetamine ⁴	8	13	13	6	4	2	30	13
Sedatives	5	3	8	4	4	2	28	26
ILLICIT DRUGS OTHER THAN MARIJUANA²	166	177	323	290	156	122	606	592

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.12B Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	57.6	60.1	37.9	37.9	22.3	18.8	4.8	5.0
Marijuana and Hashish	49.4	53.7	31.5	32.0	15.6	14.3	2.4	2.6
Cocaine	5.2	5.5	2.2	2.0	1.0	0.6	0.1	0.1
Crack	0.8	1.1	0.2	0.2	0.2	0.1	*	0.0
Heroin	1.0	0.2	0.2	*	0.1	0.1	0.0	*
Hallucinogens	9.2	7.5	2.7	4.0	1.1	1.2	0.3	0.2
LSD	1.9	1.0	0.5	0.7	0.2	0.2	0.0	0.0
PCP	0.8	0.1	0.2	0.2	*	0.1	0.0	0.0
Ecstasy	5.2	3.4	1.2	1.3	0.5	0.5	0.1	0.1
Inhalants	4.4	6.2	3.7	3.2	3.9 ^b	1.7	0.8	0.8
Nonmedical Use of Psychotherapeutics ^{3,4}	21.5	24.3	10.3	10.4	4.9	5.7	2.0	1.9
Pain Relievers	18.4	21.3	8.8	8.3	3.7	5.3	1.6	1.4
OxyContin [®]	1.0 ^a	3.0	0.4	0.7	0.2	0.4	0.1	0.0
Tranquilizers	5.1 ^a	9.1	2.3	3.1	0.9	0.8	0.2	0.3
Stimulants ⁴	4.6	4.4	2.4	2.0	1.0	0.6	0.4	0.3
Methamphetamine ⁴	1.3	2.2	0.6	0.3	0.2	0.1	0.1	0.1
Sedatives	0.8	0.5	0.4	0.2	0.2	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	27.6	29.9	16.1	15.6	9.7	7.8	2.9	2.8

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.13A Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	2,366	2,331	2,388	2,482	920	882	809	769
Marijuana and Hashish	1,997	2,021	1,998	2,079	745	729	582	550
Cocaine	407 ^a	309	245	199	50	38	27	26
Crack	29	24	30	14	7	9	6	10
Heroin	25 ^a	8	15	17	3	13	11	9
Hallucinogens	263	235	198	171	70 ^a	36	32	33
LSD	33	41	12	19	13	1	1	10
PCP	5	4	7	2	*	*	2	9
Ecstasy	146 ^a	91	127	101	40	21	12	16
Inhalants	42	43	44	49	16	12	27	25
Nonmedical Use of Psychotherapeutics ^{3,4}	847	747	753	698	249	239	265	266
Pain Relievers	655	595	584	538	184	168	197	216
OxyContin [®]	54	58	60	59	11	19	13	20
Tranquilizers	273	269	230 ^a	167	79	76	64	59
Stimulants ⁴	218 ^a	146	167	120	35	49	37	37
Methamphetamine ⁴	69	44	82 ^a	38	12	21	19	12
Sedatives	26	38	7 ^a	21	11	3	11	10
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,209	1,067	1,008	957	350	295	337	324

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.13B Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	46.5	48.4	27.4	28.0	14.2	13.9	6.5	6.1
Marijuana and Hashish	39.2	41.9	22.9	23.5	11.5	11.5	4.7	4.3
Cocaine	8.0 ^a	6.4	2.8	2.2	0.8	0.6	0.2	0.2
Crack	0.6	0.5	0.3	0.2	0.1	0.1	0.1	0.1
Heroin	0.5	0.2	0.2	0.2	0.0	0.2	0.1	0.1
Hallucinogens	5.2	4.9	2.3	1.9	1.1 ^a	0.6	0.3	0.3
LSD	0.6	0.9	0.1	0.2	0.2	0.0	0.0	0.1
PCP	0.1	0.1	0.1	0.0	*	*	0.0	0.1
Ecstasy	2.9	1.9	1.5	1.1	0.6	0.3	0.1	0.1
Inhalants	0.8	0.9	0.5	0.6	0.2	0.2	0.2	0.2
Nonmedical Use of Psychotherapeutics ^{3,4}	16.6	15.5	8.6	7.9	3.9	3.8	2.1	2.1
Pain Relievers	12.9	12.3	6.7	6.1	2.8	2.6	1.6	1.7
OxyContin [®]	1.1	1.2	0.7	0.7	0.2	0.3	0.1	0.2
Tranquilizers	5.4	5.6	2.6 ^a	1.9	1.2	1.2	0.5	0.5
Stimulants ⁴	4.3	3.0	1.9	1.4	0.5	0.8	0.3	0.3
Methamphetamine ⁴	1.4	0.9	0.9 ^b	0.4	0.2	0.3	0.1	0.1
Sedatives	0.5	0.8	0.1 ^a	0.2	0.2	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	23.7	22.1	11.6	10.8	5.4	4.6	2.7	2.5

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.14A Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	2,820	2,643	3,192	3,361	3,136	2,630	2,246	2,349
Marijuana and Hashish	2,241	1,931	2,406	2,399	1,996	1,828	1,158	1,227
Cocaine	740	590	404	374	307	215	130	220
Crack	203	174	177	117	189	97	49	149
Heroin	125	19	12	*	67	50	64	32
Hallucinogens	67	88	83	118	58	72	54	68
LSD	4	4	10	24	21	16	5	*
PCP	*	*	3	9	*	*	*	6
Ecstasy	45	69	22	55	26	63	27	22
Inhalants	60	69	46	51	148 ^a	38	37	34
Nonmedical Use of Psychotherapeutics ^{3,4}	940	889	907	992	1,174	994	1,115	1,241
Pain Relievers	633	731	624	656	760	643	892	957
OxyContin [®]	30	33	26	45	18	11	35	78
Tranquilizers	207	258	268	276	286	285	232	263
Stimulants ⁴	177	143	183	145	268	132	130	152
Methamphetamine ⁴	153	98	114	108	132	70	95	104
Sedatives	71	36	27	93	114	52	73	59
ILLCIT DRUGS OTHER THAN MARIJUANA²	1,414	1,371	1,295	1,409	1,477	1,217	1,274	1,449

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.14B Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS²	25.1	22.8	11.0	11.2	5.2 ^a	4.3	2.6	2.7
Marijuana and Hashish	19.9	16.6	8.3	8.0	3.3	3.0	1.3	1.4
Cocaine	6.6	5.1	1.4	1.2	0.5	0.4	0.1	0.3
Crack	1.8	1.5	0.6	0.4	0.3	0.2	0.1	0.2
Heroin	1.1	0.2	0.0	*	0.1	0.1	0.1	0.0
Hallucinogens	0.6	0.8	0.3	0.4	0.1	0.1	0.1	0.1
LSD	0.0	0.0	0.0	0.1	0.0	0.0	0.0	*
PCP	*	*	0.0	0.0	*	*	*	0.0
Ecstasy	0.4	0.6	0.1	0.2	0.0	0.1	0.0	0.0
Inhalants	0.5	0.6	0.2	0.2	0.2 ^a	0.1	0.0	0.0
Nonmedical Use of Psychotherapeutics ^{3,4}	8.4	7.7	3.1	3.3	1.9	1.6	1.3	1.4
Pain Relievers	5.6	6.3	2.2	2.2	1.3	1.1	1.0	1.1
OxyContin [®]	0.3	0.3	0.1	0.2	0.0	0.0	0.0	0.1
Tranquilizers	1.8	2.2	0.9	0.9	0.5	0.5	0.3	0.3
Stimulants ⁴	1.6	1.2	0.6	0.5	0.4	0.2	0.1	0.2
Methamphetamine ⁴	1.4	0.8	0.4	0.4	0.2	0.1	0.1	0.1
Sedatives	0.6	0.3	0.1	0.3	0.2	0.1	0.1	0.1
ILLICIT DRUGS OTHER THAN MARIJUANA²	12.6	11.8	4.5	4.7	2.4	2.0	1.5	1.7

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.15A Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Numbers in Thousands, 2006 and 2007

Drug	Total		UNDERAGE AND LEGAL AGE DRINKING GROUP			
			12-20		21 or Older	
	2006	2007	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month						
ILLCIT DRUGS^{1,2}	16,288	15,686	3,871	3,751	12,417	11,935
Marijuana or Hashish	12,566	12,122	3,244	3,224	9,322	8,897
Cocaine (Including Crack)	2,246 ^a	1,806	416 ^b	300	1,829	1,505
Heroin	259	111	37	16	223	95
Hallucinogens	866	856	423	360	444	496
Inhalants	520	385	234	180	286	205
Pain Relievers ³	3,789	3,694	957	914	2,832	2,780
Tranquilizers ³	1,436	1,453	329	315	1,106	1,138
Stimulants ^{3,4}	1,138 ^b	809	289 ^b	190	850	619
Methamphetamine ⁴	588 ^a	400	101 ^a	53	487	347
Sedatives ³	272	251	39	28	233	224
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	7,398	6,905	1,812 ^b	1,562	5,586	5,343
Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month						
ILLCIT DRUGS^{1,2}	7,180	6,951	1,701	1,716	5,480	5,235
Marijuana or Hashish	6,138	5,983	1,594	1,616	4,544	4,367
Cocaine (Including Crack)	831	614	70	71	761	543
Heroin	91	38	2	6	89	32
Hallucinogens	78	99	21	35	58	63
Inhalants	31	16	13	12	17	*
Pain Relievers ³	498	519	128 ^a	84	371	435
Tranquilizers ³	106	89	26	19	80	69
Stimulants ³	43	39	15	11	28	29
Methamphetamine ⁶	162	123	22	11	140	111
Sedatives ³	19	49	1	2	18	47
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	1,616	1,386	267	217	1,349	1,170

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.15B Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Percentages, 2006 and 2007

Drug	Total		UNDERAGE AND LEGAL AGE DRINKING GROUP			
			12-20		21 or Older	
	2006	2007	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month						
ILLICIT DRUGS^{1,2}	13.0	12.4	35.8	35.0	10.8	10.3
Marijuana or Hashish	10.0	9.6	30.0	30.0	8.1	7.7
Cocaine (Including Crack)	1.8 ^a	1.4	3.8 ^b	2.8	1.6	1.3
Heroin	0.2	0.1	0.3	0.1	0.2	0.1
Hallucinogens	0.7	0.7	3.9	3.4	0.4	0.4
Inhalants	0.4	0.3	2.2	1.7	0.3	0.2
Pain Relievers ³	3.0	2.9	8.8	8.5	2.5	2.4
Tranquilizers ³	1.1	1.1	3.0	2.9	1.0	1.0
Stimulants ^{3,4}	0.9 ^b	0.6	2.7 ^b	1.8	0.7 ^a	0.5
Methamphetamine ⁴	0.5 ^a	0.3	0.9 ^a	0.5	0.4	0.3
Sedatives ³	0.2	0.2	0.4	0.3	0.2	0.2
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	5.9	5.4	16.7 ^b	14.6	4.9	4.6
Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month						
ILLICIT DRUGS^{1,2}	5.7	5.5	16.0	16.3	4.8	4.5
Marijuana or Hashish	4.9	4.7	15.0	15.3	4.0	3.8
Cocaine (Including Crack)	0.7	0.5	0.7	0.7	0.7	0.5
Heroin	0.1	0.0	0.0	0.1	0.1	0.0
Hallucinogens	0.1	0.1	0.2	0.3	0.1	0.1
Inhalants	0.0	0.0	0.1	0.1	0.0	*
Pain Relievers ³	0.4	0.4	1.2 ^a	0.8	0.3	0.4
Tranquilizers ³	0.1	0.1	0.2	0.2	0.1	0.1
Stimulants ³	0.0	0.0	0.1	0.1	0.0	0.0
Methamphetamine ⁶	0.1	0.1	0.2	0.1	0.1	0.1
Sedatives ³	0.0	0.0	0.0	0.0	0.0	0.0
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	1.3	1.1	2.5	2.0	1.2	1.0

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.16A Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Drug	AGE GROUP					
	12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007
	Used Specified Drug Anytime in the Past Month					
ILLCIT DRUGS^{1,2}	1,467	1,356	5,674	5,695	9,148	8,635
Marijuana or Hashish	1,182	1,136	4,740	4,829	6,644	6,157
Cocaine (Including Crack)	91	79	703 ^b	547	1,452	1,180
Heroin	12 ^a	3	43	38	204	70
Hallucinogens	127	136	531	442	208	278
Inhalants	164	123	102	105	254	157
Pain Relievers ³	348	363	1,423	1,301	2,017	2,030
Tranquilizers ³	92	123	583	512	761	818
Stimulants ^{3,4}	91	72	420 ^a	316	628	421
Methamphetamine ⁴	24	20	164 ^a	103	400	276
Sedatives ³	16	9	43	62	213	180
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	645	589	2,567 ^a	2,319	4,186	3,998
Drug⁵	Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month					
ILLCIT DRUGS^{1,2}	566	580	2,653	2,739	3,961	3,632
Marijuana or Hashish	536	539	2,429	2,535	3,173	2,909
Cocaine (Including Crack)	18	17	163	134	650	463
Heroin	1	2	2	11	88	25
Hallucinogens	10	20	26	41	43	38
Inhalants	12	8	4	6	15	*
Pain Relievers ³	32	33	212	198	254	288
Tranquilizers ³	7	12	39	46	60	31
Stimulants ³	4	3	24	22	15	14
Methamphetamine ⁶	4	3	35	32	123	87
Sedatives ³	1	2	1 ^a	13	16	33
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	75	79	460	420	1,080	888

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.16B Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Drug	AGE GROUP					
	12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month						
ILLCIT DRUGS^{1,2}	34.7	33.7	28.0	28.4	9.1	8.4
Marijuana or Hashish	28.0	28.3	23.4	24.1	6.6	6.0
Cocaine (Including Crack)	2.2	2.0	3.5 ^b	2.7	1.4	1.1
Heroin	0.3	0.1	0.2	0.2	0.2	0.1
Hallucinogens	3.0	3.4	2.6	2.2	0.2	0.3
Inhalants	3.9	3.0	0.5	0.5	0.3	0.2
Pain Relievers ³	8.2	9.0	7.0	6.5	2.0	2.0
Tranquilizers ³	2.2	3.1	2.9	2.6	0.8	0.8
Stimulants ^{3,4}	2.2	1.8	2.1 ^a	1.6	0.6	0.4
Methamphetamine ⁴	0.6	0.5	0.8 ^a	0.5	0.4	0.3
Sedatives ³	0.4	0.2	0.2	0.3	0.2	0.2
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	15.3	14.6	12.7	11.6	4.2	3.9
Drug ⁵	Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month					
	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS^{1,2}	13.7	14.8	13.2	13.8	3.9	3.5
Marijuana or Hashish	13.0	13.7	12.1	12.7	3.2	2.8
Cocaine (Including Crack)	0.4	0.4	0.8	0.7	0.6	0.5
Heroin	0.0	0.1	0.0	0.1	0.1	0.0
Hallucinogens	0.2	0.5	0.1	0.2	0.0	0.0
Inhalants	0.3	0.2	0.0	0.0	0.0	*
Pain Relievers ³	0.8	0.8	1.1	1.0	0.3	0.3
Tranquilizers ³	0.2	0.3	0.2	0.2	0.1	0.0
Stimulants ³	0.1	0.1	0.1	0.1	0.0	0.0
Methamphetamine ⁶	0.1	0.1	0.2	0.2	0.1	0.1
Sedatives ³	0.0	0.1	0.0 ^a	0.1	0.0	0.0
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	1.8	2.0	2.3	2.1	1.1	0.9

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.