

Table 7.22A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	11,307	10,870
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3,093	3,215	5,138	4,836
Cigars	7,699	7,316	6,010	5,947
Pipe Tobacco	1,272	1,138	1,049	907
ALCOHOL	40,846	40,166	84,462	86,594
Binge Alcohol Use ²	26,851	27,031	29,724	30,747
Heavy Alcohol Use ²	9,874	9,877	7,072	7,134

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.22B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	6.1	5.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	5.0	5.4	2.8	2.6
Cigars	12.5	12.2	3.3	3.2
Pipe Tobacco	2.1	1.9	0.6	0.5
ALCOHOL	66.3	66.9	45.8	46.1
Binge Alcohol Use ²	43.6	45.0	16.1	16.4
Heavy Alcohol Use ²	16.0	16.4	3.8	3.8

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.23A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	641	646
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	299	324	301	293
Cigars	677	699	369	371
Pipe Tobacco	111	135	57	46
ALCOHOL	1,619	1,512	2,603	2,509
Binge Alcohol Use ²	1,265	1,220	1,349	1,227
Heavy Alcohol Use ²	382	422	221	169

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.23B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	2.8	2.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	11.4	13.0	1.3	1.3
Cigars	25.7	28.2	1.6	1.6
Pipe Tobacco	4.2	5.5	0.3	0.2
ALCOHOL	61.5	60.9	11.4	11.0
Binge Alcohol Use ²	48.0	49.2	5.9	5.4
Heavy Alcohol Use ²	14.5	17.0	1.0	0.7

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.24A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	1,801	1,859
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,116	1,125	591	596
Cigars	2,671	2,554	1,301	1,319
Pipe Tobacco	323	279	107	115
ALCOHOL	9,987 ^a	9,361	10,295	10,673
Binge Alcohol Use ²	7,895	7,446	5,919	6,228
Heavy Alcohol Use ²	3,459 ^a	3,140	1,633	1,680

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.24B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	8.9	8.9
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	8.9	9.5	2.9	2.9
Cigars	21.2	21.6	6.5	6.3
Pipe Tobacco	2.6	2.4	0.5	0.6
ALCOHOL	79.4	79.1	51.1	51.1
Binge Alcohol Use ²	62.7	62.9	29.4	29.8
Heavy Alcohol Use ²	27.5	26.5	8.1	8.0

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.25A Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	8,865	8,365
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	1,678	1,766	4,246	3,947
Cigars	4,350	4,064	4,340	4,256
Pipe Tobacco	838	724	885	746
ALCOHOL	29,240	29,293	71,564	73,412
Binge Alcohol Use ²	17,691	18,364	22,456	23,292
Heavy Alcohol Use ²	6,034	6,315	5,217	5,285

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.25B Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	6.3	5.8
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	3.6	3.9	3.0	2.7
Cigars	9.4	8.9	3.1	3.0
Pipe Tobacco	1.8	1.6	0.6	0.5
ALCOHOL	63.1	64.0	50.6	50.9
Binge Alcohol Use ²	38.2	40.1	15.9	16.2
Heavy Alcohol Use ²	13.0	13.8	3.7	3.7

*Low precision; no estimate reported.

N/A: Not applicable.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.