

Table 7.26A Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	11,603	11,554	20,317	20,350	16,946	16,102	24,006	22,933
Cigarettes	9,874	9,877	16,976	17,154	13,996	13,135	20,719	19,903
Smokeless Tobacco	1,937	2,099	2,224	2,291	1,441	1,408	2,628	2,253
Cigars	3,170	2,983	4,771	4,617	3,182	2,964	2,584	2,700
Pipe Tobacco	464	396	575	520	539	533	744	597

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.26B Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	68.5	67.9	51.3	49.9	24.7	23.3	19.9	18.9
Cigarettes	58.3	58.1	42.8	42.1	20.4	19.0	17.2	16.4
Smokeless Tobacco	11.4	12.3	5.6	5.6	2.1	2.0	2.2	1.9
Cigars	18.7	17.5	12.0	11.3	4.6	4.3	2.1	2.2
Pipe Tobacco	2.7	2.3	1.5	1.3	0.8	0.8	0.6	0.5

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.27A Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	454	490	1,058	977	454	380	1,309	1,281
Cigarettes	382	422	883	798	354	291	1,015	970
Smokeless Tobacco	134	129	178	183	63	56	225	248
Cigars	215	216	361	357	138	105	333	393
Pipe Tobacco	49	49	56	55	21	22	43	55

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.27B Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	75.3 ^a	82.9	52.6	52.6	28.2	24.2	6.2	6.0
Cigarettes	63.3 ^a	71.5	43.9	43.0	22.0	18.5	4.8	4.6
Smokeless Tobacco	22.3	21.8	8.8	9.9	3.9	3.6	1.1	1.2
Cigars	35.7	36.5	17.9	19.2	8.6	6.7	1.6	1.8
Pipe Tobacco	8.1	8.4	2.8	3.0	1.3	1.4	0.2	0.3

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.28A Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	3,951	3,643	5,065	5,004	2,409	2,205	2,958	2,841
Cigarettes	3,459 ^a	3,140	4,436	4,307	2,092	1,915	2,595	2,474
Smokeless Tobacco	764	758	541	571	161	116	241	276
Cigars	1,375	1,260	1,385	1,477	578	528	634	608
Pipe Tobacco	155	119	155	168	46	53	74	54

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.28B Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	77.6	75.6	58.1	56.5	37.2	34.7	23.7	22.4
Cigarettes	67.9	65.1	50.9	48.6	32.3	30.1	20.8	19.5
Smokeless Tobacco	15.0	15.7	6.2	6.4	2.5	1.8	1.9	2.2
Cigars	27.0	26.1	15.9	16.7	8.9	8.3	5.1	4.8
Pipe Tobacco	3.0	2.5	1.8	1.9	0.7	0.8	0.6	0.4

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.29A Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	7,198	7,420	14,194	14,369	14,083	13,517	19,739	18,811
Cigarettes	6,034	6,315	11,657	12,049	11,549	10,929	17,109	16,459
Smokeless Tobacco	1,039	1,212	1,506	1,537	1,217	1,235	2,162	1,729
Cigars	1,580	1,507	3,026	2,783	2,466	2,331	1,617	1,699
Pipe Tobacco	260	227	364	297	472	459	627	488

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.29B Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	64.0	64.0	49.1	47.8	23.2	22.1	22.7	21.6
Cigarettes	53.6	54.4	40.3	40.1	19.0	17.9	19.6	18.9
Smokeless Tobacco	9.2	10.4	5.2	5.1	2.0	2.0	2.5	2.0
Cigars	14.0	13.0	10.5	9.3	4.1	3.8	1.9	1.9
Pipe Tobacco	2.3	2.0	1.3	1.0	0.8	0.8	0.7	0.6

*Low precision; no estimate reported.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.