

Table 7.45A Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	2,214	2,061	313	260	522	506	1,378	1,295
From More Than One Doctor	190	292	42	25	61	47	86 ^a	219
Wrote Fake Prescription	10	26	3	2	8	7	*	17
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	52	29	8	8	5	14	39	7
From Friend or Relative for Free	6,472	6,427	685	637	2,128	2,095	3,659	3,694
Bought from Friend or Relative	1,083	1,012	125	123	496	455	461	434
Took from Friend or Relative without Asking	632	595	151	148	226	205	256	242
Bought from Drug Dealer or Other Stranger	455	464	63	68	217	208	175	188
Bought on the Internet	11	52	1	3	4	9	6	39
Some Other Way	503	422	112	98	130	141	261	183

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.45B Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	19.1	18.1	20.8	18.9	13.8	13.7	21.8	20.5
From More Than One Doctor	1.6	2.6	2.8	1.8	1.6	1.3	1.4 ^a	3.5
Wrote Fake Prescription	0.1	0.2	0.2	0.1	0.2	0.2	*	0.3
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.3	0.5	0.6	0.1	0.4	0.6	0.1
From Friend or Relative for Free	55.7	56.5	45.6	46.4	56.0	56.8	57.9	58.5
Bought from Friend or Relative	9.3	8.9	8.3	8.9	13.1	12.3	7.3	6.9
Took from Friend or Relative without Asking	5.4	5.2	10.0	10.8	5.9	5.6	4.1	3.8
Bought from Drug Dealer or Other Stranger	3.9	4.1	4.2	5.0	5.7	5.6	2.8	3.0
Bought on the Internet	0.1	0.5	0.1	0.2	0.1	0.3	0.1	0.6
Some Other Way	4.3	3.7	7.4	7.2	3.4	3.8	4.1	2.9

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.46A Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	4,678	4,672	350	309	1,443	1,372	2,885	2,991
From More Than One Doctor	192	165	25	26	54	56	113	82
Wrote Fake Prescription	10	6	*	1	4	6	6	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	20	48	6	7	5	6	9	36
From Friend or Relative for Free	424	379	69	71	166	147	189	161
Bought from Friend or Relative	214	274	35	34	118	139	61	101
Took from Friend or Relative without Asking	69	65	41	32	29	17	*	16
Bought from Drug Dealer or Other Stranger	91	106	15	29	42	48	34	29
Bought on the Internet	*	4	*	1	*	3	*	*
Some Other Way	96	47	29	25	30 ^a	13	37	10

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.46B Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	80.7	81.0	61.4	57.7	76.3	76.0	86.5	87.3
From More Than One Doctor	3.3	2.9	4.4	4.9	2.9	3.1	3.4	2.4
Wrote Fake Prescription	0.2	0.1	*	0.1	0.2	0.3	0.2	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3	0.8	1.1	1.3	0.2	0.3	0.3	1.0
From Friend or Relative for Free	7.3	6.6	12.1	13.3	8.8	8.2	5.7	4.7
Bought from Friend or Relative	3.7	4.8	6.2	6.4	6.2	7.7	1.8	3.0
Took from Friend or Relative without Asking	1.2	1.1	7.2	6.1	1.5	0.9	*	0.5
Bought from Drug Dealer or Other Stranger	1.6	1.8	2.6	5.5	2.2	2.6	1.0	0.9
Bought on the Internet	*	0.1	*	0.1	*	0.2	*	*
Some Other Way	1.7	0.8	5.0	4.7	1.6	0.7	1.1	0.3

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.47A Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1,212	1,278	139	138	291	280	782	861
From More Than One Doctor	192	295	26	18	60	70	105	207
Wrote Fake Prescription	26	31	3	2	6	11	*	17
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	56	43	5	12	12	15	39	16
From Friend or Relative for Free	2,756	2,671	294	273	930	859	1,532	1,539
Bought from Friend or Relative	1,017	1,008	118	114	503	446	396	449
Took from Friend or Relative without Asking	456	513	97	96	198	172	161	245
Bought from Drug Dealer or Other Stranger	533	480	65	65	246	227	222	189
Bought on the Internet	25	53	2	6	7	15	16	32
Some Other Way	317	253	77	61	79	80	161	113

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.47B Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	25.3	27.1	24.7	25.2	19.0	20.1	29.0	31.0
From More Than One Doctor	4.0	6.3	4.7	3.4	3.9	5.0	3.9	7.5
Wrote Fake Prescription	0.5	0.7	0.6	0.4	0.4	0.8	*	0.6
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.2	0.9	0.8	2.1	0.8	1.1	1.5	0.6
From Friend or Relative for Free	57.5	56.6	52.3	49.8	60.8	61.5	56.7	55.4
Bought from Friend or Relative	21.2	21.4	20.9	20.8	32.9	32.0	14.7	16.2
Took from Friend or Relative without Asking	9.5	10.9	17.2	17.5	13.0	12.4	6.0	8.8
Bought from Drug Dealer or Other Stranger	11.1	10.2	11.7	11.8	16.1	16.2	8.2	6.8
Bought on the Internet	0.5	1.1	0.3	1.1	0.5	1.1	0.6	1.2
Some Other Way	6.8	5.5	14.4	11.5	5.3	5.9	6.1	4.2

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.48A Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	2,031	1,828	144	130	625	576	1,262	1,122
From More Than One Doctor	244	222	24	31	67	90	153	101
Wrote Fake Prescription	52	28	4	*	0 ^a	13	47	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	52	62	*	10	8	16	43	*
From Friend or Relative for Free	382	412	60	50	163	143	159	219
Bought from Friend or Relative	270	351	45	35	133	157	92	159
Took from Friend or Relative without Asking	120	78	46	32	44	36	31	10
Bought from Drug Dealer or Other Stranger	160	167	18	33	85	95	57	40
Bought on the Internet	15	35	*	2	10	18	*	14
Some Other Way	70	53	25	11	31	19	13	23

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.48B Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	80.2	76.0	56.2	57.4	74.6	74.6	87.7	79.7
From More Than One Doctor	9.7	9.3	9.5	13.6	8.1	11.7	10.6	7.3
Wrote Fake Prescription	2.1	1.2	1.7	*	0.0 ^a	1.7	3.3	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2.1	2.6	*	4.5	0.9	2.1	3.0	*
From Friend or Relative for Free	15.2	17.3	24.1	22.3	19.7	18.7	11.0	15.8
Bought from Friend or Relative	10.7	14.8	17.9	15.6	15.9	20.3	6.4	11.5
Took from Friend or Relative without Asking	4.8	3.3	18.2	14.4	5.3	4.7	2.2	0.7
Bought from Drug Dealer or Other Stranger	6.4	7.0	7.3	14.6	10.3	12.3	4.0	2.9
Bought on the Internet	0.6	1.5	*	1.1	1.1	2.4	*	1.0
Some Other Way	2.8	2.3	10.6	5.4	3.9	2.5	0.9	1.6

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.49A Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	501	464	28	27	127	96	346	341
From More Than One Doctor	42	47	8	0	12	12	22	35
Wrote Fake Prescription	22	9	2	4	2	4	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	24	13	3	2	5	11	*	*
From Friend or Relative for Free	2,880	2,970	171	169	881	842	1,828	1,959
Bought from Friend or Relative	494	601	68	72	302	295	125	235
Took from Friend or Relative without Asking	156 ^a	324	33	44	72	95	51	185
Bought from Drug Dealer or Other Stranger	375 ^a	240	51	41	174	145	150	53
Bought on the Internet	20	48	*	2	10	5	8	40
Some Other Way	151	164	16	25	50	51	85	89

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.49B Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	10.7	9.5	7.4	7.0	7.8	6.2	13.0	11.6
From More Than One Doctor	0.9	1.0	2.0	0.1	0.7	0.8	0.8	1.2
Wrote Fake Prescription	0.5	0.2	0.6	1.1	0.1	0.3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	0.3	0.7	0.5	0.3	0.7	*	*
From Friend or Relative for Free	61.7	60.9	44.9	43.7	53.9	54.1	69.0	66.7
Bought from Friend or Relative	10.6	12.3	17.7	18.6	18.5	18.9	4.7	8.0
Took from Friend or Relative without Asking	3.3 ^a	6.6	8.7	11.3	4.4	6.1	1.9	6.3
Bought from Drug Dealer or Other Stranger	8.0 ^a	4.9	13.5	10.6	10.6	9.3	5.7 ^a	1.8
Bought on the Internet	0.4	1.0	*	0.6	0.6	0.3	0.3	1.4
Some Other Way	3.2	3.4	4.3	6.4	3.1	3.2	3.2	3.0

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.50A Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	2,085	2,004	65	55	517	463	1,503	1,486
From More Than One Doctor	64	62	*	6	31	26	30	30
Wrote Fake Prescription	0	2	*	*	0	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	10	*	*	2	4	*	6
From Friend or Relative for Free	189	279	36	27	101	113	52	139
Bought from Friend or Relative	158	135	*	17	59	78	84	41
Took from Friend or Relative without Asking	21	35	4	*	17	7	*	*
Bought from Drug Dealer or Other Stranger	52	88	9	14	24	26	19	49
Bought on the Internet	20	5	*	*	7	*	*	1
Some Other Way	10	50	*	1	5	19	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.50B Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	79.9	75.1	47.9	40.7	67.7	62.4	87.9	82.8
From More Than One Doctor	2.5	2.3	*	4.6	4.0	3.5	1.8	1.7
Wrote Fake Prescription	0.0	0.1	*	*	0.0	0.2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.4	0.4	*	*	0.2	0.6	*	0.3
From Friend or Relative for Free	7.3	10.5	26.6	19.7	13.3	15.3	3.0	7.7
Bought from Friend or Relative	6.1	5.1	*	12.3	7.8	10.5	4.9	2.3
Took from Friend or Relative without Asking	0.8	1.3	2.8	*	2.2	1.0	*	*
Bought from Drug Dealer or Other Stranger	2.0	3.3	6.5	10.2	3.2	3.5	1.1	2.7
Bought on the Internet	0.8	0.2	*	*	1.0	*	*	0.1
Some Other Way	0.4	1.9	*	0.8	0.7	2.5	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.51A Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	211	240	*	15	69	47	133	179
From More Than One Doctor	23	26	*	*	10	5	*	*
Wrote Fake Prescription	*	4	*	*	2	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	8	*	0	1	8	*	*
From Friend or Relative for Free	898	1,011	*	*	291	285	568	661
Bought from Friend or Relative	233	274	*	*	153	135	56	*
Took from Friend or Relative without Asking	83	152	*	10	45	52	21	*
Bought from Drug Dealer or Other Stranger	204	157	*	17	96	62	*	*
Bought on the Internet	19	33	*	*	6	7	*	*
Some Other Way	58	77	*	4	19	18	32	55

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.51B Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	13.1	14.5	*	11.6	11.6	9.1	14.5	17.5
From More Than One Doctor	1.4	1.6	*	*	1.7	0.9	*	*
Wrote Fake Prescription	*	0.3	*	*	0.3	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.5	*	0.2	0.1	1.5	*	*
From Friend or Relative for Free	56.0	60.8	*	*	48.9	55.3	62.1	64.7
Bought from Friend or Relative	14.6	16.5	*	*	25.7	26.3	6.1	*
Took from Friend or Relative without Asking	5.2	9.1	*	8.2	7.6	10.0	2.3	*
Bought from Drug Dealer or Other Stranger	12.7	9.4	*	13.8	16.1	12.1	*	*
Bought on the Internet	1.2	2.0	*	*	1.0	1.3	*	*
Some Other Way	3.7	4.7	*	3.5	3.3	3.6	3.5	5.4

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.52A Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	637	659	*	*	182	169	*	*
From More Than One Doctor	32	*	*	*	10	20	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	3	*	*	4	*	*	*
From Friend or Relative for Free	97	116	*	*	43	69	*	*
Bought from Friend or Relative	67	71	*	*	28	36	*	*
Took from Friend or Relative without Asking	15	22	*	*	9	*	*	*
Bought from Drug Dealer or Other Stranger	26	61	*	*	15	23	*	*
Bought on the Internet	*	0	*	*	5	0	*	*
Some Other Way	4	9	*	*	*	4	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.52B Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	77.6	73.0	*	*	72.9	65.2	*	*
From More Than One Doctor	3.9	*	*	*	4.2	7.7	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5	0.3	*	*	1.6	*	*	*
From Friend or Relative for Free	11.9	12.8	*	*	17.4	26.6	*	*
Bought from Friend or Relative	8.1	7.9	*	*	11.4	13.8	*	*
Took from Friend or Relative without Asking	1.8	2.4	*	*	3.5	*	*	*
Bought from Drug Dealer or Other Stranger	3.2	6.7	*	*	6.0	8.7	*	*
Bought on the Internet	*	0.0	*	*	2.2	0.2	*	*
Some Other Way	0.4	1.0	*	*	*	1.7	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.53A Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	215	191	53	27	75	47	87	117
From More Than One Doctor	22	12	4	4	4	8	14	*
Wrote Fake Prescription	16	5	3	*	*	4	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	15	25	11 ^a	2	4	13	*	11
From Friend or Relative for Free	1,260	1,061	149	134	527	427	584	500
Bought from Friend or Relative	261	319	39	49	193	184	29 ^a	87
Took from Friend or Relative without Asking	160	96	37	32	51	37	72	27
Bought from Drug Dealer or Other Stranger	117	134	29	22	52	63	36	50
Bought on the Internet	89	81	5	1	10	13	74	67
Some Other Way	125 ^a	45	22	15	25	12	*	18

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.53B Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	9.4	9.7	15.1	9.5	8.0	5.8	8.8	13.3
From More Than One Doctor	1.0	0.6	1.0	1.5	0.4	1.0	1.4	*
Wrote Fake Prescription	0.7	0.3	0.7	*	*	0.5	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	1.3	3.2 ^a	0.5	0.4	1.6	*	1.2
From Friend or Relative for Free	55.2	53.9	42.5	46.7	56.0	53.0	59.1	57.1
Bought from Friend or Relative	11.5 ^a	16.2	11.1	16.9	20.5	22.8	3.0 ^a	9.9
Took from Friend or Relative without Asking	7.0	4.9	10.6	11.3	5.4	4.6	7.3	3.1
Bought from Drug Dealer or Other Stranger	5.1	6.8	8.2	7.5	5.5	7.8	3.6	5.7
Bought on the Internet	3.9	4.1	1.4	0.3	1.1	1.7	7.5	7.6
Some Other Way	5.5	2.3	6.2	5.2	2.6	1.5	*	2.1

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.54A Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	774	671	78	*	366	306	*	*
From More Than One Doctor	26	32	8	*	15	13	*	*
Wrote Fake Prescription	*	0	*	0	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	*	*	*	*
From Friend or Relative for Free	131	118	20	13	54	45	*	*
Bought from Friend or Relative	*	66	4	*	25	29	*	*
Took from Friend or Relative without Asking	10	7	*	*	4	1	*	*
Bought from Drug Dealer or Other Stranger	13	12	3	7	7	5	*	*
Bought on the Internet	16	12	*	*	7	*	*	*
Some Other Way	1	11	*	1	*	1	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.54B Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	69.5	72.0	62.4	*	76.4	75.3	*	*
From More Than One Doctor	2.3	3.4	6.7	*	3.1	3.2	*	*
Wrote Fake Prescription	*	0.0	*	0.1	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2	0.2	*	*	*	*	*	*
From Friend or Relative for Free	11.8	12.7	16.3	10.6	11.4	11.1	*	*
Bought from Friend or Relative	*	7.1	2.9	*	5.3	7.0	*	*
Took from Friend or Relative without Asking	0.9	0.7	*	*	0.9	0.3	*	*
Bought from Drug Dealer or Other Stranger	1.1	1.3	2.8	5.8	1.4	1.3	*	*
Bought on the Internet	1.4	1.3	*	*	1.5	*	*	*
Some Other Way	0.1	1.2	*	1.1	*	0.2	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.55A Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	119	95	*	*	36	27	*	*
From More Than One Doctor	17	20	*	*	3	*	*	*
Wrote Fake Prescription	3	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	9	14	*	*	*	*	*	*
From Friend or Relative for Free	357	360	*	*	175	133	*	*
Bought from Friend or Relative	115	145	*	*	89	82	*	*
Took from Friend or Relative without Asking	95	51	*	*	36	*	*	*
Bought from Drug Dealer or Other Stranger	63	47	*	*	38	32	*	*
Bought on the Internet	50	43	*	*	7	*	*	*
Some Other Way	*	15	*	*	9	3	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.55B Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	16.0	14.3	*	*	12.0	11.1	*	*
From More Than One Doctor	2.3	2.9	*	*	1.2	*	*	*
Wrote Fake Prescription	0.3	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1.3	2.1	*	*	*	*	*	*
From Friend or Relative for Free	47.8	54.0	*	*	58.8	53.8	*	*
Bought from Friend or Relative	15.3	21.6	*	*	30.0	33.1	*	*
Took from Friend or Relative without Asking	12.8	7.6	*	*	12.0	*	*	*
Bought from Drug Dealer or Other Stranger	8.4	7.0	*	*	12.9	13.0	*	*
Bought on the Internet	6.7	6.4	*	*	2.3	*	*	*
Some Other Way	*	2.2	*	*	3.2	1.1	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.56A Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	244	*	*	*	129	*	*	*
From More Than One Doctor	10	10	*	*	3	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	81	*	*	*	*	*	*	*
Bought from Friend or Relative	33	*	*	*	26	*	*	*
Took from Friend or Relative without Asking	8	4	*	*	6	*	*	*
Bought from Drug Dealer or Other Stranger	*	18	*	*	17	*	*	*
Bought on the Internet	7	*	*	*	*	*	*	*
Some Other Way	3	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.56B Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	70.9	*	*	*	75.5	*	*	*
From More Than One Doctor	3.1	3.2	*	*	1.8	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	23.7	*	*	*	*	*	*	*
Bought from Friend or Relative	9.8	*	*	*	15.1	*	*	*
Took from Friend or Relative without Asking	2.5	1.3	*	*	3.7	*	*	*
Bought from Drug Dealer or Other Stranger	*	5.4	*	*	9.8	*	*	*
Bought on the Internet	2.0	*	*	*	*	*	*	*
Some Other Way	0.8	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.57A Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	905 ^b	526	78	*	265 ^a	181	562	*
Bought from Friend or Relative	361	361	33	*	107	87	220	260
Took from Friend or Relative without Asking	28	26	10	*	9	9	*	12
Bought from Drug Dealer or Other Stranger	356	235	*	*	97	70	227	*
Bought on the Internet	*	4	*	*	1	*	*	*
Some Other Way	16	36	4	*	11	15	1	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.57B Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	53.6	44.3	49.4	*	54.1	50.0	54.0	*
Bought from Friend or Relative	21.4	30.4	21.2	*	21.8	24.0	21.2	35.4
Took from Friend or Relative without Asking	1.7	2.2	6.3	*	1.9	2.5	*	1.6
Bought from Drug Dealer or Other Stranger	21.1	19.8	*	*	19.8	19.4	21.8	*
Bought on the Internet	*	0.4	*	*	0.2	*	*	*
Some Other Way	0.9	3.0	2.5	*	2.2	4.1	0.1	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.58A Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	150	*	*	*	57	*	*	*
Bought from Friend or Relative	322 ^a	185	*	*	85	*	*	*
Took from Friend or Relative without Asking	15	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	263	*	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	21	19	*	*	11	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.58B Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	19.5	*	*	*	23.4	*	*	*
Bought from Friend or Relative	41.7	41.9	*	*	35.0	*	*	*
Took from Friend or Relative without Asking	2.0	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	34.1	*	*	*	*	*	*	*
Bought on the Internet	0.0	*	*	*	*	*	*	*
Some Other Way	2.8	4.2	*	*	4.5	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.59A Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	229	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	11	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	2	*	*	*	*	*	*	*
Some Other Way	14	19	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.59B Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	35.7	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.8	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0.3	*	*	*	*	*	*	*
Some Other Way	2.2	4.1	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.60A Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	9	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports.

See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.60B Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	2.6	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports.

See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.61A Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	229	*	*	*	*	15	*	*
From More Than One Doctor	15	9	*	*	6	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	10	*	*	*	*	*	*
From Friend or Relative for Free	428	516	*	*	*	91	*	*
Bought from Friend or Relative	*	35	*	*	*	23	*	*
Took from Friend or Relative without Asking	24	19	*	*	12	*	*	*
Bought from Drug Dealer or Other Stranger	16	35	*	*	9	9	*	*
Bought on the Internet	5	5	*	*	*	*	*	*
Some Other Way	21	28	*	*	*	4	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.61B Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	27.4	*	*	*	*	9.8	*	*
From More Than One Doctor	1.8	1.1	*	*	3.8	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.1	1.2	*	*	*	*	*	*
From Friend or Relative for Free	51.1	65.4	*	*	*	59.3	*	*
Bought from Friend or Relative	*	4.4	*	*	*	14.9	*	*
Took from Friend or Relative without Asking	2.8	2.5	*	*	7.2	*	*	*
Bought from Drug Dealer or Other Stranger	1.9	4.5	*	*	5.2	5.8	*	*
Bought on the Internet	0.6	0.7	*	*	*	*	*	*
Some Other Way	2.5	3.6	*	*	*	2.8	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.62A Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	394	*	*	*	*	*	*
From More Than One Doctor	10	11	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0	*	*	*	*	*	*	*
From Friend or Relative for Free	9	16	*	*	*	*	*	*
Bought from Friend or Relative	5	6	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3	*	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	*	*	*	*
Some Other Way	*	1	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.62B Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	91.7	*	*	*	*	*	*
From More Than One Doctor	2.5	2.7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.0	*	*	*	*	*	*	*
From Friend or Relative for Free	2.2	3.7	*	*	*	*	*	*
Bought from Friend or Relative	1.2	1.4	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	0.3	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.8	*	*	*	*	*	*	*
Bought on the Internet	0.0	0.1	*	*	*	*	*	*
Some Other Way	*	0.2	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.63A Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.63B Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.64A Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.64B Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

^a Difference between estimate and 2007 estimate is statistically significant at the 0.05 level.

^b Difference between estimate and 2007 estimate is statistically significant at the 0.01 level.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.