

**Table 7.69A Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Drug	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>ILLCIT DRUGS<sup>2</sup></b>	5,963	5,913	97	135	5,836	5,718
Marijuana and Hashish	4,272	4,179	71	98	4,179	4,031
Cocaine	652	590	7	10	640	577
Crack	167	112	2	7	165 <sup>a</sup>	104
Heroin	38	47	3	2	35	45
Hallucinogens	274 <sup>a</sup>	345	6	4	263 <sup>a</sup>	337
LSD	29	33	3	0	26	32
PCP	9	7	*	0	8	6
Ecstasy	156 <sup>a</sup>	218	3	2	150 <sup>a</sup>	216
Inhalants	125	162	2	7	121	154
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	2,367	2,256	35	32	2,319	2,204
Pain Relievers	1,674	1,608	29	20	1,634	1,571
OxyContin <sup>®</sup>	126	98	1	1	124	97
Tranquilizers	678	676	6	8	668	662
Stimulants <sup>4</sup>	551	466	5	10	542	454
Methamphetamine <sup>4</sup>	266	209	3	2	262	206
Sedatives	101	116	1	1	100	114
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	2,929	2,887	40	49	2,872	2,815

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.69B Types of Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Drug	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>ILLCIT DRUGS<sup>2</sup></b>	9.6	9.5	3.9	5.2	9.9	9.7
Marijuana and Hashish	6.9	6.7	2.8	3.8	7.1	6.8
Cocaine	1.1	1.0	0.3	0.4	1.1	1.0
Crack	0.3	0.2	0.1	0.3	0.3 <sup>a</sup>	0.2
Heroin	0.1	0.1	0.1	0.1	0.1	0.1
Hallucinogens	0.4 <sup>a</sup>	0.6	0.2	0.1	0.4 <sup>a</sup>	0.6
LSD	0.0	0.1	0.1	0.0	0.0	0.1
PCP	0.0	0.0	*	0.0	0.0	0.0
Ecstasy	0.3 <sup>a</sup>	0.4	0.1	0.1	0.3 <sup>a</sup>	0.4
Inhalants	0.2	0.3	0.1	0.3	0.2	0.3
Nonmedical Use of Psychotherapeutics <sup>3,4</sup>	3.8	3.6	1.4	1.2	3.9	3.7
Pain Relievers	2.7	2.6	1.2	0.8	2.8	2.7
OxyContin <sup>®</sup>	0.2	0.2	0.1	0.0	0.2	0.2
Tranquilizers	1.1	1.1	0.3	0.3	1.1	1.1
Stimulants <sup>4</sup>	0.9	0.8	0.2	0.4	0.9	0.8
Methamphetamine <sup>4</sup>	0.4	0.3	0.1	0.1	0.4	0.3
Sedatives	0.2	0.2	0.0	0.0	0.2	0.2
<b>ILLCIT DRUGS OTHER THAN MARIJUANA<sup>2</sup></b>	4.7	4.7	1.6	1.9	4.9	4.8

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

<sup>3</sup> Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

<sup>4</sup> Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.70A Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	5,963	5,913	97	135	5,836	5,718
<b>AGE</b>						
15-17	908	867	9 <sup>a</sup>	20	884	832
18-25	2,499	2,528	63	69	2,425	2,430
26-44	2,556	2,519	24	46	2,526	2,456
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	5,220	5,274	82	120	5,110	5,101
White	4,185	4,135	58	92	4,105	4,012
Black or African American	750	857	22	22	723	820
American Indian or Alaska Native	50	46	*	*	49	43
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	80	116	*	*	79	111
Two or More Races	129	101	*	*	127	96
Hispanic or Latino	743	639	15	15	725	616
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	50	45	N/A	N/A
Second	N/A	N/A	26	54	N/A	N/A
Third	N/A	N/A	21	27	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.70B Illicit Drug Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	9.6	9.5	3.9	5.2	9.9	9.7
<b>AGE</b>						
15-17	14.8 <sup>a</sup>	13.5	12.3	22.6	14.7 <sup>a</sup>	13.3
18-25	15.5	15.6	7.0	7.2	16.1	16.0
26-44	6.4	6.4	1.6	3.0	6.7	6.5
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	10.0	10.2	4.1	5.8	10.2	10.4
White	10.7	10.8	3.9	6.2	10.9	10.9
Black or African American	8.8	9.9	6.8	5.7	8.9	10.1
American Indian or Alaska Native	12.9	14.3	*	*	13.0	13.8
Native Hawaiian or Other Pacific Islander	*	*	*	*	*	*
Asian	2.5	3.6	*	*	2.6	3.7
Two or More Races	19.5 <sup>a</sup>	13.3	*	*	19.8 <sup>a</sup>	13.1
Hispanic or Latino	7.7 <sup>a</sup>	6.2	3.0	2.9	7.9 <sup>a</sup>	6.3
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	7.0	5.3	N/A	N/A
Second	N/A	N/A	3.2	5.7	N/A	N/A
Third	N/A	N/A	2.3	3.5	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.71A Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	4,272	4,179	71	98	4,179	4,031
<b>AGE</b>						
15-17	679	647	8	16	662	619
18-25	1,937	1,996	45	51	1,883	1,922
26-44	1,657	1,536	18	31	1,634	1,490
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	3,818	3,775	60	90	3,738	3,641
White	3,074	2,951	46	70	3,013	2,854
Black or African American	547	630	12	18	531	599
American Indian or Alaska Native	40	33	*	*	39	32
Native Hawaiian or Other Pacific Islander	9	7	*	*	9	7
Asian	57	69	*	*	57	69
Two or More Races	90	86	*	*	89	81
Hispanic or Latino	454	404	11	8	441	390
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	36	39	N/A	N/A
Second	N/A	N/A	22	40	N/A	N/A
Third	N/A	N/A	13	13	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.71B Marijuana Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	6.9	6.7	2.8	3.8	7.1	6.8
<b>AGE</b>						
15-17	11.0	10.1	10.7	17.7	11.0 <sup>a</sup>	9.9
18-25	12.0	12.3	4.9	5.3	12.5	12.7
26-44	4.2	3.9	1.2	2.0	4.3	4.0
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	7.3	7.3	3.0	4.3	7.5	7.4
White	7.8	7.7	3.1	4.7	8.0	7.8
Black or African American	6.4	7.3	3.8	4.6	6.5	7.4
American Indian or Alaska Native	10.3	10.2	*	*	10.4	10.4
Native Hawaiian or Other Pacific Islander	4.3	2.7	*	*	4.3	2.8
Asian	1.8	2.1	*	*	1.9	2.3
Two or More Races	13.7	11.3	*	*	13.9	11.1
Hispanic or Latino	4.7	3.9	2.2	1.5	4.8	4.0
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	5.0	4.7	N/A	N/A
Second	N/A	N/A	2.6	4.2	N/A	N/A
Third	N/A	N/A	1.5	1.7	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.72A Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Substance	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOBACCO PRODUCTS<sup>2</sup></b>	18,627	18,069	423	431	18,124	17,514
Cigarettes	17,948	17,317	413	426	17,455	16,774
Smokeless Tobacco	184	211	4	3	179	200
Cigars	1,926	2,025	28	34	1,889	1,974
Pipe Tobacco	206	182	3	2	203	178
<b>ALCOHOL</b>	31,824	31,874	300	302	31,376	31,386
Binge Alcohol Use <sup>3</sup>	13,939	14,427	96	97	13,782	14,240
Heavy Alcohol Use <sup>3</sup>	3,158	3,302	18	19	3,132	3,263

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.72B Tobacco Product and Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy Status: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Substance	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOBACCO PRODUCTS<sup>2</sup></b>	30.1	29.1	17.0	16.6	30.7 <sup>a</sup>	29.7
Cigarettes	29.0 <sup>a</sup>	27.9	16.6	16.4	29.6 <sup>a</sup>	28.4
Smokeless Tobacco	0.3	0.3	0.2	0.1	0.3	0.3
Cigars	3.1	3.3	1.1	1.3	3.2	3.3
Pipe Tobacco	0.3	0.3	0.1	0.1	0.3	0.3
<b>ALCOHOL</b>	51.4	51.4	12.1	11.6	53.1	53.2
Binge Alcohol Use <sup>3</sup>	22.5	23.3	3.9	3.7	23.3	24.1
Heavy Alcohol Use <sup>3</sup>	5.1	5.3	0.7	0.7	5.3	5.5

\*Low precision; no estimate reported.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

<sup>3</sup> Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.



**Table 7.73A Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	17,948	17,317	413	426	17,455	16,774
<b>AGE</b>						
15-17	1,149 <sup>a</sup>	1,039	17	21	1,117 <sup>a</sup>	1,002
18-25	5,675 <sup>b</sup>	5,417	240	224	5,400 <sup>a</sup>	5,155
26-44	11,123	10,860	157	180	10,939	10,616
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	16,050	15,538	377	386	15,606	15,056
White	13,241	12,808	319	298	12,868	12,464
Black or African American	2,030	1,893	48	58	1,972	1,804
American Indian or Alaska Native	156	127	*	*	152	123
Native Hawaiian or Other Pacific Islander	*	53	*	*	*	49
Asian	289	390	*	*	286	359
Two or More Races	262	267	*	*	257	257
Hispanic or Latino	1,898	1,779	36	40	1,849	1,718
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	170	179	N/A	N/A
Second	N/A	N/A	105	147	N/A	N/A
Third	N/A	N/A	126 <sup>a</sup>	85	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.73B Cigarette Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	29.0 <sup>a</sup>	27.9	16.6	16.4	29.6 <sup>a</sup>	28.4
<b>AGE</b>						
15-17	18.7 <sup>b</sup>	16.2	22.3	24.3	18.5 <sup>b</sup>	16.0
18-25	35.3 <sup>b</sup>	33.3	26.4	23.3	35.8 <sup>b</sup>	33.9
26-44	28.1	27.6	10.4	11.6	28.8	28.3
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	30.8	30.1	19.0	18.5	31.3	30.6
White	33.8	33.3	21.5	19.9	34.3	33.9
Black or African American	23.8	22.0	15.0	15.0	24.2	22.2
American Indian or Alaska Native	40.6	39.5	*	*	40.6	39.8
Native Hawaiian or Other Pacific Islander	*	21.1	*	*	*	20.4
Asian	9.1	12.0	*	*	9.5	11.8
Two or More Races	39.6	35.1	*	*	40.0	35.1
Hispanic or Latino	19.5 <sup>a</sup>	17.2	7.2	7.7	20.2 <sup>a</sup>	17.6
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	23.7	21.4	N/A	N/A
Second	N/A	N/A	12.9	15.4	N/A	N/A
Third	N/A	N/A	13.7	10.8	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.74A Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	31,824	31,874	300	302	31,376	31,386
<b>AGE</b>						
15-17	1,699	1,621	10	14	1,672	1,591
18-25	8,963 <sup>b</sup>	9,345	88	94	8,826	9,193
26-44	21,163	20,908	202	194	20,879	20,601
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	28,192	27,905	266	281	27,801	27,466
White	23,095	22,595	204	217	22,790	22,280
Black or African American	3,453	3,662	43	61	3,388	3,565
American Indian or Alaska Native	170	134	*	*	169	131
Native Hawaiian or Other Pacific Islander	*	86	*	*	*	86
Asian	994	1,010	*	*	983	991
Two or More Races	400	419	*	*	397	413
Hispanic or Latino	3,632	3,969	34	21	3,575 <sup>a</sup>	3,920
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	147	147	N/A	N/A
Second	N/A	N/A	84	79	N/A	N/A
Third	N/A	N/A	61	63	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.74B Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	51.4	51.4	12.1	11.6	53.1	53.2
<b>AGE</b>						
15-17	27.6 <sup>b</sup>	25.3	13.9	15.8	27.7 <sup>b</sup>	25.4
18-25	55.7 <sup>b</sup>	57.5	9.7	9.8	58.5 <sup>b</sup>	60.5
26-44	53.4	53.1	13.5	12.5	55.0	54.9
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	54.1	54.0	13.4	13.5	55.7	55.8
White	58.9	58.8	13.8	14.5	60.7	60.6
Black or African American	40.5	42.5	13.4	15.7	41.6	43.8
American Indian or Alaska Native	44.3	41.4	*	*	45.0	42.3
Native Hawaiian or Other Pacific Islander	*	34.5	*	*	*	35.9
Asian	31.2	31.1	*	*	32.4	32.6
Two or More Races	60.5	55.1	*	*	61.8	56.4
Hispanic or Latino	37.4	38.3	6.8	4.1	39.1	40.1
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	20.6	17.6	N/A	N/A
Second	N/A	N/A	10.2	8.2	N/A	N/A
Third	N/A	N/A	6.7	8.0	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.75A Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	13,939	14,427	96	97	13,782	14,240
<b>AGE</b>						
15-17	1,023	968	6	9	1,003	947
18-25	5,162 <sup>b</sup>	5,501	45	49	5,089 <sup>a</sup>	5,424
26-44	7,754	7,959	45	39	7,690	7,868
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	12,178	12,540	78	83	12,051	12,380
White	10,002	10,421	47	51	9,915	10,323
Black or African American	1,485	1,458	28	30	1,447	1,413
American Indian or Alaska Native	121	102	*	*	120	100
Native Hawaiian or Other Pacific Islander	*	55	*	*	*	55
Asian	345	321	*	*	344	309
Two or More Races	179	183	*	*	177	179
Hispanic or Latino	1,761	1,887	19	13	1,732	1,860
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	54	56	N/A	N/A
Second	N/A	N/A	21	21	N/A	N/A
Third	N/A	N/A	15	7	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.75B Binge Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	22.5	23.3	3.9	3.7	23.3	24.1
<b>AGE</b>						
15-17	16.6 <sup>a</sup>	15.1	8.4	10.3	16.6 <sup>a</sup>	15.1
18-25	32.1 <sup>b</sup>	33.9	4.9	5.1	33.8 <sup>b</sup>	35.7
26-44	19.6	20.2	3.0	2.5	20.3	21.0
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	23.3 <sup>a</sup>	24.3	3.9	4.0	24.1 <sup>a</sup>	25.1
White	25.5 <sup>b</sup>	27.1	3.2	3.4	26.4 <sup>b</sup>	28.1
Black or African American	17.4	16.9	8.8	7.8	17.8	17.4
American Indian or Alaska Native	31.6	31.8	*	*	32.1	32.3
Native Hawaiian or Other Pacific Islander	*	22.3	*	*	*	23.2
Asian	10.8	9.9	*	*	11.3	10.2
Two or More Races	27.1	24.1	*	*	27.6	24.4
Hispanic or Latino	18.1	18.2	3.7	2.5	18.9	19.0
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	7.5	6.6	N/A	N/A
Second	N/A	N/A	2.6	2.2	N/A	N/A
Third	N/A	N/A	1.6	0.9	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.76A Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Numbers in Thousands, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	3,158	3,302	18	19	3,132	3,263
<b>AGE</b>						
15-17	216	212	2	2	211	204
18-25	1,418 <sup>b</sup>	1,587	7	9	1,408 <sup>a</sup>	1,572
26-44	1,524	1,503	9	7	1,513	1,487
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	2,904	2,968	13	16	2,883	2,933
White	2,552	2,590	12	13	2,533	2,568
Black or African American	235	244	1	3	234	232
American Indian or Alaska Native	21	30	*	*	21	30
Native Hawaiian or Other Pacific Islander	10	7	*	*	10	7
Asian	40	50	*	*	40	50
Two or More Races	45	47	*	*	44	46
Hispanic or Latino	254	334	*	3	249 <sup>a</sup>	330
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	12	14	N/A	N/A
Second	N/A	N/A	6	2	N/A	N/A
Third	N/A	N/A	1	2	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.

**Table 7.76B Heavy Alcohol Use in the Past Month among Females Aged 15 to 44, by Pregnancy and Demographic Characteristics: Percentages, Annual Averages Based on 2004-2005 and 2006-2007**

Demographic/Pregnancy Characteristic	Total <sup>1</sup>		PREGNANCY STATUS			
			Pregnant		Not Pregnant	
	2004-2005	2006-2007	2004-2005	2006-2007	2004-2005	2006-2007
<b>TOTAL</b>	5.1	5.3	0.7	0.7	5.3	5.5
<b>AGE</b>						
15-17	3.5	3.3	2.9	2.8	3.5	3.3
18-25	8.8 <sup>b</sup>	9.8	0.8	0.9	9.3 <sup>b</sup>	10.4
26-44	3.8	3.8	0.6	0.5	4.0	4.0
<b>HISPANIC ORIGIN AND RACE</b>						
Not Hispanic or Latino	5.6	5.7	0.7	0.8	5.8	6.0
White	6.5	6.7	0.8	0.8	6.8	7.0
Black or African American	2.8	2.8	0.3	0.7	2.9	2.8
American Indian or Alaska Native	5.6	9.5	*	*	5.7	9.8
Native Hawaiian or Other Pacific Islander	4.7	2.9	*	*	4.7	3.0
Asian	1.3	1.6	*	*	1.3	1.7
Two or More Races	6.8	6.1	*	*	6.9	6.3
Hispanic or Latino	2.6	3.2	*	0.6	2.7	3.4
<b>TRIMESTER<sup>2</sup></b>						
First	N/A	N/A	1.7	1.7	N/A	N/A
Second	N/A	N/A	0.7	0.2	N/A	N/A
Third	N/A	N/A	0.1	0.3	N/A	N/A

\*Low precision; no estimate reported.

N/A: Not applicable.

NOTE: Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on each of 5 or more days in the past 30 days.

<sup>a</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.05 level.

<sup>b</sup> Difference between estimate and 2006-2007 estimate is statistically significant at the 0.01 level.

<sup>1</sup> Estimates in the Total column are for all females aged 15 to 44, including those with unknown pregnancy status.

<sup>2</sup> Pregnant females aged 15 to 44 not reporting trimester were excluded.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2004, 2005, 2006, and 2007.