

Table 7.108C (REVISED) No Illicit Drug or Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drugs or Alcohol ¹		Illicit Drugs ¹		Alcohol	
	2006	2007	2006	2007	2006	2007
TOTAL	93	97	67	68	82	87
RACE²						
American Indian or Alaska Native	16	16	19	22	17	17
Asian or Pacific Islander	70	64	74	65	72	65
Asian Only	67	62	70	64	69	62
Native Hawaiian or Other Pacific Islander Only	19	*	19	*	19	*
Black or African American	34	34	26	25	28	29
White	80	84	58	58	74	77
Two or More Races	30	29	30	30	30	30
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	42	41	30	29	38	37
Not Hispanic or Latino	82	88	59	60	73	78
Black or African American	113	119	117	126	119	128
White	192	195	210	219	196	201
GENDER						
Male	64	70	46	54	57	61
Female	63	65	45	44	59	61
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	109	111	119	120	113	117
100-199%	103	105	109	112	107	111
200% or More	205	221	222	246	209	226
REGION						
Northeast	40	37	28	23	38	35
Midwest	35	35	26	27	32	32
South	57	58	38	42	51	51
West	52	54	41	39	43	49

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.108D (REVISED) No Illicit Drug or Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drugs or Alcohol ¹		Illicit Drugs ¹		Alcohol	
	2006	2007	2006	2007	2006	2007
TOTAL	0.37	0.38	0.27	0.27	0.32	0.34
RACE²						
American Indian or Alaska Native	2.63	3.74	2.12	2.60	2.13	3.82
Asian or Pacific Islander	2.09	1.56	1.39	1.22	1.64	1.56
Asian Only	2.34	1.67	1.51	1.29	1.84	1.67
Native Hawaiian or Other Pacific Islander Only	3.64	*	3.20	*	3.04	*
Black or African American	0.84	0.86	0.65	0.62	0.69	0.72
White	0.41	0.44	0.30	0.30	0.38	0.40
Two or More Races	2.32	1.50	1.74	1.19	2.12	1.30
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.91	0.88	0.66	0.62	0.84	0.80
Not Hispanic or Latino	0.39	0.43	0.28	0.29	0.35	0.38
Black or African American	0.85	0.88	0.66	0.64	0.71	0.74
White	0.44	0.49	0.31	0.33	0.42	0.44
GENDER						
Male	0.49	0.55	0.36	0.42	0.44	0.47
Female	0.50	0.53	0.36	0.35	0.47	0.49
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	0.92	0.89	0.67	0.67	0.83	0.76
100-199%	0.78	0.75	0.60	0.61	0.69	0.66
200% or More	0.45	0.49	0.32	0.32	0.41	0.45
REGION						
Northeast	0.88	0.83	0.62	0.51	0.84	0.78
Midwest	0.61	0.62	0.46	0.48	0.57	0.57
South	0.62	0.64	0.41	0.46	0.56	0.56
West	0.86	0.89	0.67	0.65	0.72	0.80

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.108P (REVISED) No Illicit Drug or Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic/Geographic Characteristic	NO PAST MONTH USE					
	Illicit Drugs or Alcohol ¹		Illicit Drugs ¹		Alcohol	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3643		0.5424		0.1687	
RACE²						
American Indian or Alaska Native	0.8711		0.8957		0.6368	
Asian or Pacific Islander	0.4131		0.5392		0.9595	
Asian Only	0.5387		0.7381		0.8413	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.6441		0.4396		0.6808	
White	0.6166		0.9701		0.1926	
Two or More Races	0.1473		0.1684		0.1130	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.9840		0.4086		0.9604	
Not Hispanic or Latino	0.3164		0.7793		0.1390	
Black or African American	0.7113		0.4096		0.6935	
White	0.4370		0.7223		0.1015	
GENDER						
Male	0.8383		0.7050		0.5950	
Female	0.2581		0.1665		0.1551	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	0.2806		0.4805		0.1412	
100-199%	0.5246		0.8376		0.3381	
200% or More	0.8564		0.7842		0.7087	
REGION						
Northeast	0.0420		0.1035		0.1321	
Midwest	0.4895		0.6594		0.5239	
South	0.8512		0.5793		0.4101	
West	0.7034		0.7831		0.3912	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-10A (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.109C (REVISED) Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	67	68	55	56	69	69
RACE³						
American Indian or Alaska Native	7	9	4	7	6	14
Asian or Pacific Islander	18	15	11	15	20	17
Asian Only	17	14	10	15	19	17
Native Hawaiian or Other Pacific Islander Only	5	*	4	2	4	1
Black or African American	26	25	20	20	20	20
White	58	58	48	48	61	62
Two or More Races	8	6	7	6	8	4
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	30	29	24	23	32	29
Not Hispanic or Latino	59	60	47	51	62	62
Black or African American	28	28	21	21	20	21
White	52	53	43	44	60	60
GENDER						
Male	46	54	39	44	47	51
Female	45	44	37	35	49	45
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	34	33	26	27	37	31
100-199%	34	35	27	27	33	29
200% or More	54	53	44	44	57	59
REGION						
Northeast	28	23	24	21	31	28
Midwest	26	27	22	23	26	27
South	38	42	31	32	43	42
West	41	39	32	34	38	38

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.109D (REVISED) Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.27	0.21	0.22	0.27	0.27
RACE³						
American Indian or Alaska Native	2.12	2.60	1.25	2.11	1.68	3.80
Asian or Pacific Islander	1.39	1.22	0.86	1.27	1.57	1.40
Asian Only	1.51	1.29	0.91	1.39	1.77	1.53
Native Hawaiian or Other Pacific Islander Only	3.20	*	2.59	1.79	2.28	0.86
Black or African American	0.65	0.62	0.50	0.50	0.49	0.50
White	0.30	0.30	0.25	0.25	0.32	0.32
Two or More Races	1.74	1.19	1.53	1.06	1.71	0.85
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.66	0.62	0.53	0.50	0.70	0.62
Not Hispanic or Latino	0.28	0.29	0.23	0.25	0.30	0.30
Black or African American	0.66	0.64	0.51	0.52	0.50	0.52
White	0.31	0.33	0.26	0.28	0.35	0.36
GENDER						
Male	0.36	0.42	0.30	0.34	0.36	0.40
Female	0.36	0.35	0.29	0.29	0.39	0.37
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	0.67	0.67	0.54	0.56	0.74	0.63
100-199%	0.60	0.61	0.49	0.48	0.59	0.52
200% or More	0.32	0.32	0.27	0.27	0.34	0.34
REGION						
Northeast	0.62	0.51	0.52	0.47	0.68	0.63
Midwest	0.46	0.48	0.39	0.41	0.46	0.47
South	0.41	0.46	0.33	0.35	0.47	0.47
West	0.67	0.65	0.53	0.57	0.62	0.63

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.109P (REVISED) Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic/Geographic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	0.5424		0.9184		0.1229	
RACE³						
American Indian or Alaska Native	0.8957		0.9771		0.2083	
Asian or Pacific Islander	0.5392		0.7982		0.5924	
Asian Only	0.7381		0.5027		0.8089	
Native Hawaiian or Other Pacific Islander Only	*		0.1931		0.0206	
Black or African American	0.4396		0.3291		0.3601	
White	0.9701		0.5699		0.2341	
Two or More Races	0.1684		0.4913		0.0145	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.4086		0.9109		0.9953	
Not Hispanic or Latino	0.7793		0.8506		0.0903	
Black or African American	0.4096		0.3206		0.3431	
White	0.7223		0.5342		0.1440	
GENDER						
Male	0.7050		0.1524		0.8423	
Female	0.1665		0.1251		0.0404	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	0.4805		0.9257		0.5778	
100-199%	0.8376		0.9216		0.1589	
200% or More	0.7842		0.7989		0.4048	
REGION						
Northeast	0.1035		0.4118		0.0370	
Midwest	0.6594		0.8781		0.3878	
South	0.5793		0.2092		0.9791	
West	0.7831		0.5543		0.0824	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10B and 26-11D (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.110C Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	415	441	350	353	741	763
RACE³						
American Indian or Alaska Native	28	35	23	19	75	61
Asian or Pacific Islander	66	87	44	49	138	141
Asian Only	62	81	40	48	135	131
Native Hawaiian or Other Pacific Islander Only	15	28	14	6	40	50
Black or African American	155	153	131	129	249	234
White	366	391	308	318	679	711
Two or More Races	31	43	30	42	82	68
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	140	132	103	101	268	276
Not Hispanic or Latino	391	412	334	334	682	703
Black or African American	160	168	136	140	274	277
White	354	380	304	307	721	762
GENDER						
Male	326	338	279	283	561	585
Female	232	237	175	172	395	391
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	167	151	119	110	254	264
100-199%	176	186	148	154	338	323
200% or More	342	359	287	284	708	753
EDUCATION						
< High School	173	180	131	148	346	318
High School Graduate	219	241	187	197	450	466
Some College/College Graduate	312	308	271	247	604	642

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.110D Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	0.19	0.20	0.16	0.16	0.34	0.34
RACE³						
American Indian or Alaska Native	1.45	1.64	1.16	0.96	3.19	2.66
Asian or Pacific Islander	0.63	0.82	0.43	0.46	1.26	1.28
Asian Only	0.66	0.84	0.43	0.50	1.36	1.28
Native Hawaiian or Other Pacific Islander Only	1.65	3.14	1.53	0.75	4.40	5.04
Black or African American	0.60	0.59	0.51	0.50	0.97	0.90
White	0.20	0.22	0.17	0.18	0.38	0.39
Two or More Races	1.16	1.54	1.12	1.50	2.80	2.37
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.48	0.45	0.36	0.34	0.93	0.93
Not Hispanic or Latino	0.20	0.21	0.17	0.17	0.36	0.36
Black or African American	0.59	0.60	0.50	0.51	0.98	0.92
White	0.22	0.24	0.19	0.19	0.40	0.42
GENDER						
Male	0.31	0.32	0.26	0.26	0.53	0.54
Female	0.20	0.21	0.15	0.15	0.35	0.34
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	0.60	0.56	0.45	0.42	0.88	0.92
100-199%	0.40	0.45	0.34	0.38	0.70	0.72
200% or More	0.22	0.22	0.19	0.18	0.41	0.41
EDUCATION						
< High School	0.47	0.48	0.36	0.40	0.83	0.81
High School Graduate	0.31	0.35	0.27	0.28	0.59	0.60
Some College/College Graduate	0.26	0.25	0.23	0.21	0.46	0.47

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.110P Illicit Drug Use, Marijuana Use, and Binge Alcohol Use in the Past Month among Persons Aged 18 or Older, by Demographic and Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic Characteristic	PAST MONTH USE					
	Illicit Drug Use ¹		Marijuana Use		Binge Alcohol Use ²	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3237		0.3291		0.4100	
RACE³						
American Indian or Alaska Native	0.5349		0.1279		0.6894	
Asian or Pacific Islander	0.4714		0.6247		0.6032	
Asian Only	0.5229		0.3214		0.6438	
Native Hawaiian or Other Pacific Islander Only	0.5560		0.0921		0.5451	
Black or African American	0.5380		0.7950		0.9207	
White	0.2634		0.2549		0.3914	
Two or More Races	0.1011		0.0967		0.7114	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.6646		0.3361		0.6687	
Not Hispanic or Latino	0.3683		0.1868		0.2888	
Black or African American	0.8023		0.9773		0.9425	
White	0.2268		0.0895		0.2756	
GENDER						
Male	0.7355		0.6204		0.5269	
Female	0.2102		0.2648		0.5154	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)⁴						
Less Than 100%	0.4925		0.3364		0.6142	
100-199%	0.5089		0.8045		0.4707	
200% or More	0.6745		0.5464		0.2974	
EDUCATION						
< High School	0.8917		0.3955		0.9709	
High School Graduate	0.9548		0.7492		0.8809	
Some College/College Graduate	0.1547		0.1364		0.2827	

*Low precision; no estimate reported.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-10C and 26-11C (<http://www.healthypeople.gov/>).

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days.

³ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

⁴ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.111C (REVISED) Inhalant Use in the Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2006	2007
TOTAL	45	43
RACE¹		
American Indian or Alaska Native	4	4
Asian or Pacific Islander	10	7
Asian Only	9	7
Native Hawaiian or Other Pacific Islander Only	2	1
Black or African American	14	13
White	42	39
Two or More Races	6	4
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	23	21
Not Hispanic or Latino	41	38
Black or African American	15	14
White	39	36
GENDER		
Male	32	31
Female	32	30
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²		
Less Than 100%	19	21
100-199%	24	19
200% or More	33	34
REGION		
Northeast	17	15
Midwest	17	19
South	27	27
West	26	23

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.111D (REVISED) Inhalant Use in the Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2006	2007
TOTAL	0.18	0.17
RACE¹		
American Indian or Alaska Native	1.14	1.31
Asian or Pacific Islander	0.79	0.65
Asian Only	0.89	0.71
Native Hawaiian or Other Pacific Islander Only	1.11	0.80
Black or African American	0.35	0.33
White	0.22	0.20
Two or More Races	1.25	0.80
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	0.50	0.44
Not Hispanic or Latino	0.19	0.19
Black or African American	0.36	0.34
White	0.24	0.23
GENDER		
Male	0.25	0.24
Female	0.26	0.24
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²		
Less Than 100%	0.41	0.43
100-199%	0.45	0.35
200% or More	0.20	0.21
REGION		
Northeast	0.37	0.33
Midwest	0.31	0.33
South	0.30	0.29
West	0.43	0.39

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.111P (REVISED) Inhalant Use in the Past Year among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic/Geographic Characteristic	INHALANT USE IN PAST YEAR	
	2006	2007
TOTAL	0.0829	
RACE¹		
American Indian or Alaska Native	0.5658	
Asian or Pacific Islander	0.3841	
Asian Only	0.4598	
Native Hawaiian or Other Pacific Islander Only	0.2959	
Black or African American	0.4300	
White	0.2092	
Two or More Races	0.3055	
HISPANIC ORIGIN AND RACE		
Hispanic or Latino	0.3715	
Not Hispanic or Latino	0.1576	
Black or African American	0.3748	
White	0.2997	
GENDER		
Male	0.7299	
Female	0.0347	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²		
Less Than 100%	0.8337	
100-199%	0.0004	
200% or More	0.9032	
REGION		
Northeast	0.2119	
Midwest	1.0000	
South	0.1064	
West	0.6232	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Number 26-15 (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.112C (REVISED) Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2006	2007	2006	2007	2006	2007
TOTAL	22	21	46	48	34	37
RACE¹						
American Indian or Alaska Native	1	*	5	3	3	3
Asian or Pacific Islander	*	1	8	8	3	6
Asian Only	*	1	8	8	2	5
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	9	7	20	22	14	13
White	19	19	39	42	32	33
Two or More Races	3	2	7	6	4	3
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	13	10	20	21	19	16
Not Hispanic or Latino	17	18	41	43	29	33
Black or African American	9	7	20	21	13	13
White	14	16	34	36	26	29
GENDER						
Male	15	15	33	33	25	26
Female	16	14	31	32	24	24
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²						
Less Than 100%	13	10	21	21	18	17
100-199%	14	12	25	25	21	18
200% or More	11	13	34	34	23	25
REGION						
Northeast	8	5	21	16	11	10
Midwest	8	8	20	18	15	15
South	15	14	29	32	24	26
West	12	12	22	24	18	16

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4A (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.112D (REVISED) Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.27	0.25	0.52	0.53	0.41	0.43
RACE¹						
American Indian or Alaska Native	1.36	*	4.20	2.93	2.52	3.14
Asian or Pacific Islander	*	0.38	2.26	2.13	0.97	1.63
Asian Only	*	0.19	2.55	2.36	0.54	1.62
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	0.73	0.50	1.45	1.44	1.06	0.91
White	0.31	0.30	0.58	0.62	0.48	0.50
Two or More Races	1.94	0.98	3.98	2.71	2.26	1.87
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.86	0.64	1.22	1.28	1.18	1.00
Not Hispanic or Latino	0.26	0.27	0.56	0.57	0.42	0.47
Black or African American	0.75	0.51	1.49	1.47	1.09	0.93
White	0.29	0.33	0.65	0.67	0.50	0.57
GENDER						
Male	0.36	0.36	0.74	0.72	0.58	0.61
Female	0.39	0.34	0.73	0.74	0.59	0.56
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²						
Less Than 100%	0.80	0.63	1.25	1.19	1.09	1.02
100-199%	0.77	0.65	1.23	1.28	1.08	0.98
200% or More	0.24	0.27	0.66	0.65	0.47	0.51
REGION						
Northeast	0.55	0.33	1.38	1.09	0.73	0.72
Midwest	0.41	0.42	0.94	0.92	0.75	0.82
South	0.50	0.45	0.90	0.95	0.77	0.81
West	0.62	0.60	1.09	1.09	0.94	0.79

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4A (<http://www.healthypeople.gov/>).¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.112P (REVISED) Marijuana Use, Alcohol Use, and Cigarette Use in Lifetime among Persons Aged 12 to 13, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic/Geographic Characteristic	LIFETIME USE					
	Marijuana Use		Alcohol Use		Cigarette Use	
	2006	2007	2006	2007	2006	2007
TOTAL	0.3181		0.2359		0.0726	
RACE¹						
American Indian or Alaska Native	*		0.3150		0.3171	
Asian or Pacific Islander	*		0.9007		0.2719	
Asian Only	*		0.8542		0.2026	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.4133		0.7199		0.4641	
White	0.3662		0.1355		0.0595	
Two or More Races	0.5504		0.8089		0.8765	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.1151		0.7131		0.1940	
Not Hispanic or Latino	0.8611		0.2479		0.1829	
Black or African American	0.3964		0.6194		0.4626	
White	0.8234		0.0976		0.1814	
GENDER						
Male	0.5066		0.4476		0.7443	
Female	0.4482		0.3458		0.0263	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²						
Less Than 100%	0.2320		0.9641		0.6539	
100-199%	0.1573		0.4151		0.0480	
200% or More	0.3857		0.3068		0.5047	
REGION						
Northeast	0.0871		0.0505		0.1912	
Midwest	0.8077		0.2202		0.3803	
South	0.5387		0.8530		0.7536	
West	0.9654		0.6166		0.1456	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9 and 27-4A (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.113C (REVISED) Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	109	107	107	110	108	112
RACE²						
American Indian or Alaska Native	11	12	12	19	11	12
Asian or Pacific Islander	36	38	44	42	45	40
Asian Only	33	36	42	38	43	38
Native Hawaiian or Other Pacific Islander Only	15	*	*	*	15	*
Black or African American	43	45	45	44	45	44
White	92	91	93	96	91	95
Two or More Races	12	14	18	20	17	20
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	56	47	53	52	51	53
Not Hispanic or Latino	94	95	94	98	96	98
Black or African American	60	63	77	83	77	78
White	109	110	131	142	115	117
GENDER						
Male	73	73	75	74	76	76
Female	76	76	78	80	72	83
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	64	68	80	77	71	72
100-199%	63	64	74	79	72	77
200% or More	114	122	137	152	123	129
REGION						
Northeast	38	41	47	43	44	43
Midwest	43	41	42	45	43	40
South	64	65	66	66	64	71
West	63	54	62	60	59	61

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.113D (REVISED) Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	0.44	0.43	0.43	0.44	0.43	0.45
RACE²						
American Indian or Alaska Native	2.46	3.31	2.65	3.35	2.38	3.34
Asian or Pacific Islander	2.37	2.54	2.30	2.84	2.42	2.42
Asian Only	2.49	2.71	2.55	2.92	2.71	2.60
Native Hawaiian or Other Pacific Islander Only	5.92	*	*	*	5.17	*
Black or African American	1.09	1.15	1.14	1.13	1.14	1.12
White	0.48	0.48	0.49	0.51	0.47	0.50
Two or More Races	2.62	2.29	2.59	2.32	2.52	2.57
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	1.24	1.02	1.19	1.15	1.13	1.16
Not Hispanic or Latino	0.46	0.47	0.46	0.49	0.46	0.48
Black or African American	1.11	1.18	1.16	1.16	1.16	1.14
White	0.50	0.52	0.52	0.56	0.50	0.53
GENDER						
Male	0.57	0.58	0.59	0.59	0.59	0.60
Female	0.62	0.62	0.63	0.66	0.59	0.68
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	1.04	1.09	1.08	1.13	1.01	1.05
100-199%	0.91	0.94	0.97	0.96	0.99	0.98
200% or More	0.52	0.55	0.54	0.55	0.53	0.56
REGION						
Northeast	0.86	0.94	1.07	0.98	1.00	0.97
Midwest	0.77	0.74	0.75	0.82	0.77	0.73
South	0.72	0.72	0.73	0.74	0.71	0.79
West	1.05	0.91	1.04	1.02	0.98	1.02

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.113P (REVISED) Perceptions of Great Risk of Smoking Marijuana Once a Month, Using Cocaine Once a Month, or Having Five or More Drinks of an Alcoholic Beverage Once or Twice a Week among Persons Aged 12 to 17, by Demographic, Socioeconomic, and Geographic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic/Geographic Characteristic	PERCEPTIONS OF GREAT RISK					
	Smoke Marijuana Once a Month ¹		Use Cocaine Once a Month ¹		Have Five or More Drinks Once or Twice a Week ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	0.6717		0.3344		0.9797	
RACE²						
American Indian or Alaska Native	0.6223		0.0159		0.5465	
Asian or Pacific Islander	0.1427		0.7477		0.9637	
Asian Only	0.1142		0.8212		0.8291	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.9351		0.4685		0.9451	
White	0.2337		0.6752		0.8738	
Two or More Races	0.1672		0.1621		0.3552	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.4485		0.9162		0.9276	
Not Hispanic or Latino	0.9452		0.2421		0.9500	
Black or African American	0.7817		0.4437		0.7788	
White	0.3893		0.5344		0.8713	
GENDER						
Male	0.7630		0.4416		0.9575	
Female	0.3802		0.5434		0.9343	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	0.5982		0.6439		0.2506	
100-199%	0.0780		0.1769		0.6684	
200% or More	0.8933		0.4666		0.7043	
REGION						
Northeast	0.7416		0.2459		0.3712	
Midwest	0.5910		0.7876		0.2417	
South	0.9265		0.2195		0.7169	
West	0.5753		0.2964		0.8074	

*Low precision; no estimate reported.

NOTE: This table was revised in May 2012 due to updates (for details, see the introduction to these 2007 detailed tables).

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-17B, 26-17C, and 26-17A (<http://www.healthypeople.gov/>).

¹ Respondents with unknown data were excluded.

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.114C Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Standard Errors of Numbers in Thousands, 2006 and 2007

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALTY FACILITY					
	Illicit Drugs		Alcohol		Either Illicit Drugs or Alcohol ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	126	112	124	130	163	164
RACE²						
American Indian or Alaska Native	7	4	9	*	11	*
Asian or Pacific Islander	10	*	10	1	14	*
Asian Only	10	*	*	0	13	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	67	52	50	66	72	96
White	105	100	113	108	144	132
Two or More Races	9	13	9	*	10	*
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	72	22	76	31	97	34
Not Hispanic or Latino	105	111	99	127	134	162
Black or African American	51	52	39	65	57	96
White	90	98	89	105	119	128
GENDER						
Male	110	98	99	112	137	145
Female	66	51	73	64	90	76
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	72	59	66	75	84	86
100-199%	54	45	63	49	81	78
200% or More	93	86	83	90	116	114
AGE GROUP						
12-17	18	13	14	12	20	15
18 or Older	125	110	124	130	162	164

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.114D Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: Standard Errors of Percentages, 2006 and 2007

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALTY FACILITY					
	Illicit Drugs		Alcohol		Either Illicit Drugs or Alcohol ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	0.52	0.46	0.52	0.54	0.65	0.66
RACE²						
American Indian or Alaska Native	1.73	1.51	2.13	*	2.63	*
Asian or Pacific Islander	1.68	*	1.59	0.11	2.28	*
Asian Only	2.05	*	*	0.09	2.68	*
Native Hawaiian or Other Pacific Islander Only	*	*	*	*	*	*
Black or African American	2.08	1.78	1.61	2.18	2.19	2.97
White	0.53	0.50	0.57	0.55	0.71	0.65
Two or More Races	2.41	3.22	2.45	*	2.72	*
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	1.90	0.74	2.00	1.03	2.45	1.15
Not Hispanic or Latino	0.51	0.52	0.49	0.60	0.64	0.74
Black or African American	1.72	1.81	1.37	2.21	1.90	3.02
White	0.55	0.57	0.54	0.61	0.71	0.74
GENDER						
Male	0.69	0.60	0.63	0.68	0.84	0.86
Female	0.76	0.64	0.84	0.82	1.01	0.93
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	1.64	1.50	1.52	1.92	1.83	2.11
100-199%	1.11	1.05	1.29	1.14	1.59	1.74
200% or More	0.61	0.54	0.55	0.58	0.76	0.71
AGE GROUP						
12-17	0.85	0.66	0.67	0.57	0.93	0.76
18 or Older	0.56	0.50	0.57	0.59	0.71	0.72

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.114P Received Illicit Drug or Alcohol Treatment at a Specialty Facility in the Past Year among Persons Aged 12 or Older Who Needed Illicit Drug or Alcohol Treatment in the Past Year, by Demographic and Socioeconomic Characteristics: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Demographic/Socioeconomic Characteristic	SUBSTANCE FOR WHICH TREATMENT WAS RECEIVED AT A SPECIALTY FACILITY					
	Illicit Drugs		Alcohol		Either Illicit Drugs or Alcohol ¹	
	2006	2007	2006	2007	2006	2007
TOTAL	0.1846		0.8368		0.7060	
RACE²						
American Indian or Alaska Native	0.9156		*		*	
Asian or Pacific Islander	*		0.0373		*	
Asian Only	*		*		*	
Native Hawaiian or Other Pacific Islander Only	*		*		*	
Black or African American	0.2231		0.4498		0.5037	
White	0.3672		0.8072		0.3328	
Two or More Races	0.3955		*		*	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.0074		0.0431		0.0033	
Not Hispanic or Latino	0.8148		0.2255		0.3771	
Black or African American	0.5019		0.2756		0.2764	
White	0.9853		0.4331		0.7747	
GENDER						
Male	0.5359		0.5025		0.8326	
Female	0.1204		0.4984		0.2535	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)³						
Less Than 100%	0.3040		0.6716		0.9494	
100-199%	0.4270		0.7293		0.8009	
200% or More	0.7993		0.6287		0.9481	
AGE GROUP						
12-17	0.3852		0.4312		0.3695	
18 or Older	0.2233		0.7812		0.7727	

*Low precision; no estimate reported.

NOTE: Respondents were classified as needing treatment for an illicit drug or alcohol problem if they met at least one of three criteria during the past year: (1) dependent on illicit drugs or alcohol; (2) abuse of illicit drugs or alcohol; or (3) received treatment for an illicit drug or alcohol problem at a specialty facility (i.e., drug and alcohol rehabilitation facilities [inpatient or outpatient], hospitals [inpatient only], and mental health centers). Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-18 and 26-21 (<http://www.healthypeople.gov/>).

¹ Estimates include persons who received treatment specifically for illicit drugs or alcohol, as well as persons who received treatment but did not specify for what substance(s).

² These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

³ Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.115D Standard Errors of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Demographic and Socioeconomic Characteristics: 2006 and 2007

Demographic/Socioeconomic Characteristic	MEAN AGE					
	Alcohol		Marijuana		Cigarettes	
	2006	2007	2006	2007	2006	2007
TOTAL	0.04	0.04	0.05	0.05	0.06	0.06
RACE¹						
American Indian or Alaska Native	0.24	0.20	0.27	0.76	0.24	0.35
Asian or Pacific Islander	0.22	0.27	0.27	0.30	0.39	0.34
Asian Only	0.23	0.31	0.29	0.30	0.42	0.37
Native Hawaiian or Other Pacific Islander Only	0.42	0.24	*	*	*	*
Black or African American	0.13	0.11	0.18	0.13	0.17	0.17
White	0.04	0.04	0.06	0.06	0.07	0.06
Two or More Races	0.22	0.23	0.31	0.22	0.29	0.27
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.10	0.10	0.15	0.16	0.17	0.15
Not Hispanic or Latino	0.04	0.04	0.06	0.05	0.06	0.07
Black or African American	0.13	0.12	0.18	0.13	0.17	0.17
White	0.04	0.05	0.06	0.06	0.07	0.07
GENDER						
Male	0.06	0.05	0.07	0.07	0.09	0.09
Female	0.05	0.05	0.07	0.07	0.08	0.08
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²						
Less Than 100%	0.11	0.09	0.15	0.15	0.13	0.16
100-199%	0.09	0.09	0.12	0.12	0.16	0.14
200% or More	0.05	0.05	0.06	0.06	0.07	0.07

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used substance(s) for the first time in the 12 months prior to date of interview.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9a, 26-9b, and 27-4a (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.115P P Values from Tests of Differences of Mean Age at First Use among Past Year Initiates of Substance Use Aged 12 or Older Who Initiated Use Prior to the Age of 18, by Demographic and Socioeconomic Characteristics: 2007 Versus 2006

Demographic/Socioeconomic Characteristic	MEAN AGE					
	Alcohol		Marijuana		Cigarettes	
	2006	2007	2006	2007	2006	2007
TOTAL	0.7172		0.3747		0.5745	
RACE¹						
American Indian or Alaska Native	0.3322		0.9221		0.7366	
Asian or Pacific Islander	0.0398		0.5491		0.1743	
Asian Only	0.0860		0.7211		0.1793	
Native Hawaiian or Other Pacific Islander Only	0.4450		*		*	
Black or African American	0.6945		0.8272		0.7965	
White	0.3681		0.1571		0.5038	
Two or More Races	0.1815		0.0267		0.0541	
HISPANIC ORIGIN AND RACE						
Hispanic or Latino	0.7189		0.4311		0.6904	
Not Hispanic or Latino	0.8202		0.5588		0.6895	
Black or African American	0.5014		0.8789		0.9061	
White	0.3329		0.2449		0.5700	
GENDER						
Male	0.1754		1.0000		0.6871	
Female	0.4256		0.1973		0.2137	
POVERTY LEVEL (% of Census Bureau Poverty Threshold)²						
Less Than 100%	0.9590		0.3265		0.7204	
100-199%	0.4420		0.2380		0.1246	
200% or More	0.9187		0.2003		0.8557	

*Low precision; no estimate reported.

NOTE: Past Year Initiates are defined as persons who used substance(s) for the first time in the 12 months prior to date of interview.

NOTE: Estimates shown on this table correspond to Healthy People 2010 Objective Numbers 26-9a, 26-9b, and 27-4a (<http://www.healthypeople.gov/>).

¹ These racial categories do not distinguish among ethnic origin (i.e., Hispanic or Latino origin), so they include respondents who are either Hispanic or not Hispanic.

² Estimates are based on a definition of Poverty Level that incorporates information on family income, size, and composition and is calculated as a percentage of the U.S. Census Bureau's poverty thresholds.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.