

Table 7.11C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	208	219	227	232	209	203	199	195
Marijuana and Hashish	186	192	203	188	169	166	138	140
Cocaine	105	95	75	75	69	53	47	58
Crack	58	42	55	32	57	37	25	47
Heroin	58	12	11	7	44	28	20	18
Hallucinogens	38	38	39	45	27	32	32	30
LSD	11	10	12	26	17	17	6	7
PCP	5	2	6	8	*	2	*	9
Ecstasy	30	30	27	31	19	30	24	14
Inhalants	26	45	31	28	52	22	29	35
Nonmedical Use of Psychotherapeutics ^{3,4}	122	110	100	123	128	131	134	139
Pain Relievers	97	100	87	87	110	109	126	126
OxyContin [®]	19	17	21	24	11	10	18	43
Tranquilizers	48	55	56	62	66	66	54	70
Stimulants ⁴	56	48	47	49	63	37	40	55
Methamphetamine ⁴	48	42	37	40	41	28	35	48
Sedatives	36	20	17	51	39	23	40	33
ILLCIT DRUGS OTHER THAN MARIJUANA²	148	141	128	148	145	143	146	152

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.11D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	1.06	1.03	0.53	0.53	0.30	0.29	0.16	0.16
Marijuana and Hashish	0.97	0.97	0.49	0.45	0.24	0.24	0.11	0.11
Cocaine	0.60	0.54	0.19	0.18	0.10	0.08	0.04	0.05
Crack	0.34	0.25	0.14	0.08	0.08	0.05	0.02	0.04
Heroin	0.34	0.07	0.03	0.02	0.06	0.04	0.02	0.01
Hallucinogens	0.22	0.22	0.10	0.11	0.04	0.05	0.03	0.02
LSD	0.06	0.06	0.03	0.06	0.02	0.02	0.00	0.01
PCP	0.03	0.01	0.02	0.02	*	0.00	*	0.01
Ecstasy	0.18	0.17	0.07	0.08	0.03	0.04	0.02	0.01
Inhalants	0.15	0.26	0.08	0.07	0.08	0.03	0.02	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.68	0.61	0.25	0.29	0.18	0.19	0.11	0.12
Pain Relievers	0.55	0.56	0.22	0.21	0.16	0.16	0.10	0.10
OxyContin [®]	0.11	0.10	0.05	0.06	0.02	0.01	0.01	0.04
Tranquilizers	0.28	0.32	0.14	0.15	0.10	0.10	0.04	0.06
Stimulants ⁴	0.33	0.28	0.12	0.12	0.09	0.05	0.03	0.05
Methamphetamine ⁴	0.28	0.25	0.09	0.10	0.06	0.04	0.03	0.04
Sedatives	0.21	0.12	0.04	0.13	0.06	0.03	0.03	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA²	0.81	0.75	0.31	0.35	0.21	0.21	0.12	0.13

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.11P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS ²	0.3725		0.9399		0.0266		0.7389	
Marijuana and Hashish	0.2278		0.6833		0.3437		0.6993	
Cocaine	0.0680		0.3404		0.2006		0.1626	
Crack	0.6482		0.1976		0.1741		0.0414	
Heroin	0.0310		0.2649		0.8937		0.1645	
Hallucinogens	0.7102		0.7656		0.6368		0.9824	
LSD	0.8669		0.4514		0.4720		0.5046	
PCP	0.3087		0.9369		*		*	
Ecstasy	0.3175		0.9847		0.6533		0.8288	
Inhalants	0.6909		0.8016		0.0070		0.8231	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.3608		0.8116		0.2899		0.6232	
Pain Relievers	0.7237		0.5309		0.4487		0.8066	
OxyContin®	0.4789		0.5019		0.8164		0.3225	
Tranquilizers	0.3477		0.4782		0.9215		0.5744	
Stimulants ⁴	0.1368		0.1113		0.0780		0.9718	
Methamphetamine ⁴	0.2279		0.2536		0.2487		0.8135	
Sedatives	0.5397		0.1681		0.1100		0.7275	
ILLCIT DRUGS OTHER THAN MARIJUANA ²	0.3315		0.8143		0.0708		0.4921	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.12C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	26	27	37	37	25	24	46	51
Marijuana and Hashish	24	25	34	34	21	21	31	34
Cocaine	7	7	8	7	5	4	6	7
Crack	3	3	2	3	2	1	*	3
Heroin	3	1	2	*	1	1	3	*
Hallucinogens	9	9	9	11	5	5	11	8
LSD	4	3	4	5	2	2	3	4
PCP	3	1	3	3	*	1	2	3
Ecstasy	7	6	6	7	3	3	6	5
Inhalants	7	9	11	10	11	7	19	21
Nonmedical Use of Psychotherapeutics ^{3,4}	16	16	18	19	10	14	30	29
Pain Relievers	14	16	16	16	9	13	27	25
OxyContin [®]	2	5	3	4	2	3	4	4
Tranquilizers	8	10	8	10	5	5	7	11
Stimulants ⁴	6	7	8	7	5	3	13	9
Methamphetamine ⁴	2	5	4	3	2	1	9	5
Sedatives	2	2	3	3	2	1	7	8
ILLCIT DRUGS OTHER THAN MARIJUANA²	18	18	22	22	16	15	36	38

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.12D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	2.78	2.85	1.51	1.53	1.34	1.40	0.21	0.23
Marijuana and Hashish	2.87	2.87	1.40	1.49	1.21	1.22	0.15	0.16
Cocaine	1.13	1.20	0.40	0.39	0.33	0.23	0.03	0.03
Crack	0.49	0.56	0.08	0.15	0.14	0.09	*	0.01
Heroin	0.52	0.16	0.12	*	0.08	0.09	0.01	*
Hallucinogens	1.46	1.50	0.45	0.58	0.31	0.32	0.05	0.04
LSD	0.67	0.53	0.22	0.28	0.14	0.13	0.01	0.02
PCP	0.51	0.12	0.13	0.17	*	0.09	0.01	0.01
Ecstasy	1.13	1.01	0.31	0.38	0.21	0.21	0.03	0.02
Inhalants	1.07	1.49	0.56	0.52	0.64	0.42	0.09	0.10
Nonmedical Use of Psychotherapeutics ^{3,4}	2.39	2.41	0.85	0.94	0.63	0.84	0.14	0.14
Pain Relievers	2.19	2.33	0.78	0.82	0.55	0.83	0.12	0.12
OxyContin [®]	0.41	0.88	0.15	0.21	0.12	0.18	0.02	0.02
Tranquilizers	1.21	1.64	0.39	0.53	0.29	0.30	0.04	0.05
Stimulants ⁴	0.92	1.08	0.41	0.40	0.29	0.19	0.06	0.04
Methamphetamine ⁴	0.37	0.79	0.19	0.18	0.13	0.08	0.04	0.02
Sedatives	0.36	0.29	0.13	0.14	0.13	0.09	0.03	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	2.53	2.58	1.04	1.07	0.92	0.92	0.17	0.17

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.12P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS ²	0.5274		0.9770		0.0728		0.5828	
Marijuana and Hashish	0.2876		0.8240		0.4477		0.3707	
Cocaine	0.8497		0.7352		0.3497		0.5289	
Crack	0.7337		0.9599		0.5670		*	
Heroin	0.1673		*		0.9872		*	
Hallucinogens	0.3918		0.0836		0.9248		0.2635	
LSD	0.2953		0.7322		0.8853		0.7297	
PCP	0.2042		0.8285		*		0.8764	
Ecstasy	0.2448		0.9416		0.9053		0.5368	
Inhalants	0.3236		0.4942		0.0032		0.8767	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.3957		0.9435		0.4634		0.5131	
Pain Relievers	0.3579		0.6461		0.1130		0.3136	
OxyContin®	0.0400		0.1918		0.3021		0.6955	
Tranquilizers	0.0476		0.2475		0.6626		0.0591	
Stimulants ⁴	0.9104		0.4526		0.2667		0.1231	
Methamphetamine ⁴	0.3026		0.2579		0.4280		0.1196	
Sedatives	0.5177		0.4018		0.3676		0.7939	
ILLCIT DRUGS OTHER THAN MARIJUANA ²	0.5166		0.7568		0.1393		0.7676	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.13C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	88	90	84	91	51	46	48	45
Marijuana and Hashish	82	84	76	81	44	43	39	39
Cocaine	32	27	24	24	11	9	8	7
Crack	8	6	9	5	3	5	3	5
Heroin	8	3	5	7	2	6	5	4
Hallucinogens	28	25	27	21	15	8	10	9
LSD	9	9	5	6	6	1	0	6
PCP	4	2	5	2	*	*	2	5
Ecstasy	21	15	24	17	12	5	5	6
Inhalants	10	9	10	12	6	6	8	9
Nonmedical Use of Psychotherapeutics ^{3,4}	50	48	48	45	26	24	27	23
Pain Relievers	40	43	41	37	23	19	23	19
OxyContin [®]	10	10	13	12	4	7	4	5
Tranquilizers	25	28	25	20	15	14	13	13
Stimulants ⁴	28	19	20	20	8	13	10	10
Methamphetamine ⁴	15	11	15	10	5	10	7	6
Sedatives	11	9	3	6	4	2	5	6
ILLCIT DRUGS OTHER THAN MARIJUANA²	61	56	56	53	32	26	31	26

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.13D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	1.10	1.17	0.73	0.78	0.72	0.68	0.36	0.34
Marijuana and Hashish	1.11	1.17	0.71	0.72	0.65	0.64	0.30	0.30
Cocaine	0.59	0.54	0.27	0.26	0.17	0.15	0.06	0.05
Crack	0.15	0.13	0.10	0.06	0.05	0.08	0.02	0.04
Heroin	0.16	0.06	0.06	0.08	0.03	0.09	0.04	0.04
Hallucinogens	0.52	0.51	0.31	0.24	0.22	0.13	0.08	0.07
LSD	0.18	0.18	0.06	0.07	0.10	0.01	0.00	0.05
PCP	0.08	0.04	0.05	0.02	*	*	0.01	0.04
Ecstasy	0.39	0.31	0.27	0.19	0.18	0.09	0.04	0.05
Inhalants	0.19	0.20	0.11	0.14	0.10	0.09	0.07	0.07
Nonmedical Use of Psychotherapeutics ^{3,4}	0.88	0.87	0.51	0.47	0.39	0.37	0.21	0.18
Pain Relievers	0.74	0.80	0.44	0.40	0.34	0.30	0.18	0.15
OxyContin [®]	0.19	0.21	0.14	0.14	0.06	0.10	0.03	0.04
Tranquilizers	0.48	0.56	0.28	0.22	0.23	0.21	0.10	0.10
Stimulants ⁴	0.53	0.38	0.22	0.23	0.12	0.20	0.08	0.08
Methamphetamine ⁴	0.28	0.22	0.17	0.11	0.08	0.15	0.05	0.05
Sedatives	0.21	0.19	0.03	0.07	0.07	0.03	0.04	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.01	0.98	0.57	0.54	0.46	0.39	0.24	0.20

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.13P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS ²	0.2356		0.5520		0.7157		0.3641	
Marijuana and Hashish	0.0926		0.5755		0.9429		0.4046	
Cocaine	0.0463		0.1383		0.4297		0.8883	
Crack	0.7555		0.1099		0.6825		0.5352	
Heroin	0.0576		0.8096		0.1245		0.8240	
Hallucinogens	0.6905		0.4096		0.0474		0.9943	
LSD	0.4131		0.3798		0.0541		0.1082	
PCP	0.7782		0.3052		*		0.2106	
Ecstasy	0.0513		0.3752		0.1441		0.6259	
Inhalants	0.7769		0.7951		0.6830		0.8153	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.3583		0.2837		0.8531		0.8939	
Pain Relievers	0.6384		0.2872		0.6542		0.6049	
OxyContin [®]	0.6333		0.9148		0.2877		0.3554	
Tranquilizers	0.7757		0.0303		0.9025		0.7284	
Stimulants ⁴	0.0622		0.0773		0.3033		0.9324	
Methamphetamine ⁴	0.2342		0.0090		0.3914		0.4141	
Sedatives	0.3056		0.0332		0.1175		0.8100	
ILLCIT DRUGS OTHER THAN MARIJUANA ²	0.2544		0.3556		0.1990		0.6085	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.14C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	183	191	205	206	197	196	182	184
Marijuana and Hashish	166	162	182	164	160	160	126	133
Cocaine	100	91	70	70	68	52	46	57
Crack	57	42	54	32	57	36	25	46
Heroin	58	12	7	*	44	27	19	17
Hallucinogens	25	27	27	38	22	31	28	28
LSD	4	3	10	24	15	16	5	*
PCP	*	*	3	7	*	*	*	6
Ecstasy	21	25	13	24	13	30	23	11
Inhalants	23	42	26	23	50	20	21	26
Nonmedical Use of Psychotherapeutics ^{3,4}	108	98	87	114	124	126	126	136
Pain Relievers	87	89	75	77	107	106	118	122
OxyContin [®]	16	12	15	21	10	7	17	43
Tranquilizers	41	49	50	60	65	65	52	69
Stimulants ⁴	47	43	40	43	62	34	37	53
Methamphetamine ⁴	45	41	33	39	40	26	33	47
Sedatives	34	18	16	51	38	23	39	32
ILLCIT DRUGS OTHER THAN MARIJUANA²	130	130	115	136	138	137	136	145

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.14D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS²	1.43	1.38	0.67	0.66	0.31	0.31	0.21	0.21
Marijuana and Hashish	1.33	1.24	0.61	0.54	0.26	0.26	0.14	0.15
Cocaine	0.87	0.76	0.24	0.23	0.11	0.08	0.05	0.07
Crack	0.50	0.36	0.19	0.11	0.09	0.06	0.03	0.05
Heroin	0.51	0.10	0.02	*	0.07	0.04	0.02	0.02
Hallucinogens	0.22	0.23	0.09	0.12	0.04	0.05	0.03	0.03
LSD	0.04	0.03	0.04	0.08	0.03	0.03	0.01	*
PCP	*	*	0.01	0.02	*	*	*	0.01
Ecstasy	0.18	0.22	0.04	0.08	0.02	0.05	0.03	0.01
Inhalants	0.21	0.36	0.09	0.08	0.08	0.03	0.02	0.03
Nonmedical Use of Psychotherapeutics ^{3,4}	0.92	0.81	0.29	0.37	0.20	0.20	0.14	0.16
Pain Relievers	0.75	0.74	0.26	0.25	0.18	0.17	0.14	0.14
OxyContin [®]	0.14	0.11	0.05	0.07	0.02	0.01	0.02	0.05
Tranquilizers	0.36	0.42	0.17	0.20	0.11	0.11	0.06	0.08
Stimulants ⁴	0.41	0.37	0.14	0.14	0.10	0.06	0.04	0.06
Methamphetamine ⁴	0.40	0.35	0.11	0.13	0.07	0.04	0.04	0.05
Sedatives	0.30	0.15	0.06	0.17	0.06	0.04	0.04	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA²	1.10	1.05	0.38	0.44	0.23	0.22	0.16	0.17

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.14P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS ²	0.2538		0.8831		0.0496		0.6845	
Marijuana and Hashish	0.0753		0.6647		0.4222		0.6944	
Cocaine	0.2001		0.6453		0.2709		0.1735	
Crack	0.6284		0.2918		0.1693		0.0558	
Heroin	0.0680		*		0.7474		0.2125	
Hallucinogens	0.6274		0.5121		0.7168		0.7205	
LSD	0.9737		0.6124		0.8248		*	
PCP	*		0.4930		*		*	
Ecstasy	0.4930		0.2398		0.2661		0.8291	
Inhalants	0.8819		0.9389		0.0395		0.9206	
Nonmedical Use of Psychotherapeutics ^{3,4}	0.5811		0.7317		0.2682		0.4989	
Pain Relievers	0.5302		0.9512		0.4066		0.7050	
OxyContin®	0.9233		0.4750		0.5428		0.3549	
Tranquilizers	0.4930		0.9705		0.9765		0.7201	
Stimulants ⁴	0.5392		0.4590		0.0504		0.7296	
Methamphetamine ⁴	0.3265		0.8335		0.1785		0.8802	
Sedatives	0.3345		0.2285		0.1625		0.7902	
ILLCIT DRUGS OTHER THAN MARIJUANA ²	0.6266		0.7229		0.1550		0.3677	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

⁴ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.15C Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	Total		UNDERAGE AND LEGAL AGE DRINKING GROUP			
			12-20		21 or Older	
	2006	2007	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month						
ILLCIT DRUGS^{1,2}	385	403	109	114	357	377
Marijuana or Hashish	335	330	99	105	311	304
Cocaine (Including Crack)	146	134	30	26	142	131
Heroin	74	31	9	6	73	31
Hallucinogens	61	68	35	29	49	61
Inhalants	65	57	22	19	62	54
Pain Relievers ³	172	173	45	50	165	165
Tranquilizers ³	101	107	26	27	98	105
Stimulants ^{3,4}	96	77	27	21	91	75
Methamphetamine ⁴	72	65	17	11	71	64
Sedatives ³	56	60	11	7	55	60
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	245	266	68	65	229	255
Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month						
ILLCIT DRUGS^{1,2}	253	230	66	68	240	217
Marijuana or Hashish	220	209	64	67	208	194
Cocaine (Including Crack)	106	76	12	12	105	75
Heroin	45	18	1	4	45	17
Hallucinogens	21	24	5	9	20	23
Inhalants	11	6	6	5	9	*
Pain Relievers ³	55	58	17	13	52	57
Tranquilizers ³	25	18	7	5	24	17
Stimulants ³	13	10	6	4	11	9
Methamphetamine ⁶	39	37	9	7	38	36
Sedatives ³	16	18	1	2	16	18
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	130	103	25	22	127	102

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.15D Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: Standard Errors of Percentages, 2006 and 2007

Drug	Total		UNDERAGE AND LEGAL AGE DRINKING GROUP			
			12-20		21 or Older	
	2006	2007	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month						
ILLCIT DRUGS^{1,2}	0.29	0.30	0.71	0.75	0.30	0.31
Marijuana or Hashish	0.26	0.25	0.70	0.72	0.26	0.26
Cocaine (Including Crack)	0.12	0.10	0.27	0.23	0.12	0.11
Heroin	0.06	0.02	0.09	0.05	0.06	0.03
Hallucinogens	0.05	0.05	0.31	0.26	0.04	0.05
Inhalants	0.05	0.04	0.20	0.18	0.05	0.05
Pain Relievers ³	0.14	0.13	0.39	0.43	0.14	0.14
Tranquilizers ³	0.08	0.08	0.24	0.25	0.09	0.09
Stimulants ^{3,4}	0.08	0.06	0.25	0.19	0.08	0.06
Methamphetamine ⁴	0.06	0.05	0.16	0.11	0.06	0.05
Sedatives ³	0.04	0.05	0.10	0.07	0.05	0.05
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.19	0.20	0.53	0.53	0.20	0.21
Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month						
Drug⁵						
ILLCIT DRUGS^{1,2}	0.20	0.18	0.57	0.55	0.21	0.19
Marijuana or Hashish	0.18	0.16	0.56	0.55	0.18	0.17
Cocaine (Including Crack)	0.08	0.06	0.12	0.12	0.09	0.06
Heroin	0.04	0.01	0.01	0.03	0.04	0.01
Hallucinogens	0.02	0.02	0.05	0.08	0.02	0.02
Inhalants	0.01	0.00	0.06	0.05	0.01	*
Pain Relievers ³	0.04	0.05	0.16	0.12	0.05	0.05
Tranquilizers ³	0.02	0.01	0.07	0.05	0.02	0.01
Stimulants ³	0.01	0.01	0.06	0.04	0.01	0.01
Methamphetamine ⁶	0.03	0.03	0.08	0.06	0.03	0.03
Sedatives ³	0.01	0.01	0.01	0.01	0.01	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.10	0.08	0.22	0.20	0.11	0.09

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.15P Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Underage and Legal Age Drinking Groups: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	Total		UNDERAGE AND LEGAL AGE DRINKING GROUP	
			12-20	21 or Older
	2006	2007	2006	2007
Used Specified Drug Anytime in the Past Month				
ILLCIT DRUGS^{1,2}	0.1282		0.4385	0.1888
Marijuana or Hashish	0.1879		0.9386	0.1889
Cocaine (Including Crack)	0.0177		0.0055	0.0702
Heroin	0.0509		0.0614	0.0881
Hallucinogens	0.8344		0.1762	0.5674
Inhalants	0.1027		0.0657	0.2995
Pain Relievers ³	0.5589		0.5833	0.6938
Tranquilizers ³	1.0000		0.7523	0.9048
Stimulants ^{3,4}	0.0051		0.0048	0.0406
Methamphetamine ⁴	0.0426		0.0215	0.1198
Sedatives ³	0.7705		0.4050	0.8775
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.0930		0.0038	0.3380
Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month				
ILLCIT DRUGS^{1,2}	0.3459		0.7534	0.3176
Marijuana or Hashish	0.4458		0.7293	0.3999
Cocaine (Including Crack)	0.0905		0.9312	0.0854
Heroin	0.2096		0.2864	0.1744
Hallucinogens	0.5394		0.1298	0.8715
Inhalants	0.2504		0.9234	*
Pain Relievers ³	0.8455		0.0482	0.4438
Tranquilizers ³	0.5479		0.4236	0.7053
Stimulants ³	0.7969		0.5938	0.9904
Methamphetamine ⁶	0.4432		0.3500	0.5636
Sedatives ³	0.2227		0.5658	0.2383
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.1317		0.1552	0.2209

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.16C Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	AGE GROUP					
	12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007
	Used Specified Drug Anytime in the Past Month					
ILLCIT DRUGS^{1,2}	51	52	143	153	338	347
Marijuana or Hashish	46	47	131	140	294	279
Cocaine (Including Crack)	12	11	41	38	140	127
Heroin	4	2	10	10	73	30
Hallucinogens	14	15	42	34	42	56
Inhalants	17	15	16	17	61	52
Pain Relievers ³	24	25	63	62	158	157
Tranquilizers ³	12	16	40	37	91	101
Stimulants ^{3,4}	11	10	33	30	89	70
Methamphetamine ⁴	5	6	20	18	70	62
Sedatives ³	4	3	12	12	54	59
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	33	32	89	87	220	243
Drug⁵	Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month					
ILLCIT DRUGS^{1,2}	31	32	91	94	229	200
Marijuana or Hashish	30	31	86	91	197	178
Cocaine (Including Crack)	6	5	21	17	102	74
Heroin	1	2	1	7	45	16
Hallucinogens	3	6	7	11	19	21
Inhalants	6	4	2	4	9	*
Pain Relievers ³	7	8	24	24	48	52
Tranquilizers ³	3	4	9	12	23	12
Stimulants ³	2	2	8	7	9	7
Methamphetamine ⁶	2	2	10	10	37	35
Sedatives ³	1	2	1	5	16	17
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	12	12	36	33	122	98

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.16D Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Drug	AGE GROUP					
	12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007
	Used Specified Drug Anytime in the Past Month					
ILLICIT DRUGS^{1,2}	0.96	1.03	0.51	0.53	0.32	0.32
Marijuana or Hashish	0.91	0.98	0.50	0.52	0.28	0.27
Cocaine (Including Crack)	0.28	0.26	0.20	0.18	0.14	0.12
Heroin	0.10	0.04	0.05	0.05	0.07	0.03
Hallucinogens	0.32	0.37	0.20	0.17	0.04	0.05
Inhalants	0.40	0.37	0.08	0.08	0.06	0.05
Pain Relievers ³	0.55	0.59	0.29	0.28	0.16	0.15
Tranquilizers ³	0.29	0.39	0.19	0.18	0.09	0.10
Stimulants ^{3,4}	0.26	0.26	0.16	0.15	0.09	0.07
Methamphetamine ⁴	0.12	0.15	0.10	0.09	0.07	0.06
Sedatives ³	0.10	0.09	0.06	0.06	0.05	0.06
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.71	0.73	0.39	0.38	0.22	0.23
Drug⁵	Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month					
ILLICIT DRUGS^{1,2}	0.70	0.75	0.42	0.40	0.23	0.19
Marijuana or Hashish	0.68	0.74	0.40	0.39	0.19	0.17
Cocaine (Including Crack)	0.14	0.13	0.10	0.09	0.10	0.07
Heroin	0.02	0.04	0.01	0.03	0.04	0.02
Hallucinogens	0.07	0.16	0.03	0.06	0.02	0.02
Inhalants	0.14	0.09	0.01	0.02	0.01	*
Pain Relievers ³	0.17	0.20	0.12	0.12	0.05	0.05
Tranquilizers ³	0.07	0.11	0.04	0.06	0.02	0.01
Stimulants ³	0.05	0.04	0.04	0.03	0.01	0.01
Methamphetamine ⁶	0.05	0.05	0.05	0.05	0.04	0.03
Sedatives ³	0.02	0.04	0.01	0.03	0.02	0.02
ILLICIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.28	0.30	0.17	0.16	0.12	0.09

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.16P Types of Illicit Drug Use in the Past Month and Types of Illicit Drugs Used in the Past Month with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol among Past Month Alcohol Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	AGE GROUP					
	12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007
	Used Specified Drug Anytime in the Past Month					
ILLCIT DRUGS^{1,2}	0.4619		0.5383		0.1418	
Marijuana or Hashish	0.8511		0.3020		0.1216	
Cocaine (Including Crack)	0.6294		0.0075		0.1151	
Heroin	0.0513		0.7494		0.0701	
Hallucinogens	0.4504		0.1310		0.3663	
Inhalants	0.1205		0.8598		0.2097	
Pain Relievers ³	0.3333		0.1838		0.9090	
Tranquilizers ³	0.0811		0.2177		0.7505	
Stimulants ^{3,4}	0.3092		0.0259		0.0541	
Methamphetamine ⁴	0.7789		0.0259		0.1600	
Sedatives ³	0.1908		0.2304		0.6545	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.5368		0.0508		0.4033	
Drug⁵	Used Specified Drug with Alcohol or within 2 Hours of Alcohol Use on the Most Recent Use of Alcohol in the Past Month					
ILLCIT DRUGS^{1,2}	0.3137		0.3033		0.1823	
Marijuana or Hashish	0.4882		0.2409		0.2259	
Cocaine (Including Crack)	0.9744		0.3167		0.1274	
Heroin	0.4167		0.1820		0.1271	
Hallucinogens	0.1340		0.2341		0.8496	
Inhalants	0.5699		0.6374		*	
Pain Relievers ³	0.8535		0.7294		0.6727	
Tranquilizers ³	0.2948		0.6488		0.2669	
Stimulants ³	0.7039		0.9136		0.9143	
Methamphetamine ⁶	0.7740		0.8915		0.4605	
Sedatives ³	0.5414		0.0273		0.4875	
ILLCIT DRUGS OTHER THAN MARIJUANA^{1,2}	0.6675		0.4473		0.1745	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically.

² Estimates in these designated rows do not include data from the methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

³ Refers to nonmedical use; does not include over-the-counter drugs.

⁴ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

⁵ Estimates are based on non-imputed responses to questions in the core and noncore modules; therefore, respondents with unknown responses to the questions about other drugs used with alcohol were excluded. Respondents could indicate multiple other drugs used with alcohol; thus, these response categories are not mutually exclusive.

⁶ Estimates in the designated rows include data from methamphetamine use items added in 2005 and 2006.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.