

Table 7.17C Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	300	306	141	124	224	245	136	153
Marijuana and Hashish	264	258	112	96	189	199	74	97
Cocaine	131	121	37	52	58	49	29	25
Crack	92	61	23	46	32	21	10	5
Heroin	72	31	19	18	20	4	4	*
Hallucinogens	49	60	14	25	36	33	28	17
LSD	18	32	5	5	15	4	*	5
PCP	7	9	*	8	4	*	3	4
Ecstasy	38	46	7	8	25	27	23	11
Inhalants	53	35	18	4	38	44	23	34
Nonmedical Use of Psychotherapeutics ^{2,3}	157	151	81	83	128	143	107	119
Pain Relievers	132	132	73	72	110	113	101	108
OxyContin [®]	26	30	18	40	17	9	3	17
Tranquilizers	76	74	29	47	66	74	45	52
Stimulants ³	80	69	37	48	52	31	16	27
Methamphetamine ³	64	63	33	47	34	17	12	9
Sedatives	44	26	16	15	34	54	37	29
ILLCIT DRUGS OTHER THAN MARIJUANA¹	199	195	88	99	142	166	115	124

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.17D Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.64	0.66	0.67	0.61	0.26	0.27	0.13	0.15
Marijuana and Hashish	0.57	0.58	0.54	0.48	0.22	0.22	0.07	0.10
Cocaine	0.31	0.29	0.18	0.26	0.07	0.06	0.03	0.02
Crack	0.22	0.15	0.11	0.23	0.04	0.02	0.01	0.00
Heroin	0.17	0.08	0.09	0.09	0.02	0.00	0.00	*
Hallucinogens	0.12	0.15	0.07	0.13	0.04	0.04	0.03	0.02
LSD	0.04	0.08	0.03	0.03	0.02	0.01	*	0.00
PCP	0.02	0.02	*	0.04	0.00	*	0.00	0.00
Ecstasy	0.09	0.12	0.04	0.04	0.03	0.03	0.02	0.01
Inhalants	0.13	0.09	0.09	0.02	0.05	0.05	0.02	0.03
Nonmedical Use of Psychotherapeutics ^{2,3}	0.37	0.35	0.39	0.41	0.15	0.16	0.11	0.12
Pain Relievers	0.32	0.31	0.35	0.36	0.13	0.13	0.10	0.11
OxyContin [®]	0.06	0.08	0.09	0.20	0.02	0.01	0.00	0.02
Tranquilizers	0.19	0.18	0.14	0.24	0.08	0.08	0.04	0.05
Stimulants ³	0.19	0.17	0.18	0.24	0.06	0.04	0.02	0.03
Methamphetamine ³	0.15	0.15	0.16	0.23	0.04	0.02	0.01	0.01
Sedatives	0.11	0.06	0.08	0.08	0.04	0.06	0.04	0.03
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.45	0.44	0.42	0.48	0.17	0.19	0.11	0.12

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.17P Types of Illicit Drug Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.5891		0.9742		0.2427		0.4224	
Marijuana and Hashish	0.5393		0.8925		0.4490		0.3728	
Cocaine	0.1231		0.0267		0.0427		0.5370	
Crack	0.1912		0.0277		0.1868		0.7592	
Heroin	0.0878		0.2905		0.3184		*	
Hallucinogens	0.7018		0.3457		0.4809		0.4197	
LSD	0.3815		0.9803		0.1447		*	
PCP	0.8807		*		*		0.9208	
Ecstasy	0.9601		0.7894		0.7324		0.7533	
Inhalants	0.1607		0.0211		0.4144		0.4548	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.3979		0.9620		0.5243		0.4921	
Pain Relievers	0.9563		0.6812		0.6305		0.9084	
OxyContin [®]	0.2020		0.5613		0.9610		0.1719	
Tranquilizers	0.7298		0.7581		0.5649		0.5805	
Stimulants ³	0.1431		0.9250		0.0046		0.9438	
Methamphetamine ³	0.2695		0.8479		0.0252		0.1680	
Sedatives	0.5082		0.7879		0.8435		0.8110	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.2927		0.4930		0.3244		0.6292	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.18C Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	39	40	26	23	33	32	37	42
Marijuana and Hashish	35	38	24	22	28	28	20	24
Cocaine	10	10	4	4	6	4	4	6
Crack	4	4	0	1	2	1	*	3
Heroin	4	1	*	0	2	1	3	*
Hallucinogens	13	14	8	6	5	7	7	6
LSD	6	6	2	4	2	1	2	2
PCP	4	3	*	*	*	1	2	3
Ecstasy	9	8	4	4	2	5	4	3
Inhalants	13	12	10	4	12	10	16	20
Nonmedical Use of Psychotherapeutics ^{2,3}	22	23	13	11	15	15	27	26
Pain Relievers	20	20	10	10	13	15	24	23
OxyContin [®]	4	6	4	3	2	4	2	3
Tranquilizers	12	15	4	4	4	6	6	11
Stimulants ³	8	9	7	6	7	4	11	7
Methamphetamine ³	5	5	6	3	0	3	7	4
Sedatives	4	3	1	3	2	2	7	8
ILLCIT DRUGS OTHER THAN MARIJUANA¹	27	26	16	13	19	18	31	35

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.18D Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	1.63	1.90	2.04	1.99	1.07	1.12	0.18	0.20
Marijuana and Hashish	1.64	1.88	1.92	1.89	0.95	1.01	0.10	0.12
Cocaine	0.62	0.63	0.39	0.43	0.24	0.18	0.02	0.03
Crack	0.23	0.28	0.03	0.15	0.06	0.06	*	0.01
Heroin	0.24	0.06	*	0.03	0.06	0.06	0.02	*
Hallucinogens	0.76	0.88	0.82	0.62	0.19	0.28	0.03	0.03
LSD	0.37	0.41	0.17	0.41	0.06	0.06	0.01	0.01
PCP	0.25	0.21	*	*	*	0.06	0.01	0.01
Ecstasy	0.57	0.54	0.43	0.38	0.08	0.22	0.02	0.01
Inhalants	0.75	0.74	0.93	0.41	0.43	0.39	0.08	0.10
Nonmedical Use of Psychotherapeutics ^{2,3}	1.26	1.38	1.19	1.09	0.54	0.60	0.13	0.13
Pain Relievers	1.16	1.23	0.97	0.98	0.51	0.58	0.12	0.11
OxyContin [®]	0.25	0.40	0.38	0.35	0.08	0.15	0.01	0.01
Tranquilizers	0.70	0.93	0.42	0.40	0.15	0.24	0.03	0.05
Stimulants ³	0.52	0.61	0.70	0.58	0.26	0.18	0.05	0.03
Methamphetamine ³	0.30	0.34	0.56	0.32	0.02	0.12	0.04	0.02
Sedatives	0.25	0.19	0.11	0.35	0.06	0.07	0.04	0.04
ILLCIT DRUGS OTHER THAN MARIJUANA¹	1.45	1.50	1.49	1.30	0.70	0.68	0.15	0.17

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.18P Types of Illicit Drug Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.9761		0.8040		0.4360		0.2942	
Marijuana and Hashish	0.4120		0.6996		0.7417		0.1858	
Cocaine	0.9449		0.8959		0.4510		0.4367	
Crack	0.8828		0.2677		0.8506		*	
Heroin	0.0389		*		0.7729		*	
Hallucinogens	0.9377		0.7626		0.0619		0.2080	
LSD	0.8643		0.7023		0.7509		0.9116	
PCP	0.5556		*		*		0.8747	
Ecstasy	0.1261		0.8871		0.0986		0.4718	
Inhalants	0.4007		0.0225		0.1536		0.4515	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4529		0.7902		0.5216		0.4401	
Pain Relievers	0.5467		0.7718		0.3007		0.2385	
OxyContin [®]	0.0307		0.8517		0.1636		0.7342	
Tranquilizers	0.1681		0.8363		0.1134		0.0409	
Stimulants ³	0.8313		0.3100		0.1396		0.0111	
Methamphetamine ³	0.3834		0.7096		0.1598		0.0984	
Sedatives	0.1115		0.2291		0.7454		0.4958	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.9363		0.6033		0.4399		1.0000	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.19C Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	115	115	37	37	69	81	28	27
Marijuana and Hashish	109	106	33	34	60	73	21	24
Cocaine	39	35	7	6	17	14	2	3
Crack	11	10	3	5	4	1	*	*
Heroin	10	9	5	4	1	3	*	*
Hallucinogens	37	30	8	7	19	15	6	5
LSD	10	10	0	4	7	4	*	5
PCP	6	3	*	4	*	*	2	3
Ecstasy	31	22	5	5	14	7	*	3
Inhalants	14	15	5	1	7	7	6	9
Nonmedical Use of Psychotherapeutics ^{2,3}	63	59	20	18	38	37	18	16
Pain Relievers	53	53	17	15	32	31	15	14
OxyContin [®]	16	16	3	3	2	7	2	4
Tranquilizers	36	31	10	11	17	18	8	7
Stimulants ³	30	26	8	9	18	13	6	6
Methamphetamine ³	19	17	7	5	10	2	1	3
Sedatives	11	11	5	5	4	4	2	2
ILLCIT DRUGS OTHER THAN MARIJUANA¹	76	70	23	19	44	42	20	19

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.19D Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.78	0.79	1.27	1.28	0.58	0.62	0.28	0.26
Marijuana and Hashish	0.80	0.78	1.17	1.22	0.53	0.58	0.20	0.24
Cocaine	0.37	0.36	0.28	0.25	0.16	0.13	0.02	0.03
Crack	0.11	0.10	0.12	0.19	0.04	0.01	*	*
Heroin	0.10	0.10	0.21	0.18	0.01	0.03	*	*
Hallucinogens	0.36	0.31	0.31	0.29	0.18	0.13	0.06	0.05
LSD	0.10	0.11	0.02	0.14	0.06	0.04	*	0.04
PCP	0.06	0.03	*	0.17	*	*	0.02	0.03
Ecstasy	0.30	0.23	0.20	0.21	0.14	0.07	*	0.03
Inhalants	0.14	0.15	0.21	0.03	0.07	0.07	0.06	0.08
Nonmedical Use of Psychotherapeutics ^{2,3}	0.56	0.56	0.74	0.70	0.36	0.33	0.18	0.15
Pain Relievers	0.49	0.51	0.64	0.58	0.30	0.28	0.15	0.14
OxyContin [®]	0.16	0.17	0.13	0.13	0.02	0.07	0.02	0.04
Tranquilizers	0.34	0.33	0.39	0.43	0.17	0.17	0.08	0.06
Stimulants ³	0.29	0.27	0.29	0.35	0.17	0.12	0.06	0.06
Methamphetamine ³	0.19	0.18	0.25	0.20	0.10	0.02	0.01	0.03
Sedatives	0.11	0.12	0.19	0.22	0.04	0.04	0.02	0.02
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.64	0.63	0.85	0.74	0.40	0.37	0.20	0.18

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.19P Types of Illicit Drug Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.3663		0.8296		0.1407		0.6187	
Marijuana and Hashish	0.3050		0.7037		0.0486		0.9056	
Cocaine	0.1487		0.9819		0.0180		0.9547	
Crack	0.6315		0.4461		0.0428		*	
Heroin	0.7143		0.8991		0.4785		*	
Hallucinogens	0.3842		0.7749		0.2744		0.8043	
LSD	0.4391		0.4113		0.5089		*	
PCP	0.3809		*		*		0.7560	
Ecstasy	0.1399		0.9152		0.0490		*	
Inhalants	0.9743		0.2047		0.5138		0.7074	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.4712		0.8226		0.4191		0.8877	
Pain Relievers	0.5110		0.4426		0.6063		0.8637	
OxyContin [®]	0.9662		0.7213		0.0190		0.0756	
Tranquilizers	0.2157		0.9621		0.7429		0.7502	
Stimulants ³	0.2471		0.9798		0.0513		0.9494	
Methamphetamine ³	0.2110		0.3150		0.0478		0.6129	
Sedatives	0.1584		0.6782		0.9517		0.5538	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.3242		0.7420		0.2923		0.9358	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.20C Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	265	262	129	120	204	226	126	142
Marijuana and Hashish	229	219	102	92	173	182	70	91
Cocaine	126	115	36	51	55	46	29	24
Crack	92	60	23	46	32	21	10	4
Heroin	70	30	18	17	19	*	*	*
Hallucinogens	30	48	7	23	30	29	27	15
LSD	14	30	5	*	13	*	*	*
PCP	*	8	*	6	4	*	*	*
Ecstasy	19	39	3	5	20	26	23	10
Inhalants	50	30	14	2	36	43	16	26
Nonmedical Use of Psychotherapeutics ^{2,3}	141	136	78	81	121	136	99	115
Pain Relievers	120	119	70	71	104	106	93	104
OxyContin [®]	18	25	17	40	17	3	*	16
Tranquilizers	66	65	27	46	64	72	44	51
Stimulants ³	74	64	36	47	49	28	10	24
Methamphetamine ³	62	60	32	47	33	17	10	7
Sedatives	42	24	16	14	34	53	36	28
ILLCIT DRUGS OTHER THAN MARIJUANA¹	178	178	84	97	131	157	107	118

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.20D Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLICIT DRUGS¹	0.82	0.80	0.74	0.70	0.27	0.30	0.18	0.20
Marijuana and Hashish	0.72	0.70	0.59	0.54	0.24	0.24	0.10	0.13
Cocaine	0.42	0.38	0.21	0.31	0.08	0.06	0.04	0.03
Crack	0.31	0.20	0.13	0.28	0.04	0.03	0.01	0.01
Heroin	0.24	0.10	0.11	0.11	0.03	*	*	*
Hallucinogens	0.10	0.16	0.04	0.14	0.04	0.04	0.04	0.02
LSD	0.05	0.10	0.03	*	0.02	*	*	*
PCP	*	0.03	*	0.04	0.00	*	*	*
Ecstasy	0.06	0.13	0.02	0.03	0.03	0.04	0.03	0.01
Inhalants	0.17	0.10	0.08	0.01	0.05	0.06	0.02	0.04
Nonmedical Use of Psychotherapeutics ^{2,3}	0.47	0.44	0.45	0.48	0.17	0.18	0.14	0.16
Pain Relievers	0.40	0.39	0.41	0.42	0.14	0.14	0.13	0.15
OxyContin [®]	0.06	0.09	0.10	0.24	0.02	0.00	*	0.02
Tranquilizers	0.22	0.22	0.16	0.28	0.09	0.10	0.06	0.07
Stimulants ³	0.25	0.22	0.21	0.28	0.07	0.04	0.01	0.03
Methamphetamine ³	0.21	0.20	0.19	0.28	0.05	0.02	0.01	0.01
Sedatives	0.14	0.08	0.09	0.09	0.05	0.07	0.05	0.04
ILLICIT DRUGS OTHER THAN MARIJUANA¹	0.58	0.57	0.49	0.57	0.18	0.21	0.15	0.17

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.20P Types of Illicit Drug Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette and Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Drug	CIGARETTE AND ALCOHOL USE IN PAST MONTH							
	Cigarettes and Alcohol		Cigarettes, But Not Alcohol		Alcohol, But Not Cigarettes		Neither Alcohol Nor Cigarettes	
	2006	2007	2006	2007	2006	2007	2006	2007
ILLCIT DRUGS¹	0.6537		0.9300		0.1130		0.5110	
Marijuana and Hashish	0.4723		0.8268		0.1644		0.5628	
Cocaine	0.2761		0.0244		0.1700		0.4428	
Crack	0.1939		0.0352		0.2760		0.6015	
Heroin	0.1190		0.2834		*		*	
Hallucinogens	0.1100		0.2563		0.5659		0.6608	
LSD	0.4382		*		*		*	
PCP	*		*		*		*	
Ecstasy	0.0552		0.6092		0.7764		0.7301	
Inhalants	0.2329		0.3891		0.5833		0.7808	
Nonmedical Use of Psychotherapeutics ^{2,3}	0.6139		0.9738		0.6276		0.3517	
Pain Relievers	0.7238		0.8259		0.6375		0.7087	
OxyContin [®]	0.1964		0.5156		0.1528		*	
Tranquilizers	0.9007		0.7535		0.7027		0.8067	
Stimulants ³	0.3277		0.9614		0.0373		0.2598	
Methamphetamine ³	0.5094		0.7181		0.0581		0.4815	
Sedatives	0.3358		0.7260		0.8479		0.9035	
ILLCIT DRUGS OTHER THAN MARIJUANA¹	0.6280		0.3772		0.5445		0.5945	

*Low precision; no estimate reported.

¹ Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. Illicit Drugs Other Than Marijuana include cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically. The estimates for Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine incorporated in these summary estimates do not include data from the methamphetamine items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

² Nonmedical use of prescription-type psychotherapeutics includes the nonmedical use of pain relievers, tranquilizers, stimulants, or sedatives and does not include over-the-counter drugs.

³ Estimates of Nonmedical Use of Psychotherapeutics, Stimulants, and Methamphetamine in the designated rows include data from methamphetamine items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.21C Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
TOTAL	426	445	67	68	122	127	376	394
Heavy Alcohol Use ¹	208	219	26	27	88	90	183	191
Binge Alcohol Use But Not Heavy Use ¹	227	232	37	37	84	91	205	206
Alcohol Use But Not Binge Use ¹	209	203	25	24	51	46	197	196
No Alcohol Use	199	195	46	51	48	45	182	184
PAST MONTH CIGARETTE USE	328	336	48	46	124	122	292	294
Heavy Alcohol Use ¹	175	175	22	25	74	75	155	151
Binge Alcohol Use But Not Heavy Use ¹	180	187	28	28	71	68	159	167
Alcohol Use But Not Binge Use ¹	148	127	17	15	41	37	140	120
No Alcohol Use	141	124	26	23	37	37	129	120
NO PAST MONTH CIGARETTE USE	264	295	50	53	76	88	243	272
Heavy Alcohol Use ¹	105	129	14	10	40	40	94	122
Binge Alcohol Use But Not Heavy Use ¹	138	128	23	23	42	55	128	116
Alcohol Use But Not Binge Use ¹	147	155	18	18	31	31	139	151
No Alcohol Use	136	153	37	42	28	27	126	142

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.21D Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: Standard Errors of Percentages, 2006 and 2007

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
TOTAL	0.17	0.18	0.27	0.27	0.37	0.39	0.20	0.21
Heavy Alcohol Use ¹	1.06	1.03	2.78	2.85	1.10	1.17	1.43	1.38
Binge Alcohol Use But Not Heavy Use ¹	0.53	0.53	1.51	1.53	0.73	0.78	0.67	0.66
Alcohol Use But Not Binge Use ¹	0.30	0.29	1.34	1.40	0.72	0.68	0.31	0.31
No Alcohol Use	0.16	0.16	0.21	0.23	0.36	0.34	0.21	0.21
PAST MONTH CIGARETTE USE	0.50	0.52	1.29	1.39	0.69	0.69	0.60	0.60
Heavy Alcohol Use ¹	1.44	1.39	3.19	3.33	1.27	1.40	2.17	1.96
Binge Alcohol Use But Not Heavy Use ¹	0.95	0.97	2.26	2.43	1.14	1.13	1.26	1.26
Alcohol Use But Not Binge Use ¹	0.99	0.93	3.27	4.06	1.63	1.60	1.12	1.05
No Alcohol Use	0.67	0.61	2.04	1.99	1.27	1.28	0.74	0.70
NO PAST MONTH CIGARETTE USE	0.14	0.15	0.21	0.22	0.34	0.38	0.17	0.18
Heavy Alcohol Use ¹	1.37	1.59	4.67	4.95	1.97	1.85	1.66	2.07
Binge Alcohol Use But Not Heavy Use ¹	0.60	0.53	1.82	1.89	0.88	1.01	0.72	0.63
Alcohol Use But Not Binge Use ¹	0.27	0.27	1.30	1.35	0.66	0.66	0.28	0.29
No Alcohol Use	0.13	0.15	0.18	0.20	0.28	0.26	0.18	0.20

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.21P Illicit Drug Use in the Past Month, by Levels of Past Month Cigarette Use and Past Month Alcohol Use and Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Level of Cigarette Use/ Alcohol Use	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
TOTAL	0.2782		0.5464		0.9222		0.3307	
Heavy Alcohol Use ¹	0.3725		0.5274		0.2356		0.2538	
Binge Alcohol Use But Not Heavy Use ¹	0.9399		0.9770		0.5520		0.8831	
Alcohol Use But Not Binge Use ¹	0.0266		0.0728		0.7157		0.0496	
No Alcohol Use	0.7389		0.5828		0.3641		0.6845	
PAST MONTH CIGARETTE USE	0.7216		0.8100		0.5023		0.8262	
Heavy Alcohol Use ¹	0.4277		0.9672		0.0702		0.2592	
Binge Alcohol Use But Not Heavy Use ¹	0.3521		0.9831		0.8542		0.1909	
Alcohol Use But Not Binge Use ¹	0.0644		0.4098		0.6969		0.1005	
No Alcohol Use	0.9742		0.8040		0.8296		0.9300	
NO PAST MONTH CIGARETTE USE	0.6263		0.8669		0.2646		0.3557	
Heavy Alcohol Use ¹	0.6988		0.9818		0.7424		0.6323	
Binge Alcohol Use But Not Heavy Use ¹	0.4299		0.8373		0.1438		0.1779	
Alcohol Use But Not Binge Use ¹	0.3970		0.3500		0.5621		0.3711	
No Alcohol Use	0.4224		0.2942		0.6187		0.5110	

*Low precision; no estimate reported.

NOTE: Illicit Drugs include marijuana/hashish, cocaine (including crack), heroin, hallucinogens, inhalants, or prescription-type psychotherapeutics used nonmedically, based on data from original questions not including methamphetamine items added in 2005 and 2006.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.