

Table 7.22C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	356	364
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	143	147	248	246
Cigars	256	252	251	251
Pipe Tobacco	115	124	121	117
ALCOHOL	722	721	1,223	1,355
Binge Alcohol Use ²	550	558	588	658
Heavy Alcohol Use ²	293	317	295	300

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.22D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	0.19	0.19
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.23	0.24	0.13	0.13
Cigars	0.38	0.39	0.13	0.13
Pipe Tobacco	0.18	0.20	0.07	0.06
ALCOHOL	0.69	0.73	0.45	0.48
Binge Alcohol Use ²	0.66	0.68	0.29	0.32
Heavy Alcohol Use ²	0.44	0.48	0.16	0.16

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.22P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A		0.1851	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.3247		0.2430	
Cigars	0.5621		0.6150	
Pipe Tobacco	0.5276		0.3426	
ALCOHOL	0.6042		0.5964	
Binge Alcohol Use ²	0.1403		0.5539	
Heavy Alcohol Use ²	0.5281		0.8752	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.23C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	33	37
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	21	24	24	25
Cigars	34	36	24	27
Pipe Tobacco	14	17	12	8
ALCOHOL	55	57	70	75
Binge Alcohol Use ²	50	51	51	52
Heavy Alcohol Use ²	27	31	22	21

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.23D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: Standard Errors of Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	0.14	0.16
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.75	0.90	0.10	0.11
Cigars	1.13	1.23	0.10	0.12
Pipe Tobacco	0.51	0.67	0.05	0.04
ALCOHOL	1.31	1.34	0.29	0.30
Binge Alcohol Use ²	1.37	1.37	0.22	0.22
Heavy Alcohol Use ²	0.94	1.11	0.09	0.09

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.23P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 12 to 17, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A		0.9083	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.1513		0.8153	
Cigars	0.1457		0.9508	
Pipe Tobacco	0.1350		0.4554	
ALCOHOL	0.7613		0.3261	
Binge Alcohol Use ²	0.5396		0.0791	
Heavy Alcohol Use ²	0.0802		0.0770	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.24C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	72	76
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	54	55	37	38
Cigars	87	86	63	65
Pipe Tobacco	29	27	16	19
ALCOHOL	195	193	237	245
Binge Alcohol Use ²	172	173	157	170
Heavy Alcohol Use ²	109	104	74	74

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.24D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: Standard Errors of Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	0.33	0.34
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.40	0.42	0.18	0.18
Cigars	0.57	0.58	0.29	0.30
Pipe Tobacco	0.23	0.22	0.08	0.09
ALCOHOL	0.62	0.64	0.66	0.67
Binge Alcohol Use ²	0.73	0.77	0.60	0.58
Heavy Alcohol Use ²	0.68	0.68	0.34	0.32

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.24P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 18 to 25, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A		0.9319	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.2563		0.7504	
Cigars	0.6723		0.7369	
Pipe Tobacco	0.4991		0.8749	
ALCOHOL	0.7437		0.9957	
Binge Alcohol Use ²	0.8668		0.5793	
Heavy Alcohol Use ²	0.2990		0.8895	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.25C Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	347	347
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	127	132	242	238
Cigars	236	226	241	233
Pipe Tobacco	110	120	120	115
ALCOHOL	673	660	1,173	1,306
Binge Alcohol Use ²	501	504	556	619
Heavy Alcohol Use ²	270	293	279	289

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.25D Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: Standard Errors of Percentages, 2006 and 2007

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A	N/A	0.24	0.23
Cigarettes	N/A	N/A	N/A	N/A
Smokeless Tobacco	0.27	0.29	0.17	0.16
Cigars	0.47	0.47	0.16	0.16
Pipe Tobacco	0.23	0.26	0.08	0.08
ALCOHOL	0.88	0.93	0.56	0.60
Binge Alcohol Use ²	0.83	0.84	0.36	0.40
Heavy Alcohol Use ²	0.54	0.59	0.19	0.20

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.25P Tobacco Product and Alcohol Use in the Past Month among Persons Aged 26 or Older, by Past Month Cigarette Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	CIGARETTE USE IN PAST MONTH			
	Any Use		No Use	
	2006	2007	2006	2007
TOBACCO PRODUCTS¹	N/A		0.1585	
Cigarettes	N/A		N/A	
Smokeless Tobacco	0.5446		0.2527	
Cigars	0.4623		0.6167	
Pipe Tobacco	0.5127		0.3541	
ALCOHOL	0.4652		0.6244	
Binge Alcohol Use ²	0.0946		0.5866	
Heavy Alcohol Use ²	0.3212		0.9448	

*Low precision; no estimate reported.

N/A: Not applicable.

¹ Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

² Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.