

Table 7.26C Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	321	346	482	478	484	459	565	614
Cigarettes	293	317	439	436	429	405	525	569
Smokeless Tobacco	113	121	144	133	133	124	185	176
Cigars	151	148	196	205	207	193	153	163
Pipe Tobacco	69	78	77	60	84	87	103	109

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.26D Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	1.15	1.23	0.81	0.85	0.61	0.59	0.43	0.44
Cigarettes	1.19	1.24	0.79	0.84	0.56	0.54	0.41	0.42
Smokeless Tobacco	0.64	0.65	0.36	0.32	0.19	0.18	0.15	0.14
Cigars	0.81	0.78	0.46	0.48	0.29	0.27	0.13	0.13
Pipe Tobacco	0.40	0.45	0.19	0.15	0.12	0.13	0.08	0.09

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.26P Tobacco Product Use in the Past Month among Persons Aged 12 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS ²	0.7408		0.2522		0.1035		0.1209	
Cigarettes	0.9023		0.5109		0.0746		0.2088	
Smokeless Tobacco	0.3147		0.9877		0.8298		0.1318	
Cigars	0.3069		0.2734		0.3915		0.6348	
Pipe Tobacco	0.4910		0.4773		0.9471		0.3219	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.27C Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	29	33	45	44	28	27	50	52
Cigarettes	27	31	42	40	24	24	44	42
Smokeless Tobacco	15	15	16	19	11	10	20	25
Cigars	20	21	26	26	17	15	23	29
Pipe Tobacco	10	10	11	10	6	7	9	11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.27D Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	2.46	2.71	1.54	1.65	1.43	1.51	0.23	0.24
Cigarettes	2.82	2.99	1.54	1.59	1.31	1.34	0.20	0.19
Smokeless Tobacco	2.31	2.27	0.78	0.95	0.70	0.60	0.09	0.11
Cigars	2.66	2.79	1.22	1.31	0.98	0.89	0.11	0.13
Pipe Tobacco	1.64	1.64	0.55	0.51	0.35	0.42	0.04	0.05

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.27P Tobacco Product Use in the Past Month among Persons Aged 12 to 17, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS ²	0.0371		0.9949		0.0555		0.6482	
Cigarettes	0.0499		0.6687		0.0650		0.4113	
Smokeless Tobacco	0.8979		0.3988		0.6956		0.4660	
Cigars	0.8181		0.4707		0.1534		0.1025	
Pipe Tobacco	0.9027		0.7902		0.8355		0.3750	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

80403

Table 7.28C Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	123	115	127	123	82	79	92	90
Cigarettes	109	104	119	114	75	74	86	85
Smokeless Tobacco	45	47	37	37	18	16	22	24
Cigars	66	62	58	65	38	38	40	43
Pipe Tobacco	20	18	19	22	10	11	14	11

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.28D Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	0.94	1.02	0.89	0.90	1.03	1.05	0.67	0.63
Cigarettes	1.03	1.10	0.88	0.92	0.99	1.01	0.63	0.60
Smokeless Tobacco	0.76	0.81	0.42	0.41	0.27	0.25	0.17	0.19
Cigars	0.98	1.04	0.61	0.65	0.56	0.57	0.31	0.33
Pipe Tobacco	0.37	0.37	0.22	0.24	0.16	0.17	0.11	0.09

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.28P Tobacco Product Use in the Past Month among Persons Aged 18 to 25, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS ²	0.1439		0.2147		0.0731		0.1279	
Cigarettes	0.0629		0.0691		0.1142		0.1048	
Smokeless Tobacco	0.5026		0.6771		0.0721		0.3537	
Cigars	0.5426		0.3723		0.4074		0.5174	
Pipe Tobacco	0.2643		0.7024		0.6512		0.2452	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.29C Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Numbers in Thousands, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	295	316	452	447	481	447	549	597
Cigarettes	270	293	407	406	426	391	512	555
Smokeless Tobacco	101	109	135	126	131	123	182	171
Cigars	136	129	185	187	203	188	146	154
Pipe Tobacco	65	75	75	56	82	86	102	107

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.29D Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: Standard Errors of Percentages, 2006 and 2007

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS²	1.64	1.74	1.07	1.10	0.68	0.64	0.58	0.60
Cigarettes	1.69	1.74	1.03	1.08	0.62	0.58	0.55	0.57
Smokeless Tobacco	0.86	0.88	0.46	0.41	0.21	0.20	0.21	0.19
Cigars	1.13	1.02	0.60	0.59	0.32	0.30	0.17	0.17
Pipe Tobacco	0.57	0.64	0.26	0.18	0.13	0.14	0.12	0.12

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.29P Tobacco Product Use in the Past Month among Persons Aged 26 or Older, by Levels of Past Month Alcohol Use: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Substance	LEVEL OF ALCOHOL USE IN PAST MONTH							
	Heavy Use ¹		Binge Use But Not Heavy Use ¹		Use But Not Binge Use ¹		No Use	
	2006	2007	2006	2007	2006	2007	2006	2007
TOBACCO PRODUCTS ²	0.9973		0.3915		0.2284		0.1855	
Cigarettes	0.7307		0.8662		0.1619		0.3270	
Smokeless Tobacco	0.3240		0.8712		0.9515		0.0800	
Cigars	0.5001		0.1465		0.5682		0.7099	
Pipe Tobacco	0.6798		0.3846		0.8894		0.3395	

*Low precision; no estimate reported.

¹ Binge Alcohol Use is defined as drinking five or more drinks on the same occasion (i.e., at the same time or within a couple of hours of each other) on at least 1 day in the past 30 days. Heavy Alcohol Use is defined as drinking five or more drinks on the same occasion on each of 5 or more days in the past 30 days; all heavy alcohol users are also binge alcohol users.

² Tobacco Products include cigarettes, smokeless tobacco (i.e., chewing tobacco or snuff), cigars, or pipe tobacco.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.