

**Table 7.45C Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, 2006 and 2007**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	150	139	25	22	38	36	142	134
From More Than One Doctor	32	57	10	6	12	12	27	55
Wrote Fake Prescription	5	11	2	1	4	5	*	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	19	10	4	5	3	6	19	7
From Friend or Relative for Free	215	226	35	36	80	81	194	201
Bought from Friend or Relative	89	75	15	14	35	32	80	66
Took from Friend or Relative without Asking	61	55	16	15	26	22	53	48
Bought from Drug Dealer or Other Stranger	54	59	11	12	22	20	47	56
Bought on the Internet	5	22	1	2	3	5	4	21
Some Other Way	60	48	13	14	21	23	54	39

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.45D Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.13	1.08	1.50	1.42	0.90	0.91	1.92	1.84
From More Than One Doctor	0.27	0.49	0.68	0.40	0.32	0.32	0.42	0.85
Wrote Fake Prescription	0.04	0.10	0.12	0.07	0.12	0.12	*	0.16
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.08	0.24	0.33	0.07	0.15	0.30	0.10
From Friend or Relative for Free	1.27	1.25	1.73	1.78	1.32	1.31	2.18	2.07
Bought from Friend or Relative	0.73	0.64	0.95	0.96	0.86	0.82	1.22	1.01
Took from Friend or Relative without Asking	0.52	0.47	0.97	1.01	0.65	0.57	0.82	0.74
Bought from Drug Dealer or Other Stranger	0.46	0.52	0.70	0.83	0.57	0.53	0.74	0.88
Bought on the Internet	0.04	0.19	0.04	0.17	0.07	0.13	0.07	0.33
Some Other Way	0.50	0.41	0.85	1.00	0.53	0.60	0.84	0.60

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 7.45P Source Where Pain Relievers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007 Versus 2006**

Source for Most Recent Nonmedical Use among Past Year Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.5513		0.3548		0.9759		0.6283	
From More Than One Doctor	0.0987		0.2205		0.4676		0.0259	
Wrote Fake Prescription	0.1950		0.7461		0.9272		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2968		0.8754		0.1387		0.1031	
From Friend or Relative for Free	0.6552		0.7569		0.6845		0.8420	
Bought from Friend or Relative	0.6584		0.6602		0.5275		0.7887	
Took from Friend or Relative without Asking	0.7591		0.5889		0.6639		0.8354	
Bought from Drug Dealer or Other Stranger	0.8144		0.4671		0.9112		0.8566	
Bought on the Internet	0.0664		0.2877		0.3381		0.1237	
Some Other Way	0.3367		0.8304		0.6063		0.2260	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.46C Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	188	201	23	25	61	63	176	188
From More Than One Doctor	39	28	6	6	10	10	37	25
Wrote Fake Prescription	7	4	*	1	4	4	6	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	10	27	4	5	3	3	9	26
From Friend or Relative for Free	49	49	11	12	21	17	43	44
Bought from Friend or Relative	28	38	8	6	16	18	23	32
Took from Friend or Relative without Asking	13	16	10	7	8	6	*	12
Bought from Drug Dealer or Other Stranger	22	24	5	7	12	13	18	19
Bought on the Internet	*	3	*	1	*	3	*	*
Some Other Way	23	11	7	7	8	4	20	7

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.46D Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.29	1.37	2.63	3.02	1.53	1.55	1.95	1.97
From More Than One Doctor	0.65	0.49	0.99	1.15	0.54	0.57	1.09	0.74
Wrote Fake Prescription	0.12	0.07	*	0.13	0.19	0.21	0.18	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.17	0.46	0.62	0.84	0.14	0.18	0.27	0.76
From Friend or Relative for Free	0.83	0.83	1.78	2.05	1.05	0.93	1.28	1.27
Bought from Friend or Relative	0.48	0.64	1.36	1.18	0.81	0.95	0.69	0.92
Took from Friend or Relative without Asking	0.22	0.27	1.64	1.32	0.43	0.36	*	0.35
Bought from Drug Dealer or Other Stranger	0.38	0.41	0.81	1.33	0.63	0.68	0.54	0.55
Bought on the Internet	*	0.05	*	0.14	*	0.16	*	*
Some Other Way	0.39	0.19	1.20	1.31	0.41	0.23	0.61	0.20

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.46P Source Where Friend or Relative Obtained Pain Relievers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Pain Relievers from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.8824		0.3383		0.8876		0.7846	
From More Than One Doctor	0.5858		0.7276		0.7596		0.4621	
Wrote Fake Prescription	0.6686		*		0.6879		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.3138		0.8056		0.7882		0.3399	
From Friend or Relative for Free	0.5237		0.6693		0.6512		0.5875	
Bought from Friend or Relative	0.1804		0.9369		0.2397		0.3222	
Took from Friend or Relative without Asking	0.8497		0.6006		0.2825		*	
Bought from Drug Dealer or Other Stranger	0.6320		0.0670		0.6573		0.8314	
Bought on the Internet	*		*		*		*	
Some Other Way	0.0519		0.8491		0.0528		0.1916	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.47C Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	115	114	16	17	27	28	110	111
From More Than One Doctor	44	55	8	5	11	14	42	53
Wrote Fake Prescription	16	12	2	1	3	6	*	10
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	22	12	3	5	6	6	21	9
From Friend or Relative for Free	142	141	23	22	48	50	132	130
Bought from Friend or Relative	80	77	16	14	37	33	67	70
Took from Friend or Relative without Asking	47	61	12	13	23	22	38	56
Bought from Drug Dealer or Other Stranger	66	54	10	11	23	22	60	50
Bought on the Internet	11	22	1	4	4	6	10	20
Some Other Way	46	40	12	11	14	18	43	34

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.47D Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.96	1.97	2.45	2.63	1.57	1.75	3.27	3.21
From More Than One Doctor	0.91	1.13	1.39	0.84	0.70	0.98	1.54	1.85
Wrote Fake Prescription	0.34	0.25	0.38	0.24	0.23	0.40	*	0.37
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.47	0.26	0.49	0.91	0.37	0.43	0.79	0.34
From Friend or Relative for Free	2.05	2.20	2.92	2.91	1.93	2.15	3.42	3.53
Bought from Friend or Relative	1.55	1.57	2.43	2.31	1.93	2.03	2.36	2.37
Took from Friend or Relative without Asking	0.97	1.25	1.99	2.11	1.42	1.46	1.39	1.96
Bought from Drug Dealer or Other Stranger	1.30	1.12	1.73	1.83	1.41	1.48	2.09	1.75
Bought on the Internet	0.22	0.46	0.21	0.69	0.25	0.44	0.37	0.74
Some Other Way	0.95	0.85	2.13	1.83	0.95	1.28	1.58	1.24

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.47P Sources Where Pain Relievers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Past Month Nonmedical Use among Past Month Pain Reliever Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.5223		0.8932		0.6501		0.6587	
From More Than One Doctor	0.1185		0.4006		0.3670		0.1377	
Wrote Fake Prescription	0.7776		0.7626		0.3612		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6320		0.2011		0.5761		0.3088	
From Friend or Relative for Free	0.7440		0.5425		0.8120		0.7805	
Bought from Friend or Relative	0.9449		0.9642		0.7370		0.6572	
Took from Friend or Relative without Asking	0.3878		0.9146		0.7673		0.2312	
Bought from Drug Dealer or Other Stranger	0.5649		0.9552		0.9366		0.5898	
Bought on the Internet	0.2332		0.3095		0.2318		0.4812	
Some Other Way	0.3253		0.2991		0.6991		0.3440	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained pain relievers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.48C Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	126	117	17	14	37	41	120	108
From More Than One Doctor	47	36	7	7	12	16	45	31
Wrote Fake Prescription	25	15	2	*	0	6	25	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	25	27	*	4	4	6	24	*
From Friend or Relative for Free	44	55	10	9	20	18	37	52
Bought from Friend or Relative	33	42	9	7	17	22	26	35
Took from Friend or Relative without Asking	22	13	10	7	10	9	16	7
Bought from Drug Dealer or Other Stranger	29	23	6	8	14	16	23	15
Bought on the Internet	7	11	*	2	4	7	*	8
Some Other Way	15	14	7	4	9	6	10	12

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.48D Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.84	2.40	4.40	4.14	2.16	2.45	2.63	3.74
From More Than One Doctor	1.78	1.45	2.53	2.84	1.34	1.91	2.96	2.16
Wrote Fake Prescription	0.99	0.62	0.98	*	0.02	0.72	1.71	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.97	1.13	*	1.88	0.45	0.76	1.67	*
From Friend or Relative for Free	1.62	2.13	3.57	3.58	2.11	2.18	2.45	3.42
Bought from Friend or Relative	1.28	1.65	3.22	2.97	1.81	2.51	1.78	2.44
Took from Friend or Relative without Asking	0.87	0.55	3.59	2.98	1.19	1.10	1.13	0.50
Bought from Drug Dealer or Other Stranger	1.11	0.98	2.19	3.25	1.60	1.98	1.58	1.07
Bought on the Internet	0.28	0.46	*	0.78	0.51	0.87	*	0.61
Some Other Way	0.60	0.63	2.89	1.65	1.06	0.85	0.67	0.90

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.48P Sources Where Friend or Relative Obtained Pain Relievers among Past Month Users Aged 12 or Older Who Obtained Pain Relievers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Pain Relievers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.1607		0.8475		0.9849		0.0797	
From More Than One Doctor	0.8712		0.2864		0.1336		0.3691	
Wrote Fake Prescription	0.4459		*		0.0215		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7296		*		0.2052		*	
From Friend or Relative for Free	0.4118		0.7176		0.7520		0.2612	
Bought from Friend or Relative	0.0505		0.5998		0.1500		0.0918	
Took from Friend or Relative without Asking	0.1418		0.4137		0.6995		0.2371	
Bought from Drug Dealer or Other Stranger	0.6594		0.0680		0.4180		0.5640	
Bought on the Internet	0.1078		*		0.2303		*	
Some Other Way	0.5256		0.1131		0.3264		0.5330	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained pain relievers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.49C Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	63	72	7	7	19	16	60	70
From More Than One Doctor	16	22	4	0	7	6	14	21
Wrote Fake Prescription	18	5	2	3	2	4	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	17	7	2	1	2	7	*	*
From Friend or Relative for Free	162	162	18	19	48	47	153	155
Bought from Friend or Relative	49	64	12	11	30	28	35	57
Took from Friend or Relative without Asking	24	77	6	8	13	16	19	74
Bought from Drug Dealer or Other Stranger	52	29	11	9	22	19	46	19
Bought on the Internet	9	23	*	2	5	3	7	23
Some Other Way	35	41	6	6	10	16	33	37

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.49D Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.30	1.38	1.87	1.70	1.09	0.98	2.15	2.21
From More Than One Doctor	0.35	0.45	0.97	0.13	0.41	0.39	0.53	0.71
Wrote Fake Prescription	0.39	0.11	0.48	0.82	0.11	0.26	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.37	0.14	0.52	0.37	0.13	0.42	*	*
From Friend or Relative for Free	1.98	2.11	3.47	3.35	1.99	1.98	3.06	3.36
Bought from Friend or Relative	1.00	1.25	2.69	2.55	1.53	1.58	1.28	1.83
Took from Friend or Relative without Asking	0.51	1.50	1.67	2.02	0.77	0.98	0.70	2.42
Bought from Drug Dealer or Other Stranger	1.08	0.59	2.72	2.25	1.23	1.13	1.68	0.65
Bought on the Internet	0.19	0.48	*	0.47	0.28	0.21	0.28	0.78
Some Other Way	0.74	0.81	1.44	1.54	0.62	1.01	1.23	1.21

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 7.49P Source Where Tranquilizers Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007 Versus 2006**

Source for Most Recent Nonmedical Use among Past Year Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.5177		0.8881		0.2760		0.6352	
From More Than One Doctor	0.8993		0.0540		0.9662		0.6922	
Wrote Fake Prescription	0.4525		0.5838		0.5845		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5213		0.7660		0.3180		*	
From Friend or Relative for Free	0.7633		0.8109		0.9406		0.6182	
Bought from Friend or Relative	0.2776		0.8032		0.8303		0.1461	
Took from Friend or Relative without Asking	0.0380		0.3192		0.1731		0.0822	
Bought from Drug Dealer or Other Stranger	0.0101		0.4215		0.4453		0.0341	
Bought on the Internet	0.2792		*		0.3959		0.1958	
Some Other Way	0.9140		0.3034		0.8694		0.9093	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.50C Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	149	138	11	10	37	34	144	134
From More Than One Doctor	16	16	*	3	8	7	14	14
Wrote Fake Prescription	0	2	*	*	0	2	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	8	7	*	*	1	3	*	6
From Friend or Relative for Free	28	51	8	7	17	17	21	47
Bought from Friend or Relative	30	24	*	6	11	14	28	18
Took from Friend or Relative without Asking	6	15	2	*	6	4	*	*
Bought from Drug Dealer or Other Stranger	15	24	4	5	7	7	12	23
Bought on the Internet	14	3	*	*	4	*	*	1
Some Other Way	5	25	*	1	4	8	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.50D Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.90	2.35	5.88	5.58	2.59	2.80	2.38	3.17
From More Than One Doctor	0.64	0.61	*	2.19	1.07	0.95	0.82	0.78
Wrote Fake Prescription	0.01	0.06	*	*	0.02	0.22	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.30	0.25	*	*	0.16	0.43	*	0.32
From Friend or Relative for Free	1.09	1.81	4.73	4.66	2.05	2.11	1.24	2.53
Bought from Friend or Relative	1.13	0.91	*	3.85	1.38	1.82	1.60	1.02
Took from Friend or Relative without Asking	0.23	0.56	1.53	*	0.72	0.59	*	*
Bought from Drug Dealer or Other Stranger	0.56	0.88	2.94	3.42	0.90	0.90	0.72	1.23
Bought on the Internet	0.52	0.13	*	*	0.50	*	*	0.06
Some Other Way	0.20	0.92	*	0.59	0.53	1.10	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.50P Source Where Friend or Relative Obtained Tranquilizers among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Tranquilizers from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.1152		0.3942		0.1818		0.2107	
From More Than One Doctor	0.8502		*		0.6881		0.9100	
Wrote Fake Prescription	0.3646		*		0.3614		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9478		*		0.3926		*	
From Friend or Relative for Free	0.1336		0.2892		0.4873		0.0970	
Bought from Friend or Relative	0.5036		*		0.2330		0.1805	
Took from Friend or Relative without Asking	0.3743		*		0.2013		*	
Bought from Drug Dealer or Other Stranger	0.2103		0.4026		0.8053		0.2610	
Bought on the Internet	0.2631		*		*		*	
Some Other Way	0.1141		*		0.1323		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.51C Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	42	50	*	5	13	10	39	49
From More Than One Doctor	13	17	*	*	5	3	*	*
Wrote Fake Prescription	*	4	*	*	2	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	5	*	0	1	5	*	*
From Friend or Relative for Free	83	97	*	*	26	27	78	92
Bought from Friend or Relative	28	47	*	*	21	19	17	*
Took from Friend or Relative without Asking	16	42	*	4	11	13	11	*
Bought from Drug Dealer or Other Stranger	44	43	*	5	15	13	*	*
Bought on the Internet	14	19	*	*	4	4	*	*
Some Other Way	19	21	*	2	7	8	17	19

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.51D Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	2.41	2.74	*	3.58	2.01	1.88	3.99	4.25
From More Than One Doctor	0.82	1.01	*	*	0.88	0.51	*	*
Wrote Fake Prescription	*	0.24	*	*	0.30	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.33	*	0.18	0.15	1.04	*	*
From Friend or Relative for Free	3.51	3.37	*	*	3.23	3.31	5.62	5.26
Bought from Friend or Relative	1.76	2.65	*	*	2.82	3.11	1.90	*
Took from Friend or Relative without Asking	1.02	2.40	*	3.01	1.83	2.31	1.15	*
Bought from Drug Dealer or Other Stranger	2.51	2.40	*	4.05	2.24	2.29	*	*
Bought on the Internet	0.84	1.12	*	*	0.64	0.75	*	*
Some Other Way	1.17	1.22	*	1.77	1.20	1.47	1.83	1.83

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.51P Sources Where Tranquilizers Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Past Month Nonmedical Use among Past Month Tranquilizer Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.7164		*		0.3811		0.6034	
From More Than One Doctor	0.9023		*		0.4516		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		0.1999		*	
From Friend or Relative for Free	0.3221		*		0.1511		0.7265	
Bought from Friend or Relative	0.5484		*		0.8925		*	
Took from Friend or Relative without Asking	0.1299		*		0.4242		*	
Bought from Drug Dealer or Other Stranger	0.3437		*		0.2022		*	
Bought on the Internet	0.5674		*		0.7081		*	
Some Other Way	0.5517		*		0.8938		0.4584	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained tranquilizers for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.52C Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	74	74	*	*	21	21	*	*
From More Than One Doctor	13	*	*	*	5	7	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	2	3	*	*	2	*	*	*
From Friend or Relative for Free	21	20	*	*	8	13	*	*
Bought from Friend or Relative	20	14	*	*	6	9	*	*
Took from Friend or Relative without Asking	6	9	*	*	4	*	*	*
Bought from Drug Dealer or Other Stranger	7	18	*	*	5	7	*	*
Bought on the Internet	*	0	*	*	3	0	*	*
Some Other Way	3	5	*	*	*	2	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.52D Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	3.70	4.25	*	*	3.80	4.56	*	*
From More Than One Doctor	1.55	*	*	*	1.75	2.49	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.30	0.30	*	*	0.98	*	*	*
From Friend or Relative for Free	2.50	2.30	*	*	3.15	4.30	*	*
Bought from Friend or Relative	2.41	1.65	*	*	2.27	3.28	*	*
Took from Friend or Relative without Asking	0.68	1.05	*	*	1.51	*	*	*
Bought from Drug Dealer or Other Stranger	0.90	1.89	*	*	2.01	2.67	*	*
Bought on the Internet	*	0.03	*	*	1.33	0.12	*	*
Some Other Way	0.37	0.61	*	*	*	0.91	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.52P Sources Where Friend or Relative Obtained Tranquilizers among Past Month Users Aged 12 or Older Who Obtained Tranquilizers Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Tranquilizers ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.4141		*		0.2059		*	
From More Than One Doctor	*		*		0.2523		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6765		*		*		*	
From Friend or Relative for Free	0.7805		*		0.0864		*	
Bought from Friend or Relative	0.9319		*		0.5433		*	
Took from Friend or Relative without Asking	0.6363		*		*		*	
Bought from Drug Dealer or Other Stranger	0.0889		*		0.4273		*	
Bought on the Internet	*		*		0.1251		*	
Some Other Way	0.4002		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained tranquilizers; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.53C Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	38	41	11	6	16	11	32	39
From More Than One Doctor	9	5	2	3	2	4	8	*
Wrote Fake Prescription	14	4	2	*	*	3	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	5	10	4	1	3	6	*	8
From Friend or Relative for Free	103	93	16	16	41	36	92	82
Bought from Friend or Relative	27	36	8	9	23	22	12	26
Took from Friend or Relative without Asking	31	19	8	7	12	11	28	15
Bought from Drug Dealer or Other Stranger	22	22	7	6	12	12	17	19
Bought on the Internet	24	25	3	1	4	4	23	25
Some Other Way	36	14	6	4	6	4	*	12

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.53D Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	1.58	1.94	2.96	2.17	1.66	1.29	3.08	4.06
From More Than One Doctor	0.39	0.25	0.64	0.88	0.26	0.52	0.84	*
Wrote Fake Prescription	0.61	0.19	0.55	*	*	0.42	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.24	0.50	1.23	0.44	0.32	0.68	*	0.92
From Friend or Relative for Free	2.64	2.88	3.43	3.79	2.59	2.59	5.39	5.76
Bought from Friend or Relative	1.22	1.69	2.18	2.78	2.03	2.22	1.22	2.96
Took from Friend or Relative without Asking	1.34	0.95	2.09	2.31	1.20	1.21	2.78	1.66
Bought from Drug Dealer or Other Stranger	0.95	1.14	1.93	2.02	1.22	1.39	1.70	2.18
Bought on the Internet	1.00	1.26	0.85	0.26	0.42	0.54	2.24	2.81
Some Other Way	1.52	0.70	1.76	1.51	0.64	0.53	*	1.40

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 7.53P Source Where Stimulants Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007 Versus 2006**

Source for Most Recent Nonmedical Use among Past Year Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.9220		0.1297		0.2910		0.3738	
From More Than One Doctor	0.4923		0.6933		0.3122		*	
Wrote Fake Prescription	0.4827		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2620		0.0434		0.1131		*	
From Friend or Relative for Free	0.7210		0.4074		0.4067		0.7953	
Bought from Friend or Relative	0.0233		0.0920		0.4624		0.0268	
Took from Friend or Relative without Asking	0.1982		0.8320		0.6314		0.1947	
Bought from Drug Dealer or Other Stranger	0.2609		0.8114		0.2464		0.4795	
Bought on the Internet	0.9013		0.2079		0.4097		0.9692	
Some Other Way	0.0558		0.6602		0.1663		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.54C Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	74	63	11	*	33	30	*	*
From More Than One Doctor	8	10	4	*	6	5	*	*
Wrote Fake Prescription	*	0	*	0	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	3	2	*	*	*	*	*	*
From Friend or Relative for Free	18	41	5	4	11	11	*	*
Bought from Friend or Relative	*	18	2	*	7	8	*	*
Took from Friend or Relative without Asking	5	4	*	*	3	1	*	*
Bought from Drug Dealer or Other Stranger	5	5	2	3	4	3	*	*
Bought on the Internet	7	8	*	*	4	*	*	*
Some Other Way	1	7	*	1	*	1	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.54D Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	4.07	4.11	5.52	*	2.87	3.34	*	*
From More Than One Doctor	0.69	1.03	2.88	*	1.23	1.29	*	*
Wrote Fake Prescription	*	0.02	*	0.13	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.23	0.19	*	*	*	*	*	*
From Friend or Relative for Free	1.72	3.92	3.71	3.45	2.16	2.48	*	*
Bought from Friend or Relative	*	1.88	1.56	*	1.41	1.85	*	*
Took from Friend or Relative without Asking	0.48	0.45	*	*	0.66	0.23	*	*
Bought from Drug Dealer or Other Stranger	0.45	0.49	1.39	2.68	0.82	0.78	*	*
Bought on the Internet	0.63	0.87	*	*	0.75	*	*	*
Some Other Way	0.07	0.71	*	0.61	*	0.20	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.54P Source Where Friend or Relative Obtained Stimulants among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Stimulants from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.6621		*		0.8091		*	
From More Than One Doctor	0.3562		*		0.9327		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.9802		*		*		*	
From Friend or Relative for Free	0.8307		0.2572		0.9449		*	
Bought from Friend or Relative	*		*		0.4594		*	
Took from Friend or Relative without Asking	0.8044		*		0.3780		*	
Bought from Drug Dealer or Other Stranger	0.7957		0.3106		0.9088		*	
Bought on the Internet	0.8566		*		*		*	
Some Other Way	0.1096		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Year Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.55C Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	26	25	*	*	10	9	*	*
From More Than One Doctor	7	8	*	*	2	*	*	*
Wrote Fake Prescription	2	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	4	8	*	*	*	*	*	*
From Friend or Relative for Free	44	62	*	*	23	17	*	*
Bought from Friend or Relative	18	24	*	*	16	18	*	*
Took from Friend or Relative without Asking	27	17	*	*	11	*	*	*
Bought from Drug Dealer or Other Stranger	19	11	*	*	13	8	*	*
Bought on the Internet	18	19	*	*	4	*	*	*
Some Other Way	*	6	*	*	4	1	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.55D Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	3.20	3.51	*	*	2.97	3.39	*	*
From More Than One Doctor	1.00	1.24	*	*	0.83	*	*	*
Wrote Fake Prescription	0.26	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.61	1.26	*	*	*	*	*	*
From Friend or Relative for Free	4.43	5.29	*	*	4.52	4.62	*	*
Bought from Friend or Relative	2.55	3.55	*	*	4.33	5.46	*	*
Took from Friend or Relative without Asking	3.36	2.44	*	*	3.39	*	*	*
Bought from Drug Dealer or Other Stranger	2.42	1.73	*	*	3.86	2.99	*	*
Bought on the Internet	2.35	2.70	*	*	1.18	*	*	*
Some Other Way	*	0.90	*	*	1.36	0.56	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.55P Sources Where Stimulants Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Past Month Nonmedical Use among Past Month Stimulant Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	0.7147		*		0.8447		*	
From More Than One Doctor	0.6728		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.5581		*		*		*	
From Friend or Relative for Free	0.3646		*		0.4410		*	
Bought from Friend or Relative	0.1486		*		0.6605		*	
Took from Friend or Relative without Asking	0.2156		*		*		*	
Bought from Drug Dealer or Other Stranger	0.6490		*		0.9827		*	
Bought on the Internet	0.9270		*		*		*	
Some Other Way	*		*		0.1468		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained stimulants for past month nonmedical use; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.56C Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	37	*	*	*	21	*	*	*
From More Than One Doctor	4	5	*	*	2	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	20	*	*	*	*	*	*	*
Bought from Friend or Relative	10	*	*	*	7	*	*	*
Took from Friend or Relative without Asking	3	3	*	*	3	*	*	*
Bought from Drug Dealer or Other Stranger	*	8	*	*	6	*	*	*
Bought on the Internet	4	*	*	*	*	*	*	*
Some Other Way	2	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.56D Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	5.49	*	*	*	5.05	*	*	*
From More Than One Doctor	1.20	1.59	*	*	1.24	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	5.10	*	*	*	*	*	*	*
Bought from Friend or Relative	2.81	*	*	*	4.31	*	*	*
Took from Friend or Relative without Asking	1.04	0.99	*	*	1.89	*	*	*
Bought from Drug Dealer or Other Stranger	*	2.57	*	*	3.20	*	*	*
Bought on the Internet	1.04	*	*	*	*	*	*	*
Some Other Way	0.47	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.56P Sources Where Friend or Relative Obtained Stimulants among Past Month Users Aged 12 or Older Who Obtained Stimulants Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Stimulants ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*		*		*		*	
From More Than One Doctor	0.9645		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.4183		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained stimulants; thus, these response categories are not mutually exclusive.

NOTE: Estimates in this table do not include data from methamphetamine use items added in 2005 and 2006.

¹ Past Month Stimulant Use refers only to nonmedical stimulant use and does not include those reporting only methamphetamine use. Additionally, respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.57C Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	91	60	13	*	31	22	87	*
Bought from Friend or Relative	56	57	8	*	17	15	53	54
Took from Friend or Relative without Asking	11	10	4	*	5	6	*	8
Bought from Drug Dealer or Other Stranger	56	52	*	*	16	14	53	*
Bought on the Internet	*	3	*	*	1	*	*	*
Some Other Way	5	17	2	*	5	6	1	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.57D Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	3.47	3.93	5.97	*	4.29	4.13	5.34	*
Bought from Friend or Relative	2.94	3.72	4.76	*	3.20	3.46	4.51	5.80
Took from Friend or Relative without Asking	0.67	0.83	2.71	*	0.92	1.58	*	1.04
Bought from Drug Dealer or Other Stranger	2.87	3.82	*	*	3.04	3.51	4.38	*
Bought on the Internet	*	0.26	*	*	0.21	*	*	*
Some Other Way	0.33	1.43	1.49	*	0.97	1.54	0.13	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.57P Source Where Methamphetamine was Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Most Recent Nonmedical Use among Past Year Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	0.0859		*		0.4670		*	
Bought from Friend or Relative	0.0535		*		0.6357		0.0523	
Took from Friend or Relative without Asking	0.6452		*		0.7441		*	
Bought from Drug Dealer or Other Stranger	0.7794		*		0.9400		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.1542		*		0.2881		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.58C Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	27	*	*	*	13	*	*	*
Bought from Friend or Relative	53	37	*	*	16	*	*	*
Took from Friend or Relative without Asking	9	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	52	*	*	*	*	*	*	*
Bought on the Internet	0	*	*	*	*	*	*	*
Some Other Way	7	8	*	*	5	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.58D Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	3.23	*	*	*	4.30	*	*	*
Bought from Friend or Relative	5.01	5.74	*	*	5.76	*	*	*
Took from Friend or Relative without Asking	1.14	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	5.28	*	*	*	*	*	*	*
Bought on the Internet	0.04	*	*	*	*	*	*	*
Some Other Way	0.94	1.75	*	*	2.07	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.58P Source Where Friend or Relative Obtained Methamphetamine among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Methamphetamine from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	0.9776		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.4560		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Estimates for Source Where Friend or Relative Obtained Methamphetamine include (a) past year users who reported obtaining this drug for their most recent use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Methamphetamine for respondent's most recent use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.59C Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	44	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	1	*	*	*	*	*	*	*
Some Other Way	8	8	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.59D Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	5.61	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	0.78	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	0.23	*	*	*	*	*	*	*
Some Other Way	1.25	1.90	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.59P Sources Where Methamphetamine Was Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: P Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Past Month Nonmedical Use among Past Month Methamphetamine Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	0.4049		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports. See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which they obtained methamphetamine for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.60C Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports.

See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.60D Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.60	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports.

See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.60P Sources Where Friend or Relative Obtained Methamphetamine among Past Month Users Aged 12 or Older Who Obtained Methamphetamine Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Methamphetamine ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates in this table include data from methamphetamine use items added in 2005 and 2006 and are not comparable with estimates presented in prior NSDUH reports.

See Section B.4.6 in Appendix B of the *Results from the 2007 National Survey on Drug Use and Health: National Findings*.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained methamphetamine; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 7.61C Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
Standard Errors of Numbers in Thousands, 2006 and 2007**

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	54	*	*	*	*	5	*	*
From More Than One Doctor	7	5	*	*	4	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	1	7	*	*	*	*	*	*
From Friend or Relative for Free	67	86	*	*	*	14	*	*
Bought from Friend or Relative	*	9	*	*	*	7	*	*
Took from Friend or Relative without Asking	9	7	*	*	6	*	*	*
Bought from Drug Dealer or Other Stranger	6	17	*	*	4	4	*	*
Bought on the Internet	4	5	*	*	*	*	*	*
Some Other Way	9	11	*	*	*	2	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.61D Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	5.40	*	*	*	*	3.23	*	*
From More Than One Doctor	0.86	0.64	*	*	2.14	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.12	0.92	*	*	*	*	*	*
From Friend or Relative for Free	5.79	5.73	*	*	*	5.70	*	*
Bought from Friend or Relative	*	1.12	*	*	*	4.27	*	*
Took from Friend or Relative without Asking	1.13	0.94	*	*	2.99	*	*	*
Bought from Drug Dealer or Other Stranger	0.69	2.08	*	*	2.18	2.40	*	*
Bought on the Internet	0.50	0.59	*	*	*	*	*	*
Some Other Way	1.07	1.41	*	*	*	1.47	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

**Table 7.61P Source Where Sedatives Were Obtained for Most Recent Nonmedical Use among Past Year Users Aged 12 or Older, by Age Group:
P Values from Tests of Differences of Percentages, 2007 Versus 2006**

Source for Most Recent Nonmedical Use among Past Year Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*		*		*		*	
From More Than One Doctor	0.5271		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.2370		*		*		*	
From Friend or Relative for Free	0.0774		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	0.8017		*		*		*	
Bought from Drug Dealer or Other Stranger	0.2385		*		0.8688		*	
Bought on the Internet	0.9386		*		*		*	
Some Other Way	0.5549		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source for Most Recent Nonmedical Use include (a) past month users who reported a single source of obtainment during the past 30 days, (b) past month users who identified their last source of obtainment after reporting multiple sources of obtainment in the past 30 days, and (c) all other past year users who reported their last source of obtainment in the past year.

¹ Respondents with unknown data on Source for Most Recent Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.62C Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	75	*	*	*	*	*	*
From More Than One Doctor	6	7	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0	*	*	*	*	*	*	*
From Friend or Relative for Free	4	5	*	*	*	*	*	*
Bought from Friend or Relative	3	4	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	1	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	3	*	*	*	*	*	*	*
Bought on the Internet	0	0	*	*	*	*	*	*
Some Other Way	*	1	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.62D Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	2.46	*	*	*	*	*	*
From More Than One Doctor	1.41	1.57	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.03	*	*	*	*	*	*	*
From Friend or Relative for Free	1.04	1.26	*	*	*	*	*	*
Bought from Friend or Relative	0.69	0.98	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	0.22	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	0.63	*	*	*	*	*	*	*
Bought on the Internet	0.04	0.06	*	*	*	*	*	*
Some Other Way	*	0.24	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.62P Source Where Friend or Relative Obtained Sedatives among Past Year Users Aged 12 or Older Who Obtained Most Recently Used Sedatives from a Friend or Relative for Free in the Past Year: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*		*		*		*	
From More Than One Doctor	0.9522		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	0.3594		*		*		*	
Bought from Friend or Relative	0.8724		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	0.8309		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Estimates for Source Where Friend or Relative Obtained Drug include (a) past year users who reported obtaining this drug for their most recent nonmedical use from a friend or relative for free and then reported a valid source for where their friend or relative obtained this drug and (b) past month users who reported only obtaining this drug for their past month nonmedical use from a friend or relative for free and then reported a single valid source for where their friend or relative obtained this drug.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's most recent use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.63C Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.63D Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.63P Sources Where Sedatives Were Obtained for Nonmedical Use in the Past Month among Past Month Users Aged 12 or Older, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source for Past Month Nonmedical Use among Past Month Sedative Users ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which they obtained sedatives for past month nonmedical use; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown data on Source for Past Month Nonmedical Use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.64C Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Numbers in Thousands, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.64D Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: Standard Errors of Percentages, 2006 and 2007

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*	*	*	*	*	*	*	*
From More Than One Doctor	*	*	*	*	*	*	*	*
Wrote Fake Prescription	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*
Bought from Friend or Relative	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	*	*	*	*	*	*	*	*
Bought from Drug Dealer or Other Stranger	*	*	*	*	*	*	*	*
Bought on the Internet	*	*	*	*	*	*	*	*
Some Other Way	*	*	*	*	*	*	*	*

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.

Table 7.64P Sources Where Friend or Relative Obtained Sedatives among Past Month Users Aged 12 or Older Who Obtained Sedatives Used in the Past Month from a Friend or Relative for Free, by Age Group: *P* Values from Tests of Differences of Percentages, 2007 Versus 2006

Source Where Friend or Relative Obtained Sedatives ¹	Total		AGE GROUP					
			12-17		18-25		26 or Older	
	2006	2007	2006	2007	2006	2007	2006	2007
From One Doctor	*		*		*		*	
From More Than One Doctor	*		*		*		*	
Wrote Fake Prescription	*		*		*		*	
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*		*		*		*	
From Friend or Relative for Free	*		*		*		*	
Bought from Friend or Relative	*		*		*		*	
Took from Friend or Relative without Asking	*		*		*		*	
Bought from Drug Dealer or Other Stranger	*		*		*		*	
Bought on the Internet	*		*		*		*	
Some Other Way	*		*		*		*	

*Low precision; no estimate reported.

NOTE: Respondents could indicate multiple sources from which a friend or relative obtained sedatives; thus, these response categories are not mutually exclusive.

¹ Respondents with unknown or invalid data on Source Where Friend or Relative Obtained Drug for respondent's past month use and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

Source: SAMHSA, Office of Applied Studies, National Survey on Drug Use and Health, 2006 and 2007.